

Neem cake-urea mixed applications increase growth in paddy

R. S. Bhalla^{1,*} and K. V. Devi Prasad²

¹Foundation for Ecological Research, Advocacy and Learning, No. 27, 2nd Cross, Appavou Nagar, Vazhakulam, Puducherry 605 012, India

²Sustainable Development Programme, Department of Ecology and Environmental Sciences, Pondicherry University, Kalapet, Puducherry 600 014, India

Significant increase in the growth of paddy plant parts was achieved by halving the urea used and pelleting the remaining with neem cake prior to application. Results on a non-averaged dataset showed significant increase in leaf length, number of leaves, number of panicles, number of tillers and greenness of leaves. Averaged data showed similar results barring the number of panicles, which were not significantly different from the control. The results point to a higher availability of nitrogen in the treatment, even though half the amount of urea was applied as opposed to the control. This could be attributed to inhibition of de-nitrifying bacteria by neem as well as a slower continuous release of nitrogen when urea is pelleted with neem than when it is applied directly. The study makes a strong case for cutting down on nitrogen application in paddy using low-cost, readily available materials, without compromising on the yield.

Keywords: Neem cake, nitrogen application, paddy, urea.

VARYING efficiencies of nitrogen absorption and differential rates of uptake of plants based on growth stages of crops have been known for decades¹⁻³ and have resulted in a number of formulations for slow release of fertilizers and modifications of cultivation methods⁴⁻⁶. However, adoption of these practices has been low among farmers who continue to rely on split dosages and urea as the primary source of nitrogen. In the case of paddy, nitrogen use efficiency is estimated to be only 30–40% of the application⁷. The loss of nitrogen through run-off, ammonia emissions, leaching and de-nitrification is a serious environmental issue^{8,9}. Environmental impacts of nitrogenous fertilizers include contamination of local groundwater resources and rivers^{6,7,10,11} and emission of nitrous oxide which is both a greenhouse gas as well as a major cause of ozone depletion¹²⁻¹⁴. Slow-release nitrogenous fertilizers typically cost more than conventional formulations and have had less acceptance among farmers. Low-cost alternatives to slow-release fertilizers include addition of plant-derived organic substances such as neem cake, neem leaf¹⁵ and *Pongamia* extract¹⁶. Many of these also inhibit de-nitrifying bacterial activity. Other plant-derived products such as waste from processed medicinal com-

pounds^{17,18} and tannins from forest litter¹⁹ also inhibit denitrification.

A farmer's field-based experiment-cum-demonstration was conducted in two villages, Thuruvai and Rayapud-dupakkam, in Puducherry and Tamil Nadu respectively, in 2002–03. Selected fields had clayey soil and the same variety of paddy (IR20). A random paired-block design was used with each field containing both the test and control blocks. Blocks were sub-divided into four 5 × 5 m plots. Each test block was separated from the rest of the field by a bund and linked directly to the irrigation source to prevent inflow of water and dissolved nutrient from the rest of the field. A total of 21 test and control block pairs were set up in this manner. Urea was applied in the control block at the rate of 120 kg N/ha in a split dosage with two-thirds at the time of transplanting and the rest 25 days later. The total amount of nitrogen applied was halved for the test block (60 kg N/ha) and applied in the same proportion after blending with neem seed cake in the proportion of 3 : 1 and drying two days in the shade.

We recorded relative growth of parts of the paddy plant. Fifteen plants were chosen randomly from each of the four plots per block, i.e. 60 samples each from a block-pair. Observations included leaf length, panicle length, leaf colour (using the IRRI colour card), number of leaves, number of panicles, and number of tillers. Box plots show data both as averaged reading (Figure 1) as well as disaggregated (Figure 2) for the listed variables. Yield measurements for these experiments were not possible as they interfered with harvesting operations and farmers were unwilling to allow harvesting of the experimental plots separately.

The data were analysed with the R package²⁰ using the ESS²¹ and R-Commander interfaces²². Analysis with and without averaging the values per block of samples was done. As most parameters were not normally distributed, a Bartlett and Levene's test for homogeneity of variance was performed to see if the data were suitable for analysis of variance. Both tests showed that most of the variables could not be compared using parametric tests barring the number of tillers, which had an acceptable *P* value in Bartlett's test (Tables 1 and 2).

Pairwise *t*-test (done for comparison sake though the data were not suitable for parametric tests) and Wilcoxon test were done (Tables 3 and 4). The Wilcoxon test (on paired non-averaged data) showed strong differences on all growth variables other than panicle length. On averaged data neither panicle length nor number of tillers was significantly different, while differences between all the other variables were significant. A Kruskal–Wallis rank sum test showed that all variables were significantly different with non-averaged data, while only average leaf length was strongly significant (Table 5). Average numbers of leaves and panicles were significant, while average numbers of panicle length, leaf colour and number of tillers were not significantly different. A correlation ma-

*For correspondence. (e-mail: bhalla@feralindia.org)




Figure 1. Box plots of data averaged per block.

Table 1. Tests for homogeneity of variance on non-averaged data

| Variable | Bartlett's K-squared | P value |
|---|----------------------|---------|
| Bartlett test of homogeneity of variance (non-averaged data) | | |
| Leaf length | 2.11530 | 0.14580 |
| Panicle length | 1.49220 | 0.22190 |
| Leaf colour | 0.52370 | 0.46930 |
| Number of leaves | 0.27560 | 0.59960 |
| Number of panicles | 1.56880 | 0.21040 |
| Number of tillers | 4.02330 | 0.04488 |
| Variable | F value | Pr (>F) |
| Levene's test for homogeneity of variance (non-averaged data) | | |
| Leaf length | 1.76840 | 0.18380 |
| Panicle length | 2.71120 | 0.09975 |
| Leaf colour | 2.17970 | 0.14150 |
| Number of leaves | 0.15760 | 0.69150 |
| Number of panicles | 1.70780 | 0.19270 |
| Number of tillers | 1.85490 | 0.17470 |

Table 2. Tests for homogeneity of variance on averaged data

| Variable | Bartlett's K-squared | P value |
|---|----------------------|---------|
| Bartlett test of homogeneity of variance (averaged data) | | |
| Average leaf length | 0.00810 | 0.92810 |
| Average panicle length | 0.50720 | 0.47640 |
| Average leaf colour | 0.52370 | 0.46930 |
| Average number of leaves | 2.62090 | 0.10550 |
| Average number of panicles | 1.56880 | 0.21040 |
| Average number of tillers | 4.02330 | 0.04488 |
| Variable | F value | Pr (>F) |
| Levene's test for homogeneity of variance (averaged data) | | |
| Average leaf length | 0.00280 | 0.95780 |
| Average panicle length | 1.83660 | 0.17680 |
| Average leaf colour | 2.17970 | 0.14150 |
| Average number of leaves | 1.69780 | 0.19400 |
| Average number of panicles | 1.70780 | 0.19270 |
| Average number of tillers | 1.85490 | 0.17470 |




Figure 2. Box plots of disaggregated data.

Table 3. Wilcoxon pairwise comparison on non-averaged data

| Variable | Wilcoxon test | | |
|--------------------|---------------|-----------|----------------------------------|
| | W value | P value | Evidence against null hypothesis |
| Leaf length | 373716 | < 2.2e-16 | Very strong |
| Panicle length | 94037 | 0.3663 | Little or none |
| Leaf colour | 11798.5 | 8.265e-16 | Very strong |
| Number of leaves | 5113.5 | 1.153e-07 | Very strong |
| Number of panicles | 54076 | 2.159e-14 | Very strong |
| Number of tillers | 54610 | 6.013e-14 | Very strong |

trix within the growth factors between control and test plots was run (Table 6). A significant correlation between leaf length (a proxy for leaf area, i.e. photosynthetic area)

and other growth variables was expected. However, this was true only for leaf length and panicle length. A strong correlation between the number of tillers and number of panicles was also found.

Two processes probably contributed to the increased nitrogen availability – inhibition of de-nitrifying bacteria by neem and the slow release of nitrogen caused by the binding of urea to the surface of neem cake⁶. As the experiments were conducted on farmer’s fields, farming practices between different plots could not be standardized. It should be noted however that comparisons were paired and within the same block, therefore, inter-field variations did not affect the analysis. Thus inter-field differences in soil nitrogen and irrigation water would not have influenced the results. Nitrogen provided by the neem

Table 4. Pairwise comparison on averaged data

| Variable | t value | P value | Evidence against null hypothesis |
|----------------------------|----------|-----------|----------------------------------|
| <i>T</i> -test | | | |
| Average leaf length | 4.19600 | 0.0004448 | Very strong |
| Average panicle length | 1.04970 | 0.30640 | Little or none |
| Average leaf colour | 2.31140 | 0.03361 | Fairly strong |
| Average number of leaves | 3.56980 | 0.001918 | Very strong |
| Average number of panicles | 2.00540 | 0.05864 | Suggestive |
| Average number of tillers | 1.52730 | 0.14230 | Little or none |
| Variable | W value | P value | Evidence against null hypothesis |
| Willcoxon test | | | |
| Average leaf length | 20.00000 | 0.0003538 | Very strong |
| Average panicle length | 91.00000 | 0.41200 | Little or none |
| Average leaf colour | 13.00000 | 0.04529 | Fairly strong |
| Average number of leaves | 28.50000 | 0.002631 | Very strong |
| Average number of panicles | 55.00000 | 0.03701 | Fairly strong |
| Average number of tillers | 65.50000 | 0.14510 | Little or none |

Table 5. Kruskal–Wallis test on non-averaged and averaged data

| Variable | K–W chi-squared | P value | Evidence against null hypothesis |
|---|-----------------|-----------|----------------------------------|
| Kruskal–Wallis rank sum test with $df = 1$ (test/control) and non-averaged data | | | |
| Leaf length | 55.657 | 8.629e-14 | Very strong |
| Panicle length | 6.4831 | 0.010890 | Very strong |
| Leaf colour | 8.5643 | 0.003428 | Very strong |
| Number of leaves | 15.9192 | 6.61e-05 | Very strong |
| Number of panicles | 7.0263 | 0.008032 | Very strong |
| Number of tillers | 6.9874 | 0.008209 | Very strong |
| Kruskal–Wallis rank sum test with $df = 1$ (test/control) and averaged data | | | |
| Average leaf length | 3.7034 | 0.0543 | Fairly strong |
| Average panicle length | 0.0837 | 0.7724 | Little or none |
| Average leaf colour | 2.5061 | 0.1134 | Little or none |
| Average number of leaves | 2.7666 | 0.09625 | Suggestive |
| Average number of panicles | 3.017 | 0.0824 | Suggestive |
| Average number of tillers | 2.0232 | 0.1549 | Little or none |

Table 6. Correlation matrix

| | Average leaf length | Average panicle length | Leaf colour | Number of panicles | Number of tillers | Number of leaves |
|------------------------|---------------------|------------------------|-------------|--------------------|-------------------|------------------|
| Control plots | | | | | | |
| Average leaf length | 1 | 0.51 | 0.2 | -0.13 | -0.2 | 0.13 |
| Average panicle length | 0.509 | 1 | 0.04 | -0.41 | -0.44 | -0.11 |
| Leaf colour | 0.199 | 0.04 | 1 | 0.08 | 0.21 | 0.24 |
| Number of panicles | -0.13 | -0.41 | 0.08 | 1 | 0.86 | 0.06 |
| Number of tillers | -0.2 | -0.44 | 0.21 | 0.86 | 1 | 0.19 |
| Number of leaves | 0.13 | -0.11 | 0.24 | 0.06 | 0.19 | 1 |
| Treatment plots | | | | | | |
| Average leaf length | 1 | 0.62 | 0.22 | 0.03 | 0.05 | -0.01 |
| Average panicle length | 0.62 | 1 | 0.12 | -0.15 | -0.07 | 0.09 |
| Leaf colour | 0.22 | 0.12 | 1 | 0.07 | 0.13 | 0.16 |
| Number of panicles | 0.03 | -0.15 | 0.07 | 1 | 0.9 | -0.16 |
| Number of tillers | 0.05 | -0.07 | 0.13 | 0.9 | 1 | -0.11 |
| Number of leaves | -0.01 | 0.09 | 0.16 | -0.16 | -0.11 | 1 |

cake itself would be insignificant given that the weight of neem cake used was a third of the urea and the nitrogen

percentage in neem cake is very small (2–5%)²³ compared to urea (46)²⁴. Efforts by individual farmers may

have differential impact on yield. However, it is not possible to study this effect under standardized conditions and therefore, further analysis along this line was not pursued.

Our study showed that addition of neem cake was an economical and acceptable way of reducing fertilizer application. However, it also showed higher growth of vegetative than reproductive tissue, although both were higher than the control. Better controlled conditions would allow determination of the impact of such treatments on paddy yield. Also, an additional control where only neem cake is applied, would have allowed us to further clarify its role.

- Mehrotra, O. N., Sinha, N. S. and Srivastava, R. D. L., Studies on the nutrition of indian cereals. ii. Uptake of nitrogen and phosphorus at different stages of growth by early paddy (unhulled rice) under varying fertilizer management. *Plant Soil*, 1968, **XXVIII**, 422–430.
- Ramanathan, K. M. and Krishnamoorthy, K. K., Nutrient uptake by paddy during the main three stages of growth. *Plant Soil*, 1973, **39**, 29–33.
- Srivastava, S. C., An appraisal of nitrogen fertilisation practices for paddy, wheat, sugarcane, and potato. *Plant Soil*, 1970, **33**, 265–271.
- Ryoichi, K., Reduction of effluent nitrogen and phosphorus from paddy fields. *Paddy Water Environ.*, 2003, **1**, 133–138.
- Vlek, P. L. G., Byrnes, B. H. and Craswell, E. T., Effect of urea placement on leaching losses of nitrogen from flooded rice soils. *Plant Soil*, 1980, **54**, 441–449.
- Wakimoto, K., Utilization advantages of controlled release nitrogen fertilizer on paddy rice cultivation. *Jpn Agric. Res. Quart.*, 2004, **38**, 15–20.
- Johri, B. N. and Yadav, J. S. P., Low and declining crop response to fertilizers. National Academy of Agricultural Sciences, Policy Paper 35, 2006.
- Md. Toufic, I., Measurement of ammonia emission following surface application of urea fertilizer from paddy fields. *Pak. J. Biol. Sci.*, 2005, **8**, 429–432.
- Newbould, P., The use of nitrogen fertiliser in agriculture. Where do we go practically and ecologically? *Plant Soil*, 1989, **115**, 297–311.
- Cho, J. Y. and Choi, J. K., Nitrogen and phosphorus losses from a broad paddy field in central Korea. *Commun. Soil Sci. Plant Anal.*, 2001, **32**, 2395–2410.
- Pathak, B. K., Kazama, F. and Toshiaki, I., Monitoring of nitrogen leaching from a tropical paddy in Thailand. *Agric. Eng. Int.: CIGR J. Sci. Res. Dev.*, 2004, **VI**, 1–11.
- Bhatia, A., Pathak, H. and Aggarwal, P. K., Inventory of methane and nitrous oxide emissions from agricultural soils of India and their global warming potential. *Curr. Sci.*, 2004, **87**, 1904–1912.
- Garg, A., Shukla, P. R., Kapshe, M. and Menon, D., Indian methane and nitrous oxide emissions and mitigation flexibility. *Atmos. Environ.*, 2004, **38**, 1965–1977.
- Pathak, H., Li, C. and Wassmann, R., Greenhouse gas emissions from Indian rice fields: Calibration and upscaling using the DNDC model. *Biogeosciences*, 2005, **2**, 113–123.
- Santhi, S. R., Palaniappan, S. P. and Purushothaman, D., Influence of neem leaf on nitrification in a lowland rice soil. *Plant Soil*, 1985, **93**, 133–135.
- Majumdar, D., Suppression of nitrification and N₂O emission by karanjin – a nitrification inhibitor prepared from karanja (*Pongamia glabra* Vent.). *Chemosphere*, 2002, **47**, 845–850.

- Maoa, J., Yangb, L., Shic, Y., Hua, J., Piaoa, Z., Meia, L. and Yina, S., Crude extract of *Astragalus mongholicus* root inhibits crop seed germination and soil nitrifying activity. *Soil Biol. Biochem.*, 2005, **38**, 201–208.
- Rice Elroy, L., Some roles of allelopathic compounds in plant communities. *Biochem. Syst. Ecol.*, 1977, **5**, 201–206.
- Karusa, T. E. C., Zasoskia, R. J., Dahlgrena, R. A., Horwatha, W. R. and Preston, C. M., Carbon and nitrogen dynamics in a forest soil amended with purified tannins from different plant species. *Soil Biol. Biochem.*, 2004, **36**, 309–321.
- R Development Core Team, *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria, 2004; available at <http://www.R-project.org>
- Rossini, A. J., Heiberger, R. M., Hornik, K., Maechler, M., Sparapani, R. A. and Eglén, S. J., ESS-emacs speaks statistics. ESS version 5.3.6, 2006; available at: <http://ess.r-project.org/>
- Fox, J., with contributions from Ash, M. *et al.*, Rcmdr: R-commander. R package version 1.3-5; available at <http://www.r-project.org; http://socserv.socsci.mcmaster.ca/jfox/Misc/Rcmdr/>
- Ghosh and Saha, Effects of inoculation of cyanobacteria on nitrogen status and nutrition of rice (*Oryza sativa* L.) in an entisol amended with chemical and organic sources of nitrogen. *Biol. Fertil. Soils*, 1997, **24**, 123–128; <http://dx.doi.org/10.1007/BF01420232>
- Patnaik, N., *Soil Fertility and Fertilizer Use*, Indian Council of Agricultural Research, New Delhi, 1992, 3rd edn, pp. 203–247.

ACKNOWLEDGEMENTS. This study was funded by the Department of Science and Technology, New Delhi, under the Science and Society Programme. We acknowledge useful comments by an anonymous referee.

Received 17 September 2007; revised accepted 13 March 2008

Antidiabetic activity of *Aloe vera* and histology of organs in streptozotocin-induced diabetic rats

Ayesha Noor¹, S. Gunasekaran²,
A. Soosai Manickam² and M. A. Vijayalakshmi^{1,*}

¹Centre for Bio-Separation Technology, VIT University, Vellore 632 014, India

²Diabetes Research Laboratory, Department of Physiology, Christian Medical College, Vellore 632 002, India

An attempt was made to study the beneficial effects of *Aloe vera* (L.) Burm. fil. in streptozotocin-induced diabetic rats. In diabetic induced rats fed with *A. vera* (300 mg/kg body wt), the fasting plasma glucose levels were reduced to normal and body weight was found to be increased. In the pancreatic sections of diabetic rats fed with *A. vera*, the islets were comparable to normal rats. In liver, the changes caused after induction of diabetes are granular cytoplasm, dilated sinusoids, shrunken nuclei and inflammation, which was re-

*For correspondence. (e-mail: indvijji@yahoo.com)