

TRANS-HIMADRI THRUST AND DOMAL UPWARPS IMMEDIATELY SOUTH OF COLLISION ZONE AND TECTONIC IMPLICATIONS

K. S. VALDIYA

Department of Geology, Kumaun University, Nainital 263 001, India.

ABSTRACT

A deep thrust-fault is recognized north of the Great Himalaya (=Himadri) encompassing the Zaskar, NandaDevi-PanchChuli, Dhaulagiri-Annapurna and Sagarmatha (Everest) ranges. Stretching over 1600 km from Ladakh to Sikkim, the Trans-Himadri Thrust (T-HT) sharply severs the Precambrian basement complex of high-grade metamorphics of the Great Himalayan subprovince from the late Proterozoic-to-Upper Cretaceous/Eocene sedimentaries of the Tethyan domain. It is a lag thrust that has attenuated, truncated and regionally eliminated the lower units of the Tethyan succession including the Cambrian. It has caused spectacular back-folding and back-faulting in the sedimentaries, and mylonitized and considerably sheared the basement metamorphics and granites intruding them. The T-HT developed in the continental margin of the northward advancing Indian plate following the blocking of movement along the zone of collision of the Indian and Asian crustal plates.

The leading edge of the Indian crust between Ladakh and northcentral Nepal exhibits very conspicuous domal upwarps immediately south of the line of collision. These upwarps imply great resistance encountered by the Indian crust and resultant folding up of its front. These anticlinal structures and the attendant breaking up of the crust along the T-HT indicate that the Indian plate was too buoyant to underthrust significantly under the Tibetan (Asian) crust. The fact that not much of the Tethyan sedimentary prism has been lost in the subduction rules out large-scale underthrusting of the Indian plate under the Asian crust.

INTRODUCTION

THE ruggedly youthful Great Himalaya (=Himadri) constituted of Precambrian high-grade metamorphics are succeeded by a very thick succession of Late Proterozoic-to-Upper Cretaceous/Eocene sedimentaries of the Tethyan domain, representing the shelf on the continental margin of the Indian shield. It is widely, if not universally, believed that the Great Himalayan metamorphics transitionally grade upwards into the Tethyan sediments. Metamorphism related to the principal phase of Himalayan orogeny and attendant widespread invasion of mid-Tertiary (28–20 m.y.) anatectic granite ($^{87}\text{Sr}/^{86}\text{Sr} > 0.7478$) coupled with apparent concordance of structures in earlier studied sections are responsible for this





assumption. Investigations by the author in the Kumaun Himalaya and analysis of literature demonstrate that the Great Himalayan lithotectonic subprovince is separated from the Tethyan sediments by a lag thrust of regional dimension extending perhaps discontinuously from the Sindhu valley in Ladakh to northern Sikkim. This thrust, designated as the Trans-Himadri Thrust (T-HT), is geodynamically as important as the Main Central Thrust (MCT), Main Boundary Thrust (MBT) and Himalayan Front Fault (HFF) that subdivide the Himalayan province into five latitudinal subprovinces (figure 1).

FAULT DELIMITING NORTHERN BOUNDARY OF GREAT HIMALAYA

It was in northeastern Kumaun Himalaya in the

Figure 1. The Himalayan Front Fault (HFF), Main Boundary Thrust (MBT), Main Central Thrust (MCT) and Trans-Himadri Thrust (T-HT) subdivide the Himalayan province into four lithotectonic subprovinces. Note the basement upwarps immediately south of the Indus-Tsangpo Suture (ITSZ) – the zone of collision of the Indian and Asian plates.

Figure 2. The high-grade Vaikrita metamorphics intruded by mid-Tertiary granites are sharply truncated by the T-HT against the backfolded Tethyan sedimentaries beginning with upper Proterozoic flysch in northern Kumaun.






Figure 3. The map and section⁶ of the Ladakh area showing faulted contact of the Vaikrita metamorphics of the Zanskar Range against the Tethyan sediments. Note the progressive steepening of the T-HT northwest-wards.




Figure 4a – b. a. In the Suru valley the upper Palaeozoic-Triassic Panjal Volcanics abut against the granite-injected Zaskar (Vaikrita) crystallines⁷ — the whole of Proterozoic and much of Palaeozoic missing; b. In the Spiti valley a thrust of trivial magnitude⁴ separates the Vaikrita from the Haimanta sedimentaries.

upper reaches of the Dhaulī, Gori and Kali rivers (figure 2) where the T-HT was first recognized and described as the Malari Thrust^{1-3,23}. The Malari Thrust has profoundly sheared and mylonitized the metamorphics, migmatites and mid-Tertiary granites of the Vaikrita Group, and considerably attenuated the Late Proterozoic Martoli flysch and Ralam conglomerate in the Dhaulī and Gori valleys and eliminated the upper part of the Budhi Schist (Vaikrita) and wholly the Ralam conglomerate in the Kali valley (figure 2).

To the northwest in the Spiti valley a trivial thrust separates the Vaikrita metamorphics from the Tethyan Haimanta⁴ (figure 4b). The intensity of deformation progressively increases northwestwards. The thrust between the Vaikrita crystallines of the Zaskar and Late Proterozoic Phe flysch (figure 3) dips 10–30° in the Bhaga valley in Lahaul, increasing to 45–75° in the upper Doda valley^{5,6}. In the Suru valley it is the Upper Palaeozoic-to-Triassic Panjal Volcanics that is thrust against the granite-intruded Vaikrita metamorphics⁷—the entire Proterozoic and most of the Palaeozoic having been altogether eliminated

(figure 4a). Further north the T-HT striking almost N-S is abruptly truncated obliquely by the WSW-ENE, E-W, WNW-ESE trending faults of the Indus Tsangpo Suture Zone (ITSZ) (figure 8). Not only the basement rocks and the synclorium of the cover sediments, but also the TsoMorari-Nimaling dome are obliquely cut. There is no suggestion of flexing or bending down of the rock formations, rather the plunging folds and thrusts are sharply cut down and eliminated.

Turning southeast in the Kali Gandaki Valley in west-central Nepal, disharmonic folding of incompetent beds disrupt the continuity of succession⁸ (figure 5) or the strongly lineated (NNE-SSW) Vaikrita metamorphics characterized by complex isoclinal folds give way abruptly to Tethyan sediments without these structures but exhibiting spectacular backfolds (figures 5b,c) and backfaults^{9,10}. Further east in the Sagarmatha (Everest) massif the Chomolungma Thrust¹¹ or Main North Himalayan Thrust¹² separates the Ordovician fossil-bearing limestones from the Great Himalayan hornfelsed pelitic schists intruded by granites. The metamorphics are strongly

mylonitized (figure 6). In the adjoining part of Sikkim, the Trans-Axial Thrust¹³ cuts the Great Himalayan Tongshi-Chekha schists from the Tethyan Everest Pelites-Everest Limestone succession. It is noticed that throughout Kumaun and Nepal the Cambrian sediments are missing in the Tethyan column—obviously eliminated by the T-HT.

UPWARPS AT THE LEADING EDGE OF INDIAN CRUST

The pronounced domal upwarp at the northern edge of the Indian crust in the proximity of the ITSZ (figures 1 and 7) has brought up the basement meta-

morphics against the obducted material in the Gurla Mandhata-Mansarovar area^{14,15}. The upper part of the doubly plunging TsoMorari-Nimaling anticline in Ladakh (figures 7b and 8) is constituted of anatectic granite-injected high-grade Vaikrita metamorphics and metamorphosed sedimentary cover involved in imbricate thrusting^{4,16,17}. The Nanga Parbat massif in northwestern Kashmir possibly represents a tight anticlinal structure caught in the loop of the ITSZ and delimited to the west by a N-S trending tear fault.

To the southeast, in northwestern Nepal, "strong axial rise" has brought up the crystalline basement above the sedimentary cover between Manangobhot




Figure 5a-c. a. In western Nepal the contact of the basement crystallines-sedimentary cover is marked by decollement and disharmonic folding⁸; b. North of Kanjiroba—Dhaulagiri massifs the sedimentary cover is backfolded due to gravity gliding¹⁰; c. A huge dome is developed north of Annapurna⁹.




Figure 6. T-HT separates the high-grade metamorphics of the Sagarmatha (Everest) massif from the Tethyan succession beginning with Ordovician sediments^{11,19}.

and Phijor¹⁸. The anticlinal fold (Nar Dome) north of Annapurna characterized by centripetally overturned folds⁹ (figure 5c) and the anticlinal structure related to the Kangmar Thrust north of Sagarmatha¹⁹ represent basement upwarps (figure 6b).

GEOPHYSICAL CONDITIONS OF THE COLLISION ZONE

The occurrence of 10-15 m.y. old potassic lavas in the Neogene volcanic belt immediately to the north




Figure 7. Sections of the Kailas-Mansarovar^{14,15} (north of Kumaun) and Tso Morari (SE Ladakh)^{16,17} areas showing domal upwarps exposing the basement metamorphics at the leading edge of the Indian plate.

of the ITSZ and of 6–8 m.y. old granite in the Kangmar area about 80 km south of the ITSZ²⁰ coupled with pronounced geothermal activity in the collision zone as manifest in 400 hydrothermal areas²¹ and high heat flow of the order of 125–175 mW/m² imply very shallow origin and young age of the magmatic activity²² (figure 9). The existence of high conductivity layer (parallel resistance 3–5 Ωm) at the depth of 3–14 km south of the ITSZ as defined by magnetotelluric studies²⁰ indicates partial melting in the very shallow part of the upper crust. These facts are consistent with the model of the basement upwarp in the Indian crust against the steeply dipping subduction zone.

CONCLUSIONS

Developed between ITSZ and MCT, the T-HT cuts the 10–20 km thick homoclinal Great Himalayan (Himadri) slab from the sedimentary cover and registers upthrusting of the Precambrian basement complex that initiated gravity gliding and consequent backfolding of the cover (Tethyan) sediments^{2,23}. The T-HT developed in the continental margin of the Indian shield following the blocking of the movement around 45 m.y. ago on the zone of collision (ITSZ).

The basement upwarps of considerable dimension developed at the leading edge of the northward




Figure 8. The map (modified after Thakur¹⁶) shows the ITSZ obliquely truncating the Tethyan synclinorium and the elongate Tso Morari-Nimaling dome of the basement crystallines. All Himalayan structures abruptly end against the ITSZ.

moving crust indicate that the Indian crust buoyed up, and was thus prevented from significant underthrusting beneath the Tibetan plateau. Indeed the

buoyancy resistance was so strong that the back part of the moving crust broke up along the basement-cover contact, giving rise to the T-HT. These fea-




Figure 9a - c. a. On the two sides of the collision zone (ITSZ) there are Neogene-Quaternary volcanoes and young granites²⁷. The inset shows distribution in Tibet of high-temperature geothermal areas in the proximity of the ITSZ²¹. b. High heat flow values between Great Himalaya and southern Tibet^{20,22}. c. The magnetotelluric studies outline a layer of anomalously high conductivity in the shallow part (3 - 14 km depth) of the crust, both to the north and south of the ITSZ²⁰. This layer possibly is the source of high heat flow and magmatic-volcanic activities.

tures together with the fact that very small part of the sedimentary prism of the Tethyan shelf has been lost in the obduction in the ITSZ, repudiate the postulation that a large part of the Indian crust has slid under the Tibetan crust.

This conclusion is consistent with the analysis of studies on the seismic wave velocity structure beneath Tibet implying compression and horizontal shortening²⁴, the seismic surface wave dispersion and attenuation that do not support crustal underthrusting model²⁵, the flexural rigidity of the Indian crust which would not allow entire Himalaya to underthrust Tibet²⁶ and the shortening by at least 40% of early Tertiary molasse of the collision zone and absence of systematically migrating elevated front that would have generated southern-derived molasse sediments²⁷

ACKNOWLEDGEMENT

Financial assistance from DST, New Delhi is gratefully acknowledged.

15 December 1986

1. Valdiya, K. S., *Proc. Geodynamics of Himalayan Region*, NGRI, Hyderabad, 1973, p. 204.
2. Valdiya, K. S., In: *Zagros-Hindukush-Himalaya: Geodynamic Evolution*, (eds) H. K. Gupta and F. M. Delany, Am. Geophys. Union, Washington, 1981, p. 87.
3. Kumar, G., Mehdi, S. H. and Prakash, G., *J. Palaeont. Soc. India*, 1970, 15, 80.
4. Berthelsen, A., *Medd. Dansk. Geol. Foren.*, 1951, 12, 102.
5. Nanda, M. M. and Singh, M. P., *Himalayan Geology*, WIHG, Dehradun, 1977, Vol. 6, 365.
6. Baud, A., Gaetani, M., Gerzanti, E., Fois, E., Nicova, A. and Tintovi, A., *Ecolgae. Geol. Helv.*, 1984, 77, 171.
7. Fuchs, G. R., *Jahrb. Geol. Bundes.*, 1977b, 120, 219.
8. Bordet, P., *Recherches Geologiques dans l' Himalaya du Nepal, Region du Makalu*, CNRS, Paris, 1961, p. 275.
9. Bordet, P., Colchen, M. and Le Fort, P., *Recherches Geologiques dans l' Himalaya du Nepal, Region du Nyi-Shang*, CNRS, Paris, 1975, p. 138.
10. Fuchs, G. R., *Jahrb. Geol. Bundes.*, 1977a, 120, 165.
11. Jaros, J. and Kalvoda, J., *Geological Structure of the Himalaya: Mount Everest-Makalu Section*, Akad. Nakl. Cesko, Praha, 1978, p. 67.
12. Chang Chengfa and Pan Ye-shen, *Geological and Ecological studies of the Qinghai-Xizang Plateau*, Science Press, Beijing, 1981, Vol. 1, p. 11.
13. Sinha Roy, S., *Proc. Sym. Geodynamics of Himalayan Region*, NGRI, Hyderabad, 1973, p. 156.
14. Gansser, A., *Geology of the Himalayas*, Interscience, New York-London, 1964, p. 280.
15. Gansser, A., *Ecolgae. Geol. Helv.*, 1966, 59, 831.
16. Thakur, V. C., *Trans. R. Soc. Edinburgh*, 1981, 72, 89.
17. Thakur, V. C. and Viridi, N. S., *J. Geol. Soc. India*, 1979, 22, 46.
18. Hagen, T., *Jahrb. St. Gallischen Naturf. Gessel.*, 1958, 76.
19. Brunel, M., *Tectonophysics*, 1986, 85, 247.
20. Pham, V. N., *et al.*, *Nature (London)*, 1986, 319, 310.
21. Teng Wei and Zhang Mingtao, *Geological and Ecological Studies of the Qinghai-Xizang Plateau*, Science Press, Beijing, 1981, p. 841.
22. Francheteau, J., *et al.*, *Nature (London)*, 1984, 307, 32.
23. Valdiya, K. S., *Colloq. Tectonics of Asia, Intern. Geol. Congress, Moscow*, 1984, 5, 110.
24. Chen, W. P. and Molnar, P., *J. Geophys. Res.*, 1981, 86, 4183.
25. Gupta, H. K., Rao, V. D. and Singh, J., *Tectonophysics*, 1982, 81, 213.
26. Lyon-Caen and Molnar, P., *J. Geophys. Res.*, 1983, 88, 8171.
27. Zhao Yun-Sheng, Zhang Qi, Jin Jin Cheng-wei and Deng Wan-ming, *Geological and Ecological Studies of the Qinghai-Xizang Plateau*, Science Press, Beijing, 1981, p. 363.