

22. Chauhan, O. S. and Suneethi, J., *Curr. Sci.*, 2001, **81**, 1231–1234.
23. Duplessy, J. C., Delibrias, G., Turon, J. L., Pujol, C. and Duprat, J., *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 1981, **35**, 121–144.
24. Duplessy, J. C., *Nature*, 1993, **364**, 322–324.
25. Sharma, M. and Owen, L. A., *Quat. Sci. Rev.*, 1996, **15**, 335–365.
26. Phadtare, N. R., *Quat. Res.*, 2000, **53**, 122–129.
27. Sharma, C., Chauhan, M. S., Gupta, A. and Rajagopalan, G., in Symposium on 'Recent Advances in Geological Studies of North-west Himalaya and Foredeep', Geological Survey of India, Lucknow, 1995, abstr. vol., p. 90.
28. Sarkar, A., Ramesh, R., Somayajulu, B. L. K., Agnihotri, R., Jull, A. J. T. and Burr, G. S., *Earth Planet. Sci. Lett.*, 2000, **177**, 208–218.
29. Naidu, P. D., *Curr. Sci.*, 1996, **71**, 715–718.

ACKNOWLEDGEMENTS. I thank the Director, National Institute of Oceanography, Goa. Thanks are due to M. Sarnthein, E. Vogelsang and S. Kusumgar for carbon and oxygen isotopic analyses and interpretation. Part of the work was done during the visit of the author to the University of Kiel under INSA exchange programme.

Received 22 July 2002, revised accepted 17 October 2002

Application of fossil cuticles in determining palaeoatmospheric CO₂ concentration

R. C. Mehrotra^{*†}, Rajni Tewari^{*} and Ashutosh Joshi^{**}

^{*}Birbal Sahni Institute of Palaeobotany, 53 University Road, Lucknow 226 007, India

^{**}Geological Survey of India, Lucknow 226 004, India

During the last decade stomatal index analysis from fossil leaf cuticles has emerged as a powerful tool in estimating palaeo CO₂ (pCO₂) levels. In India, this particular method has yet not been applied on fossil leaves. In order to reconstruct pCO₂ concentration of the time of deposition of the fossil, this technique is applied on a fossil leaf of *Terminalia catappa* Linn., family Combretaceae which has recently been described from the Upper Siwalik sediments of West Kameng District, Arunachal Pradesh and is considered as of Upper Plio-Pleistocene in age.

It is necessary to study the past CO₂ concentration in order to understand the future climatic changes. The amount of CO₂ is rapidly increasing in the atmosphere due to industrial revolution and human activities. The rise in temperature is related to the increase in CO₂ level in atmosphere. Thus, this increase in CO₂ is alarming as it is

expected to increase the earth's mean temperature by 2–5°C at the end of this century¹. The increase in temperature is responsible for the melting of glaciers that are ultimately responsible for the rise in sea level. El Nino, the weather phenomenon which caused havoc, displaced many people from their homes and caused about 30,000 deaths in the last decade is also linked with the increase in CO₂ level. Higher CO₂ level causes plants to develop a more extensive root system which allows the plant to exploit more water and nutrients from the soil and therefore, the good effect of the increase in CO₂ concentration is the increase in the productivity of plants due to stimulation in photosynthesis.

Quantifying palaeo CO₂ (pCO₂) is essential in order to understand climatic changes and their effect on global temperature. The instrumental record of CO₂ concentration is available only for the last six decades. Hence, a number of methods have been evolved to reconstruct pCO₂ in the geologic past, each having its own merits, demerits and limitations. The most reliable method for estimation of pCO₂ concentration during Quaternary is the analysis of air trapped in polar ice. However, this method is only applicable for the past 400 ka because of the absence of ice before this time. Though it is difficult to measure pCO₂ concentration directly with the same accuracy before this period, it is possible to measure it from the proxy signals obtained from carbon isotope analysis of palaeosols, peat or marine organic matter; boron isotope analysis of planktonic foraminifera and by modelling the long-term global carbon budget¹. Recently, stomatal index (SI) analysis from fossil leaf cuticles has emerged as a powerful tool in deducing pCO₂ levels^{1–6}. In India, this particular aspect has not yet been touched and we are applying this method on the Indian Pleistocene fossil leaf to reconstruct pCO₂ concentration.

Recently, Joshi *et al.*⁷ described a fossil of *Terminalia palaeocatappa* Awasthi and Mehrotra⁸ leaf along with its cuticle from the Upper Siwalik sediments of West Kameng District, Arunachal Pradesh. The age of the fossil ranges from the Upper Pliocene to Pleistocene. The fossil was considered similar to the extant species *Terminalia catappa* Linn., family Combretaceae because of the close resemblance in the morphological features of the leaf and cuticle.

Woodward⁹ pointed out, on the basis of the herbarium leaf material collected over the past 200 years, that the stomatal frequency of leaves of woody plants is inversely proportional to atmospheric CO₂ concentration. Stomatal frequency is generally expressed either in terms of stomatal density (no. per mm²) or SI¹⁰ [stomatal density/(stomatal density + epidermal cell density)] × 100. It has been proved through the experiments done by various workers that SI is relatively stable under varying climatic conditions and water supply because changes in the rate of leaf expansion do not affect the ratio between stomata and epidermal cells¹. Herbarium leaf material of *T. catappa*

[†]For correspondence. (e-mail: rcmehtrotra@yahoo.com)




Figure 1. Response of average stomatal index of *Terminalia catappa* to CO₂ concentration and estimation of pCO₂ of the fossil leaf.

Table 1. Stomatal index (SI) and pCO₂ data for extant and fossil leaves of *T. catappa*

Extant leaf	SI [#]	pCO ₂ (ppmv)
2002	11.3	370
1992	12.7	357
1960	17.5	317
Fossil leaf	14.5	342 (estimated)

[#]SI = [Stomatal density/(stomatal density + epidermal cell density)] × 100.

collected during 1960, 1992 and 2002 was procured to assess its SI response to CO₂ increase in the atmosphere. Quantitative pCO₂ reconstructions can be made by establishing an SI–CO₂ relationship under known pCO₂ for a species; then by using this ‘calibration set’ it is possible to estimate pCO₂ from the SI of the same species from the fossil record.

A number of readings of SIs of the herbarium leaf material of *T. catappa* were taken into account in order to calculate the average SI. Data are presented in Table 1. A perusal of the table points out that the SI decreases with the increase in CO₂ concentration. A comparison of SI of the fossil (a number of readings were taken from the type material to calculate average SI) with the historical responsiveness of that of the modern leaf clearly indicates that pCO₂ concentration during Upper Plio-Pleistocene time should not be greater than the present value of 370 ppmv. The average SI data presented in Table 1 were plotted on a graph (Figure 1) in order to infer a quantitative estimation of atmospheric CO₂ of the depositional period. When SI of the modern leaf of various ages has been plotted against the known CO₂ concentrations, it shows a linear response and serves as a calibration curve. It indicates that the pCO₂ level was about 342 ppmv during the deposition of the fossil. The result thus obtained is in good agreement with that obtained by means of geochemical methods¹ which point out CO₂ fluctuations between 280 and 360 ppmv. The above method of estimating pCO₂ may not be effective for pre-Cretaceous

sites due to its species specific nature. According to Royer *et al.*⁶, the error estimates for the Tertiary are ± 10–40 ppmv as stomatal density of trees and shrubs increases with altitude, and fossil leaves showing changes in SI through time are actually recording pCO₂ partial pressure. Therefore, for pCO₂ estimation using fossils from sites at higher altitudes, this technique may not be useful.

It is clear from the present study that SI of fossil angiosperm leaves presents an effective alternative method to measure pCO₂ concentration throughout the Cenozoic. Using the fossil cuticle of *T. catappa* from any time scale, it would be possible to deduce the pCO₂ level of that period. More such evidences, based on different fossil leaf compressions, would help in the accurate estimation of palaeoatmospheric CO₂ concentration.

- Kürschner, W. M., Wagner, F., Dilcher, D. L. and Visscher, H., *Geological Perspective of Global Climate Change* (eds Gerhard, L. C., Harrison, W. E. and Hanson, B. M.), American Association of Petroleum Geologists, USA, 2001, pp. 169–189.
- Van der Burgh, J., Visscher, H., Dilcher, D. L. and Kürschner, W. M., *Science*, 1993, **260**, 1788–1790.
- Kürschner, W. M., Van der Burgh, J., Visscher, H. and Dilcher, D. L., *Mar. Micropaleontol.*, 1996, **27**, 299–312.
- McElwin, J. and Chaloner, W. E., *Ann. Bot.*, 1995, **76**, 389–395.
- McElwin, J. and Chaloner, W. E., *Palaios*, 1996, **11**, 376–388.
- Royer, D. L., Berner, R. A. and Beerling, D. J., *Earth Sci. Rev.*, 2001, **54**, 349–392.
- Joshi, A., Tewari, R., Mehrotra, R. C., Chakraborty, P. P. and De, A., *J. Geol. Soc. India*, 2002 (in press).
- Awasthi, N. and Mehrotra, R. C., *Palaeobotanist*, 1995, **44**, 157–188.
- Woodward, F. I., *Nature*, 1987, **327**, 617–618.
- Salisbury, E. J., *Philos. Trans. R. Soc. London, Ser. B*, 1927, **216**, 1–65.

ACKNOWLEDGEMENTS. We thank the Director, Birbal Sahni Institute of Palaeobotany (BSIP), Lucknow and Deputy Directors General, Geological Survey of India for permission to undertake this study and for providing the necessary facilities. We also thank Mr Y. P. Singh, BSIP for his help in generating the computer graphics. Thanks are also due to the Deputy Director, Central National Herbarium, Howrah for permitting R.C.M. to consult the herbarium.

Received 10 June 2002, revised accepted 17 October 2002