



Figure 3. Kernel development at different days of harvest.

correlated to Arginine maturity index (AM)<sup>11</sup>. Phule-pragati in Maharashtra released<sup>2</sup> as early variety was similar to TG-3 in maturity period and other characteristics but inferior in pods yields at Trombay<sup>3</sup>. Therefore, it was not included here. TGE-1 and Chico, which matured early had 20-25% and 10 to 12% ill-filled pods respectively, while, TG-3 later maturing had not. Improvement of this character in early maturing varieties might result in higher pod yields. TGE-1 was superior to Chico not only in yield but also in other characteristics. Although it was inferior to TG-3 in yield its shells were thin like paper and therefore showed 80% shelling. The oil content of 50% in TGE-1 was an improvement in the early maturing

variety.

Presence of foliaceous stipule in TGE-1 would facilitate easy identification of the culture and hybrids during hybridization programme.

1. Baily, W. K. and Hammons, R. O., *Crop Sci.*, 1975, 15, 195.
2. Patil, G. D., Desale, S. C., Patil, P. S., and Patil, S. S., *J. Maharashtra Agr. Univ.*, 1980, 5, 47.
3. Patil, S. H., Chandra Mouli and Kale, D. M., Progress report on varietal improvement in groundnut. IInd research coordinations meeting, FAO/IAEA, Thailand, 1981.
4. Patil, S. H., *Indian. J. Genet.*, 1966, 26A 334.
5. Chandra Mouli and Patil, S. H., *J. Hered.*, 1975, 66, 28.
6. Patil, S. H., *Indian. J. Agric. Sci.*, 1973, 43, 370.
7. Chandra Mouli and Patil, S. H., *Curr. Sci.*, 1979, 48, 310.
8. Patil, S. H., and Chandra Mouli, *Proc. Induced mutation for crop improvement in Africa*, IAEA TECDOC. 1978, 222, 83.
9. Pattee, H. E., Wyne J. C. and Young, J. H., *Peanut. Sci.*, 1977, 4, 47.
10. Majumdar, P. K., Prakash, R. and Hussain, M. A., *Indian. Agric. Sci.* 1970, 40, 292.
11. Hammons, R. O., Tai, P. Y. P. and Young, C. T., *Peanut. Sci.*, 1978, 5, 68.

---

## ANNOUNCEMENT

---

### INDIAN COUNCIL OF MEDICAL RESEARCH - TALENT SEARCH SCHEME

Applications are invited from medical graduates, who have either completed or will complete their internship by the end of 1982 for Talent Search Scheme Fellowships. The candidates below 25 years of age on 1 July 1982 should have passed examinations for MBBS in the first attempt with a minimum of 60% aggregate marks. Candidates will be selected on the basis of an All India Competitive Examination to be held in May 1982. Selected candidates will commence their fellowship once they have obtained admission to a first year residency recognised for the MD degree, in the institution of

their choice. The fellowship is tenable for three years and carries a stipend equivalent to that of Government resident doctors.

The fellows who complete the fellowship successfully will be offered suitable positions in the Institutes under the Council. Full details and blank application forms are obtainable from the Director General, Indian Council of Medical Research (TSS), Ansari Nagar, P.O. Box No. 4508, New Delhi 110 029 before 10 March 1982. Last date for receipt of completed applications is 25 March 1982.