

Clusters of master control genes regulate the development of higher organisms

Edward B. Lewis

How do new genes arise from old ones? This article traces the results of testing the hypothesis that new genes will generally arise by mutation occurring within a set of tandem duplicated genes. Not only do the results support the hypothesis but quite unexpectedly they have identified one of the most ancient clusters of genes known and one that plays a vital role in controlling the fundamental body plan of all higher organisms, including human beings.

THIS article is an account of how basic research on the nature of the gene has led to the discovery of two clusters of master control genes that play a central role in programming the development of the fly. These clusters, known as the bithorax complex (BX-C) and Antennapedia complex (ANT-C), are believed to have evolved from an ancestral gene by a process of tandem duplication and divergence in function by mutation. So successful during evolution have cognates of these clusters been in programming development that they are now found in vertebrates, including humans, as well as in invertebrates.

The strategy of achieving complexity by duplicating and then diversifying is used at the developmental level, as well as at the genetic level. Thus, early in embryonic life, most higher organisms consist of little more than a tandem array of duplicated body segments. During the course of development, the segments diverge from one another and diversify to produce the variety of forms that these organisms achieve. It now seems likely that it is precisely the genes of the BX-C and ANT-C, or their cognates in other organisms, that initiate and control the diversification of the body segments.

This article is concerned primarily with the BX-C, how it came to be discovered, and how we think it functions during development.

Early history

The view of genes as beads on a string, randomly arranged by function, was forever altered in 1925 by

Sturtevant's discovery that the function of a gene is dependent on its position in the chromosome¹. He came on this phenomenon of 'position effect' in the course of analysing one of the first cases of unstable genes in *Drosophila*, the famous *Bar*-eyed mutant, found by Tice in 1913. Sturtevant showed that the occurrence of two exceptional types of progeny from homozygous *Bar* flies—non-*Bar* (normal-eyed) and double-*Bar* (tiny-eyed) types—was associated with crossing-over of an unusual kind. He assumed that non-*Bar* types had lost the *Bar* gene and that double-*Bar* was a tandem duplication of that gene. (A detailed account of this early work and of relevant crossing-over mechanisms is given by Maeda and Smithies².)

In 1936, using the then recently discovered giant salivary gland chromosomes of the *Drosophila* larva, Bridges³ and Muller *et al.*⁴ showed, independently, that the *Bar* mutation is itself a tandem duplication of seven bands, and double-*Bar* is a tandem triplication, rather than a duplication, for those bands. The origin of this triplication can be easily pictured if, for present purposes, we represent *Bar* as having the sequence ABCABC and if we introduce a crossover between two homologous ABC regions that are unequally paired. Thus,

$$\begin{array}{c} \text{ABCABC} \\ \times \longrightarrow \text{ABCABCABC} \\ \text{ABCABC} \end{array}$$

Non-*Bar* is, of course, readily accounted for as the reciprocal crossover product, ABC.

From homozygotes for the triplication a similar kind of crossing-over event is expected to generate a single copy of ABC and a fivefold tandem repetition of ABC. Just such a quintuplication was generated by Rapoport⁵ who called it quadruple-*Bar*, since at the time he would not have known that *Bar* itself was a duplication. Later,

Edward B. Lewis is in the Division of Biology, California Institute of Technology, Pasadena, CA 91125, USA

The 1991 Albert Lasker Medical Award was presented to Dr Lewis on September 27, 1991, in New York City.

Reprinted from JAMA⁶ *The Journal of the American Medical Association*, March 18, 1992, Volume 267. Copyright 1992, American Medical Association.

Rapoport obtained a sevenfold to ninefold repetition from homozygous quadruple-*Bar* females, as described by Lindsley and Grell⁶. These early studies showed for the first time that, starting with a tandem duplication, multiple linear repetition can be generated by successive rounds of crossing-over.

In a second development, Bridges⁷ called attention to several types of salivary gland chromosome banding patterns, which he interpreted as duplications that have become established in the species. Genetic evidence for such duplications was however lacking.

In 1940, Oliver⁸ reported that he could recover apparent reverse mutations associated with crossing-over between two *lozenge* eye mutants of *Drosophila*. An alternative interpretation was that the reversions were not mutations but instead were wild-type crossovers between two separate but closely linked loci. Failure to find the reciprocal crossover having both mutants in the same chromosome left the validity of the latter interpretation in limbo.

Another pair of rough-eyed mutants, *Star* (*S*) and *asteroid* (*ast*), proved more tractable to analysis. (This mutant, originally called *Star-recessive*⁹, was found by E. Novitski, who sent it to me in 1937, suggesting that I test to see if it was an allele of *S*. We were undergraduates at the time, he at Purdue University and I at the University of Minnesota, where C. P. Oliver had kindly given me space in his laboratory to work on *Drosophila*. Novitski and I had worked together on *Drosophila* while in high school in Wilkes-Barre, Pa.) From the *trans* heterozygote, *S* +/+ *ast*, not only wild-type crossovers but the reciprocal double mutant, *S ast*, could be derived as well¹⁰. Moreover, there is a striking position effect. Whereas *S* +/+ *ast* animals are nearly eyeless and have wing defects, *S ast* +/+ animals are wild-type except for a slightly roughened eye identical with that of *S*/+ animals^{10,11}. Thus, only when the two wild-type alleles (symbolized by + signs) are located together in the same chromosome or in *cis* do they function normally. The *cis* heterozygotes for *lozenge* mutants were eventually obtained by Green and Green¹²; whereas the *trans* heterozygotes have small eyes, the *cis* have normal eyes. A comparison of the *cis* and *trans* phenotypes has come to be known as the *cis-trans* test, and the position effect, if present, as the *cis-trans* effect (see Hayes¹³ for a discussion of the history of these concepts).

The *S* and *ast* loci proved to be within a double-banded structure known as 21E1-2 near the tip of the second chromosome¹⁴. Not only had Bridges⁷ interpreted such structures as tandem duplications that had become established in the species, he had actually cited 21E1-2 as the type example. Although this correlation of genetic and cytological observations seemed to lend support to the tandem gene duplication hypothesis, we still do not know the true significance of such doublet

structures, and proof that they consist of separate, homologous bands has not been forthcoming.

The BX-C

In 1946, a deliberate search was begun to find additional cases of the *S-ast* type to test the theory that a common mode of gene evolution would involve tandem gene duplication. The resultant genetic redundancy would allow one of the genes to evolve a new function while the other gene continued to carry out the old function¹¹.

Of several cases that we explored, the most illuminating from a developmental standpoint was that of a series of mutants that transform the third thoracic segment (T3) of the fly toward the second (T2). The type mutant, *bithorax* (*bx*), was discovered by Bridges in 1915 (as reported by Lindsley and Grell⁶); it is a weak and variable mutant. Fortunately, a very strong and constant mutant, *bx*³, was found later by Stern (also as reported by Lindsley and Grell⁶) and is the one that has been most useful in dissecting the *bx* function and contributing to the creation of the four-winged fly (Figure 1*b*). The details of the early work on these mutants need not concern us herein except to note that three types of mutants, *bx*, *Ultrabithorax* (*Ubx*) (formerly *bxD*), and *bithoraxoid* (*bxd*) mapped to three closely linked loci on the genetic map. Cytologically, they mapped to two adjacent doublet structures that often fuse into one structure, presumably as the result of pairing of partially homologous bands (Figure 2)¹⁵. Again, Bridges' prediction that such structures represent tandem duplications seemed well supported.

From the start, we found dramatic *cis-trans* effects between almost every double mutant combination tested¹¹. Furthermore, these effects were soon found to be conspicuously polarized. Thus, if we let *a* and *b* represent any two recessive mutants that are qualitatively different in phenotype, then in every case in which *a b* +/+ is wildtype and *a* +/+ *b* is mutant, the animal shows either the *a* phenotype or the *b* phenotype, but not both. For example, *bx*³ +/+ *pbx* (*postbithorax*) has a weak *pbx* phenotype (posterior T3 transformed toward posterior T2) but no trace of the *bx* phenotype (anterior T3 transformed toward anterior T2). Only much later were such polarized effects discovered in the operons of bacteria¹⁶.

Analysis by gene deletion

A new dimension to the analysis came with the derivation of X-ray-induced deletions for the BX-C, or for portions thereof. Thus, animals lacking all of the BX-C (homozygous for a deletion of the two 89E doublets) die as late embryos but not before they have developed the




Figure 1. Photographs of adult wild-type and homeotic mutant flies. **A**, Wild-type male (genotype: *heldout*, in order to produce spread wings) **B**, Dorsal view of a four-winged fly homozygous for the triple mutant combination *anterobithorax bithorax³ postbithorax*. The third thoracic segment is transformed into one resembling the second, resulting in two sets of nearly identical thoraces and wings. **C**, Ventral view of an eight-legged fly of the genotype *bithoraxoid/Df*, where *Df* is a deficiency for the *bithoraxoid* gene. **D**, Dorsal view of an *Antennapedia* male (genotype: *Antp⁴/Pc³* [where the *Pc* mutant enhances the *Antp* phenotype]).

intricate cuticular and tracheal systems that enable the body segments to be distinguished from one another. Such animals show a striking transformation of the abdominal segments transformed into a row of thoracic-like segments¹⁵. Thus, complete loss of the BX-C results in a more extreme homeotic transformation than that produced by the spontaneous mutants, all of which turned out to represent only partial losses of gene function.

What do the BX-C genes do?

In 1964, we reported that mutants of the BX-C are autonomously expressed in somatic mosaics¹⁷. We therefore knew that the developmental abnormalities were not due to substances, such as hormones, that diffuse readily between cells. Our mosaic studies further demonstrated that even late in development, a *bx*³

mutant expressed the *bx* transformation even in a single cell of an otherwise wild-type fly¹⁸. Thus, the wildtype gene could be inferred to function throughout development, at least until the pupal stage.

It has always been somewhat treacherous to deduce the normal function of a gene from the effects on the organism of mutations in that gene. The problem is especially severe when the mutations result in morphological changes. To assess the normal or wild-type functions of the BX-C, we adopted a new procedure. We constructed a set of genotypes in which the BX-C was entirely lacking except for the presence of one dose of various wild-type regions of the complex. In this way, we could show that all of the regions studied are involved in initiating, or in some cases suppressing, the formation of specific organs and structures, a more profound and basic type of developmental function than had been inferred from considering the homeotic effects shown by the mutants^{15,18}.




Figure 2. A correlation of the cytological and genetic maps of the bithorax complex. The bithorax complex maps to the two doublets in 89E of the third chromosome of the salivary gland chromosomes. Dominant gain-of-function mutants are in the row above that of the loss-of-function mutants. *Ubx* indicates *Ultrabithorax*; *abx*, *anterobithorax*; *bx*, *bithorax*; *Cbx*, *Contrabithorax*; *bxd*, *bithoraxoid*; *pbx*, *postbithorax*; *iab*, *infra-abdominal*; *Hab*, *Hyperabdominal*; *Mcp*, *Miscadastral pigmentation*; and *Tab*, *Transabdominal*.

Analysis of the function of the wildtype BX-C genes during development revealed two basic properties of the system. First, the genes are arranged in the chromosome in the same order in which they are expressed along the anteroposterior axis of the organism (Figure 3)^{15,19}. This colinearity has now been found in all clusters of the BX-C type in which the order in the chromosome has been established, as will be discussed herein. Second, the morphological evidence showed that once expressed in a given segment, the

gene tends to remain expressed in more posterior segments (Figure 4)¹⁵. Thus, the more posterior a segment in the organism, the more advanced in development it is as far as genes of the BX-C are concerned, the ground state being that of a thoracic state¹⁵.

Further subdivision of the complex was achieved not only by means of overlapping deletions but by rearrangement breakage points in the complex^{15,20}. The latter have proved useful in identifying a set of *cis*-regulatory regions that direct the control of gene expression in each of the abdominal segments from A2 to A9, inclusive. A loss-of-function mutation in one of these regions, designated *infra-abdominal* (*iab*), causes the segment involved to transform to the next most anterior type of segment, i.e., toward the thoracic or ground state.

Before recombinant DNA methodology was applied to the BX-C, we interpreted the wild-type action of the BX-C genes in terms of the operon model¹⁶ for gene clusters in prokaryotes. Thus, many of the genetic properties of the BX-C, including polarized *cis-trans* effects and the presence of dominant constitutive mutants (Figure 2), appeared to be remarkably similar to the genetic properties of operons. This led us to suggest that the BX-C might constitute an inducible system in which the function of its genes would be to 'repress certain systems of cellular differentiation and thereby allow other systems to come into play'¹⁷. However, the molecular basis of the BX-C, which will be discussed, proves to be entirely different from that of the operon. Our earlier concept of the function of the BX-C genes was clearly too narrow. Quite different and only partially understood mechanisms induce the expression of the BX-C genes, and we now suppose that these genes can activate, as well as repress, systems of cellular differentiation.




Figure 3. Summary of *Drosophila* bithorax complex functions and their spatial expression. The parasegmental (PS) morphological function and protein expression for PS5 to PS14 (segments T3 to A9) are shown in parentheses and are inclusive. The extent of *Ultrabithorax* function and protein expression is shown in green, that of *abdominal-A* (*abd-A*) in blue, *Abd-BI* in yellow, and *Abd-BII* in orange. *bx* indicates *bithorax*, *bxd*, *bithoraxoid*, and *iab*, *infra-abdominal*.




Figure 4. Colinearity of order of gene expression in the adult fly and the order on the molecular map of the *cis*-regulatory regions. The bithorax complex encompasses 300 kb (from +200 kb to -100 kb, where the start of the walk was 0.0). Below the map, which is approximately to scale, are five characterized transcription units, three containing homeoboxes, *Abd-B*, *abd-A*, and *Ubx*, and two noncoding, *iab-4* and *bxd*, *iab* indicates *infra-abdominal*, *pbx*, *postbithorax*, *bxd*, *bithoraxoid*, *bx*, *bithorax*, *abx*, *anterobithorax*, *abd*, *abdominal*, and *Ubx*, *Ultrabithorax*.

One of the most challenging problems remaining will be to identify the genes that the BX-C genes control. Such 'downstream' or 'target' genes are those that we assume will bring about the differentiation of tissues and organs of the third thoracic and abdominal regions of the organisms.

The molecular genetics of the BX-C

In 1978, Prof. David Hogness and his collaborators at Stanford University began a molecular analysis of the BX-C. They used chromosome walking and jumping to clone the BX-C²⁰. The current status of the molecular map of the complex is shown in the lower half of Figure 4 based on this early work and subsequent work^{21,22}. Three domains of function have come to be recognized: *Ubx*, *abdominal-A* (*abd-A*), and *Abdominal-B* (*Abd-B*), named after the protein class coded by each domain¹⁸. The *Ubx* domain has four exons and a very large intron within which lie the *anterobithorax* (*abx*) and *bx* loci. As the result of alternative splicing, a family of proteins results²³. For the sake of simplicity, only the largest transcription unit is shown in Figure 4.

The colinearity rule, first noticed for morphological effects, as already discussed, has been fully supported by studies of the expression of proteins coming from each

of the three domains of the BX-C using immunohistochemical techniques (Figure 4). Thus, in wild-type embryos, the *Ubx* proteins are first detected in posterior T2 and anterior T3^{24,25} (or in terms of the parasegmental [PS] units²⁶ of the embryonic ectoderm, in PS5). The expression declines but remains in the abdominal segments^{25,27}. Analysis of mutant embryos indicates that the *bx* and *bxd* regions direct *Ubx* protein expression in PS5 and PS6, respectively^{26,27}.

The level of *abd-A* protein in wildtype embryos remains elevated in these segments and those following up to and including PS13^{28,29}. Double labeling with antibodies to *Ubx* and *abd-A* proteins has shown that *Ubx* proteins, although reduced in amounts, are on in different cell types than is the *abd-A* protein²⁹. Analysis of mutants in the *iab-2*, *iab-3*, and *iab-4* regions shows that *abd-A* protein is expressed in a colinear fashion for PS7, PS8, and PS9^{29,30}.

The *Abd-B* region produces a large and small protein³⁰⁻³². Our analysis of the distribution of these proteins has been indirect since our monoclonal antibody is directed to an epitope in the 3' exon common to both proteins of this region. To infer their distribution, therefore, we have used mutants that eliminate the messenger RNA for either the small or the large protein. The latter is first detected in the posterior

fourth abdominal segment (PS10), while the small protein coming from the more distally located promoter is first detected in the posterior eighth abdominal segment (PS14). Using rearrangement break points disrupting the *iab-5* to *iab-9* regions, inclusively, we have shown that the pattern of expression of the large protein is colinear with the order of those regions³³.

Thus, the expression of proteins from the three domains of the complex parallels the morphological expression. It confirms and establishes the rule that the more posterior the segment, the more functions controlled by the *cis*-regulatory regions become expressed up to and including PS13. Thereafter, there is loss of detectable expression of all but the small *Abd-B* protein.

The colinearity rule also applies to RNAs coming from the *iab* regions of the complex, as shown by Sanchez-Her-rero and Akam³⁴. They point out that these transcripts may be functionless, as discussed for the *bxd* case by Lipshitz *et al.*³⁵ or the *iab* transcripts 'might function in *cis* by some unprecedented mechanism'³⁴.

The ANT-C

It is likely that the BX-C was once part of a single cluster that included another larger complex, the ANT-C³⁶. The latter lies in the same chromosome arm but at a considerable distance proximal to the BX-C. The type mutant for which this complex was named was first found in *Drosophila affinis* by Sturtevant³⁷. The phenotype is that of an antenna transformed toward a leg. A mutant with a strong transformation of this type (Figure 1 d) was induced with X-rays in 1948 by Yu as described by Lindsley and Grell⁶. In the same year, Les Calves reported the occurrence of a neutroninduced mutant that he named *Aristapedia* and suggested that it was a dominant allele of a recessive mutant and causes the antenna to transform toward a tarsus, rather than a full leg. From his determination of its cytological location in 84A, the location of the ANT-C (and of *aristapedia* in 89C⁶), it is now clear that it is in fact an allele of the *Antennapedia* locus.

Extensive genetic and cytological analyses of these and other related mutants by Kaufman and colleagues³⁸ established the existence of this second cluster of homeotic mutants and their colinearity (Figure 5). Extensive molecular analyses have been made, particularly of the *Antp* gene itself^{39,40}.

Regulation of the genes of the BX-C in trans

A class of genes that initiates the axes of the body and divides the body into segments has been identified by Nüsslein-Volhard and Wieschaus⁴¹ based on massive




Figure 5. Correlation of homeotic gene expression in the *Drosophila* embryo with that of the mouse embryo. The anterior ends of the embryos are to the left. The *Drosophila* homeotic genes form two clusters in the right arm of the third chromosome, the separation being between the *Antp* and *Ubx* genes. The homologous mouse genes are in a single cluster, among the four such clusters only that of *Hox2* in chromosome 11 is shown. The genes are color-coded; however, the homeobox sequences of the *Antp*, *Ubx*, and *abd-A* genes of *Drosophila* are so similar that homology with the mouse genes is based only on the expression patterns. In the *Drosophila* and mouse embryos, the anterior boundary of expression of each gene is shown by the corresponding color for each gene. For simplicity, the posterior boundary for a given color is not shown since it would usually continue beyond the next most posterior gene's anterior boundary. Only expression in the cuticle of the *Drosophila* embryo is shown, while in the mouse, expression is shown in the nervous system and in the prevertebrae. *lab* indicates labial; *pb*, Proboscipedia; *Dfd*, Deformed; *Scr*, Sex-combs reduced; *Ubx*, Ultra-bithorax; and *abd*, abdominal.

screenings for mutants affecting early development of *Drosophila*^{42,43}. Some of these genes appear to act as major *trans*-regulators of the BX-C and ANT-C. A possible relationship of one such gene, *hunchback* (*hb*), to the BX-C was first detected genetically when *Regulator-of-phx* (*Rg-phx*), a dominant *trans*-regulator of the *post-bithorax* function of the BX-C⁴⁴, was found by Bender *et al.*⁴⁵ to be a gain-of-function mutant of the *hb* gene. Garcia-Bellido and Capdevila⁴⁶ had earlier made extensive studies of the role of *Rg-phx* in regulating the BX-C. Recently, Qian *et al.*⁴⁷ have shown the *hb* protein binds to a specific motif in the intron of the *Ubx* gene and interacts to repress transcription of *Ubx* in regions anterior to the onset of its normal transcription in PS5 of that gene. The expression of *Ubx* has been inferred to be *trans*-

regulated by still other genes of the segment-initiation class^{43,44-46}. Thus, the genes that initiate segment formation seem to represent a major and perhaps primary set of regulators of the BX-C and ANT-C.

Several genes act as negative regulators of the BX-C; that is, if such a gene is removed or inactivated, many if not all of the genes of the complex become activated, resulting in all of the segments of the body being converted toward the eighth-abdominal segment¹⁵. The first of these genes was *Polycomb*, found by P. H. Lewis as reported by Lindsley and Grell⁶. Zink and Paro⁵² have shown that the protein coded for by the normal *Polycomb* gene is specifically associated in the salivary gland chromosomes to the BX-C and ANT-C regions, as well as other sites. Zink *et al.*⁵³ have shown in an elegant way that it binds to upstream (5') sequences of the *Antp* gene. Paro and Hogness⁵⁴ find that the sequence of the *Polycomb* gene shows some homology, not as might have been expected to a DNA-binding class of proteins but to a nonhistone chromosomal protein. Hence, the remarkably precise binding of *Polycomb* protein to bands in the salivary gland chromosomes associated with the BX-C and ANT-C suggests that the protein is complexing with one or more other proteins that do bind to those bands. If so, the *Polycomb* class of genes¹⁸ would be involved in maintaining a repressed state in regions of the embryo where it is activated^{55,56}. Ingham and Whittle⁵⁷ have shown that another gene, *trithorax*, probably allelic to *Regulator-of-bx* (*Rg-bx*), acts as a positive regulator. Ingham⁵⁸ suggests that it may be involved in maintaining the activated state of the BX-C genes.

The homeobox

Support from molecular studies for the hypothesis that the BX-C and the ANT-C owe their origin to repeated tandem gene duplication finally came with the discovery of the homeobox by McGinnis *et al.*⁵⁹ and independently by Scott and Weiner⁶⁰. This region of 180 base pairs codes for 60 amino acids that constitute the DNA binding or 'homeodomain' of each of the proteins coded by the BX-C and ANT-C genes. It is now thought likely that these proteins bind to downstream genes and thereby regulate them.

So similar are the homeobox sequences between adjacent genes that it is exceedingly improbable that they arose by convergent evolution. Only the homeodomain of the protein shows obvious conservation, possibly because most of the specificity required for DNA binding lies in the homeodomain. We would expect, however, that many motifs in enhancerlike elements of the *cis*-regulatory regions may be conserved. At present, there are insufficient data on DNA sequence in the *cis*-regulatory regions to test for conservation of such motifs.

Homeobox gene clusters in other organisms

Genetic and morphological evidence for cognates of the homeobox-containing genes of the ANT-C and BX-C have also been discovered in beetles, where they appear to lie within a single homeotic complex (HOM-C)^{61,62}. (In the silkworm *Bombyx*, a fascinating set of mutants have striking parallels to many of the mutants of the BX-C and ANT-C. They were designated as alleles of an *E* locus within which, however, crossingover and *cis-trans* effects have been reported^{63,65}, suggesting that they are in fact a gene complex; however in the absence of a genetic or molecular map, the colinear rule obviously cannot be tested. One of the *E* mutants produces an eight-legged moth [a bronze sculpture of which is in an ancient temple in Japan]. The exact cognate of that mutant in *Drosophila* is *bxd* [Figure 1c]. Another of the *E* mutants was reported to lack gonads and led us to examine its cognates in *Drosophila*, namely *iab-4* mutants. When homozygous, the latter mutants are viable and appear virtually wild-type but internally they too lack gonads in both sexes¹⁵.)

Astonishingly, mice and humans not only have cognates of the BX-C and ANT-C genes in a single HOM-C, but the complexes occur in four sets, each in a different chromosome⁶⁶⁻⁶⁹. Moreover, colinearity of spatial order of expression and molecular map order extends over the entire complex in the murine (Figure 5) and in the human cases. Other less extensively characterized clustering of cognates of the HOM-C type occur in sea urchins⁷⁰, nematodes^{71,72}, bees⁷³, frogs⁷⁴, and chickens⁷⁵. Omitted are a number of other organisms in which cognates have been found for only one or two genes of the HOM-C type^{76,77}.

In *Drosophila*, it is not yet known whether temporal order of expression is also colinear with map order. However, there is evidence that colinearity is temporal as well as spatial from experiments involving (1) chicken⁷⁵ and murine⁷⁸ limb buds; (2) expression of transcripts from HOM-C in the central nervous system and prevertebrae of the mouse embryo⁷⁹; and (3) induction of the HOM-C by retinoic acid in human cell cultures⁸⁰.

Why have the HOM-C remained intact?

A fundamental question is why genes of the HOM-C have remained linked for an estimated 500 million years or more, assuming they arose before the separation of the vertebrates and invertebrates. Even though they presumably evolved by tandem gene repetition, it is highly unlikely that the linkage would have persisted in so many organisms unless it confers a selective advantage.

Clearly, in *Drosophila*, the existence of two separate

complexes, which still remain tightly linked, suggests that at least the *Antp* and *Ubx* genes do not have to be adjacent in this organism. Whether other genes, such as *Ubx* and *abd-A*, can be separated and still function normally is uncertain. Several rearrangements with break points near the boundary of the *Ubx* and *abd-A* domains have been detected by their suppression of transvection^{81,82} in a screen that recovers break points that are unselected, with respect to having any mutant effect¹⁹. These rearrangements show a striking phenotype; namely, development of tiny sense organs on the larval abdominal segments (organs normally found only on the larval thoracic segments), an effect that corresponds to a weak *bxd* phenotype. Bender *et al.*⁸³ have shown that such break points map near the boundary of the *Ubx* and *abd-A* domains; however, from their *bxd* phenotype, it is clear that they still lie in the *Ubx* domain. Thus, if a break point can fall at or near that boundary and lack a detectable phenotype, our screen has yet to recover one.

A possible clue to understanding why gene clusters of the BX-C and ANT-C type have remained intact comes from studies of gene regulation in another cluster; namely, that coding for the family of β -globin peptides of hemoglobin. Regulation of the genes of this cluster has been shown to depend on enhancer elements residing in the *cis*-regulatory regions surrounding the protein-coding regions of the genes. Nickol and Felsenfeld⁸⁴ and Choi and Engel⁸⁵ find that the adult and embryonic β -globin genes in chickens share an enhancer.

In the BX-C, Celniker *et al.*³³ have evidence, based on morphology, that the *iab-5* region, which normally regulates the expression of the *Abd-B* gene, may be regulating expression of both *abd-A* and *Abd-B*. They suggest, therefore, that in wild-type, the *abd-A* and *Abd-B* genes share one or more enhancers located in the *iab-5* region. Duncan¹⁸ has discussed this and other cases in the complex where *cis*-regulatory elements seem to be 'bifunctional'. If enhancer sharing does occur in the BX-C and if it confers a selective advantage, then the genes involved will of course tend to remain linked.

Why four sets of complexes in mice and humans?

As just noted, homeobox (HOX) clusters have been duplicated in the genome of mice and humans four times, each in a different chromosome in both organisms. In humans, four clusters are designated HOX1, HOX2, HOX3, and HOX4 and are located in chromosomes 7, 17, 12, and 2, respectively. Boncinelli *et al.*⁶⁷ have speculated that having four sets might provide 'a finer-grained spatial control' than would be needed in organisms such as *Drosophila*, which have only one set and perhaps fewer requirements for setting up the body segments than have the higher vertebrates.

Medical implications

We can look forward to an exciting time when the functions of the genes of the four human HOX clusters are determined. Human genetic abnormalities at virtually any stage of development, but especially those traceable to abnormalities of early embryonic or fetal development, would be expected from mutations in genes of the HOX clusters. The results of genetic experiments on other vertebrates strongly support such a conjecture.

Recently, Chisaka and Capecchi⁸⁶ used the new technique of gene targeting to produce a disrupted *Hox1.5* gene of the mouse. Homozygotes for this recessive loss-of-function mutation are found to produce a constellation of defects that they point out is remarkably similar to the human congenital disorder DiGeorge syndrome. The obvious implications are that such a syndrome might result from a mutation in the human cognate of the mouse gene and that other congenital disease syndromes might result from mutations in other HOX genes.

Balling *et al.*⁸⁷ overexpressed the murine *Hox1.1* (*Hox2.3* of the HOX2 cluster) gene in transgenic mice and observed craniofacial abnormalities. Kessel *et al.*,⁸⁸ studying more severely affected mice from the same experiment, observed dramatic homeotic effects on the cervical vertebrae. It follows that a gain-of-function mutation in the human cognate of *Hox1.1* is expected to show broadly similar types of defects, depending on the penetrance of the allele.

In *Drosophila*, it is well known that such gain-of-function mutants are not uncommon (Figure 2). The first example in the BX-C was the *Contrabithorax* (*Cbx*) mutant, which causes T2 to transform toward T3, notably turning the wings into halteres^{15,17}. All of the *Antennapedia* mutants are also now known to be an overexpression of the wild-type gene in the antenna (Figure 1d). We would expect that when mutants of HOX genes in humans are found they will often be of the gain-of-function type.

The HOX genes may also be implicated in the genesis of malignancies. Blatt *et al.*⁸⁹ have reported a possible example involving a murine myeloid leukemia cell line in which a *Hox2.4* gene has been disrupted by a retroviral insertion. Their finding is confirmed by Kongsuwan *et al.*⁹⁰ who have studied the same cell line and who suggest that *Hox2.4* may interfere with terminal differentiation; then, in combination with activation by the same provirus of another gene in the same cell line, conditions would be met for development of the transformed cell. Blatt and Sachs⁹¹ also report that deletion of the *Hox4.1* gene occurs in another murine myeloid leukemia cell line; this gene is the cognate of *Hox2.7* (Figure 5). They have discussed a number of mechanisms that might involve such a

deletion in the generation of the malignancy.

To summarize the medical relevance of the discovery of HOX clusters in humans, we can expect that germinal mutations occurring within these clusters are likely to result in abnormalities at many stages of human development. On the basis of preliminary findings in murine leukemia cell lines, it is conceivable that somatic mutations in the HOX clusters may potentially play a significant role in the generation of tumors, both benign and malignant. However, the search for DNA lesions in potential cases of germinal or somatic mutations presents formidable technical problems, not only because of the large size of such clusters but because of their fourfold redundancy.

1. Sturtevant, A. H., *Genetics*, 1925, 10, 117-147.
2. Maeda, N. and Smithies, O., *Annu. Rev. Genet.*, 1986, 20, 81-108.
3. Bridges, C. B., *Science*, 1936, 83, 210-211.
4. Muller, H. J., Prokofyeva-Belgovskaya, A. A. and Kosikov, K. V., *C. R. (Doklady) Sci. URRS*, 1936, 1, 87-88.
5. Rapoport, J. A., *Bull. Biol. Med. Exp.*, 1936, 11, 242-244.
6. Lindsley, D. L. and Grell, E. H., *Genetic Variations of Drosophila Melanogaster*, Washington, DC, Carnegie Institution of Washington, 1968, pp. 1-521, Publication 627.
7. Bridges, C. B., *J. Hered.*, 1935, 26, 60-64.
8. Oliver, C. P., *Proc. Natl. Acad. Sci. USA*, 1940, 26, 452-454.
9. Lewis, E. B., *Proc. Minn. Acad. Sci.*, 1939, 7, 23-26.
10. Lewis, E. B., *Genetics*, 1942, 27, 153-154.
11. Lewis, E. B., *Cold Spring Harbor Symp. Quant. Biol.*, 1951, 16, 159-174.
12. Green, M. M. and Green, K. C., *Proc. Natl. Acad. Sci. USA*, 1949, 35, 586-591.
13. Hayes, W., *The Genetics of Bacteria and their Viruses: Studies in Basic Genetics and Molecular Biology*, John Wiley & Sons, New York, 1968, 2nd edn.
14. Lewis, E. B., *Genetics*, 1945, 30, 137-166.
15. Lewis, E. B., *Nature*, 1978, 276, 565-570.
16. Jacob, F. and Monod, J., *Cold Spring Harbor Symp.*, 1961, 26, 193-211.
17. Lewis, E. B., in *Role of Chromosomes in Development* (ed. Locke, M.) Academic Press, New York, 1964, pp. 231-252.
18. Duncan, I., *Annu. Rev. Genet.*, 1987, 21, 285-319.
19. Lewis, E. B., *Cold Spring Harbor Symp. Quant. Biol.*, 1986, 50, 155-164.
20. Bender, W. *et al.*, *Science*, 1983, 221, 23-29.
21. Karch, F. *et al.*, *Cell*, 1985, 43, 81-96.
22. Celniker, S. E. and Lewis, E. B., *Genes Dev.*, 1987, 1, 111-123.
23. Kornfeld, K., Saint, R. B., Beachy, P. A., Harte, P. J., Peattie, D. A. and Hogness, D. S., *Genes Dev.*, 1989, 3, 243-258.
24. White, R. A. H. and Wilcox, M., *Cell*, 1984, 39, 163-171.
25. Beachy, P. A., Helfand, S. L. and Hogness, D. S., *Nature*, 1985, 313, 545-551.
26. Lawrence, P. A. and Martinez-Arias, A., *Philos. Trans. R. Soc. London Biol.*, 1985, 312, 83-90.
27. Hogness, D. S. *et al.*, *Cold Spring Harbor Symp. Quant. Biol.*, 1985, 50, 181-194.
28. Karch, F., Bender, W. and Weiffenbach, B., *Genes Dev.*, 1990, 4, 1573-1587.
29. Macias, A., Casanova, J. and Morata, G., *Development*, 1990, 110, 1197-1207.
30. Celniker, S. E., Keelan, D. J. and Lewis, E. B., *Genes Dev.*, 1989, 3, 1424-1436.
31. Delorenzi, M. and Bienz, M., *Development*, 1990, 108, 323-329.
32. Boulet, A. M., Lloyd, A. and Sakonju, S., *Development*, 1991, 111, 393-405.
33. Celniker, S. E., Sharma, S., Keelan, D. J. and Lewis, E. B., *EMBO J.*, 1990, 9, 4277-4286.
34. Sanchez-Herrero, E. and Akam, M., *Development*, 1989, 107, 321-329.
35. Lipshitz, H., Peattie, D. and Hogness, D., *Genes Dev.*, 1987, 1, 307-322.
36. Kaufman, T. C., Seeger, M. A. and Olsen, G., *Adv. Genet.*, 1990, 27, 309-362.
37. Sturtevant, A. H., *Genetics*, 1940, 25, 337-353.
38. Kaufman, T. C., Lewis, R. and Wakimoto, B., *Genetics*, 1980, 94, 115-133.
39. Garber, R. L., Kuroiwa, A. and Gehring, W. J., *EMBO J.*, 1983, 2, 2027-2036.
40. Scott, M. P. *et al.*, *Cell*, 1983, 35, 763-776.
41. Nusslein-Volhard, C. and Wieschaus, E., *Nature*, 1980, 287, 795-801.
42. Nusslein-Volhard, C., Frohnhofer, H. G. and Lehmann, R., *Science*, 1987, 230, 1675-1681.
43. Ingham, P. W., *Nature*, 1988, 335, 25-34.
44. Lewis, E. B., Paper presented at the XII International Congress on Genetics, August 26, 1968, Tokyo, Japan.
45. Bender, M., Turner, F. R. and Kaufman, T. C., *Dev. Biol.*, 1987, 119, 418-432.
46. Garcia-Bellido, A. and Capdevila, M. P., in *The Clonal Basis of Development* (eds. Subtelny, S. and Sussex, I. M.), 36th Symposium of the Society for Developmental Biology, Orlando, Fla, Academic Press, 1978, pp. 3-21.
47. Quian, S., Capovilla, M. and Pirrotta, V., *EMBO J.*, 1991, 10, 1415-1425.
48. Ingham, P. W. and Martinez-Arias, A., *Nature*, 1986, 324, 592-597.
49. Ingham, P. W., Ish-Horowicz, D. and Howard, K. R., *EMBO J.*, 1986, 5, 1659-1665.
50. Martinez-Arias, A. and White, R. A. H., *Development*, 1988, 102, 325-338.
51. Irish, V. F., Martinez-Arias, A. and Akam, M., *EMBO J.*, 1989, 8, 1527-1537.
52. Zink, B. and Paro, R., *Nature*, 1989, 337, 468-471.
53. Zink, B., Engstrom, Y., Gehring, W. J. and Paro, R., *EMBO J.*, 1991, 10, 153-162.
54. Paro, R. and Hogness, D. S., *Proc. Natl. Acad. Sci. USA*, 1991, 88, 263-267.
55. Struhl, G., *Proc. Natl. Acad. Sci. USA*, 1992, 79, 7380-7384.
56. Struhl, G. and Akam, M., *EMBO J.*, 1985, 4, 3259-3264.
57. Ingham, P. W. and Whittle, J. R. S., *J. Mol. Gen. Genet.*, 1980, 179, 607.
58. Ingham, P. W., *Wilhelm Roux's Arch. Dev. Biol.*, 1981, 190, 365.
59. McGinnis, W., Levine, M., Hafen, E., Kuroiwa, A. and Gehring, W. J., *Nature*, 1984, 308, 428-433.
60. Scott, M. P. and Weiner, A. J., *Proc. Natl. Acad. Sci. USA*, 1984, 81, 4115-4119.
61. Beeman, R. W., *Nature*, 1987, 327, 247-249.
62. Beeman, R. W., Stuart, J. J., Haas, M. S. and Dennell, R. E., *Dev. Biol.*, 1989, 133, 196-209.
63. Tsujita, M., *Jpn. J. Genet.*, 1955, 30, 227-235.
64. Itikawa, N., *Bull. Seric. Expt. Station Tokyo*, 1952, 14, 23-91.
65. Tazima, Y., *The Genetics of the Silkworm*, Logos, London, England, 1964.
66. Gaunt, S. J., Sharpe, P. T. and Duboule, D., *Development*, 1988, 104, (suppl), 169-179.
67. Boncinelli, E. *et al.*, *Hum. Reprod.*, 1988, 3, 880-886.
68. Krumlauf, R., Holland, P. W. H., McVey, J. H. and Hogan, B. L. M., *Development*, 1987, 99, 603-617.
69. Schugart, K., Kappen, C. and Ruddle, F., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 7067-7071.

70. Humphreys, T., *Zool. Sci.*, 1987, 4, 1129.
71. Burglin, T. R. *et al.*, *Nature*, 1991, 351, 703.
72. Kenyon, C. and Wang, B., *Science*, 1991, 253, 516-517.
73. Walldorf, U., Fleig, R. and Gehring, W. J., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 9971-9975.
74. Wright, C. V. E., Cho, K. W. Y., Hardwicke, J., Collins, R. H. and De Robertis, E. M., *Cell*, 1989, 59, 81-93.
75. Nohno, T. *et al.*, *Cell*, 1991, 64, 1197-1205.
76. Gehring, W. J. and Hiromi, Y., *Annu. Rev. Genet.*, 1986, 20, 147-173.
77. Scott, M. P., Tamkun, J. W. and Hartzell, G. W., *Biochem. Biophys. Acta*, 1989, 989, 25-48.
78. Graham, A., Papalopolu, N. and Krumlauf, R., *Cell*, 1989, 57, 367-378.
79. Wilkinson, D. G., Bhatt, S., Cook, M., Boncinelli, E. and Krumlauf, R., *Nature*, 1989, 341, 405-409.
80. Simeone, A. *et al.*, *Nature*, 1990, 346, 763-766.
81. Lewis, E. B., *Am. Naturalist*, 1954, 88, 225-239.
82. Lewis, E. B., *Am. Naturalist*, 1955, 89, 73-89.
83. Bender, W., Weiffenbach, B., Karch, F. and Peifer, M., *Cold Spring Harbor Symp. Quant. Biol.*, 1985, 1, 173-180.
84. Nickol, J. M. and Felsenfeld, G., *Proc. Natl. Acad. Sci. USA*, 1988, 85, 2548-2552.
85. Choi, O. R. B. and Engel, J. D., *Cell*, 1988, 55, 17-26.
86. Chisaka, O. and Capecchi, M. R., *Nature*, 1991, 350, 473-479.
87. Balling, R., Mütter, G., Gruss, P. and Kessel, M., *Cell*, 1989, 58, 337-347.
88. Kessel, M., Balling, R. and Gruss, P., *Cell*, 1990, 61, 301-308.
89. Blatt, C., Aberdam, D., Schwartz, R. and Sachs, L., *EMBO J.*, 1988, 7, 4283-4290.
90. Kongsuwan, K., Allen, J. and Adams, J. M., *Nucleic Acids Res.*, 1989, 17, 1881-1892.
91. Blatt, C. and Sachs, L., *Biochem. Biophys. Res. Commun.*, 1988, 156, 1265-1270.

ACKNOWLEDGEMENTS. This study was supported in part by research grant HD06331 from the National Institutes of Health, Bethesda, Md, and by grants from the American Cancer Society, Atlanta, Ga, the March of Dimes, White Plains, NY, and the Lucille P. Markey Charitable Trust, Miami, Fla.

I thank Susan Celniker, PhD, and Welcome Bender, PhD, for a critical reading of the manuscript and Peter Chang, James Hardwick, Helen Park, and Seema Sharma, for assistance in the preparation of it. While at Caltech, a number of colleagues have directly contributed to our research on the bithorax complex, namely, Welcome Bender, PhD, Marie-Paz Capdevila, PhD, Susan Celniker, PhD, Loring Craymer, PhD, Madeline Crosby, PhD, Ian Duncan, PhD, Antonio Garcia-Bellido, PhD, William Gelbart, PhD, Alain Ghysen, PhD, Lily Jan, PhD, Margit Lohs-Schardin, PhD, and Shige Sakonju, PhD. Finally, I want to stress the close cooperation that we have had over the years with Prof David Hogness, PhD, and his colleagues at Stanford University and Prof Welcome Bender, PhD, at Harvard Medical School. It was Professor Hogness' foresight to launch the molecular analysis of the bithorax complex in 1978 in his laboratory.

Even insects experience visual illusions

Mandyam V. Srinivasan

Although the compound eyes of insects appear very different from our own eyes, research is beginning to suggest that insects experience many of the illusions that we do, and that their visual systems may process information in ways akin to ours.

It is difficult not to be intrigued by the tricks that our eyes and brain sometimes play on our perceptions. Certain visual illusions are so powerful that they cry out for an explanation. Indeed, some of the better-known illusions (reviewed, for example, by Gregory¹ and Goldstein²) have attracted nearly as many explanations as explainers!

Illusions are more than mere curiosities. They can illuminate the ways in which information is processed by the visual pathways. Recent years have witnessed mounting evidence that certain illusions are experienced not only by humans, but also by creatures such as insects, which possess relatively simple visual systems. This article describes some illusory percepts that seem

to be common to man and insect, and discusses the implications of their universality.

The 'waterfall' illusion

Many of us would have experienced that, upon staring at the water cascading down a waterfall for about a minute and then looking at the surrounding landscape, the stationary landscape appears to move distinctly *upwards* for a few seconds². This illusory percept of movement in the opposite direction can also be elicited in a contemporary urban setting by watching lines of text being scrolled on a computer screen. Curiously, insects behave as though they experience this illusion as well, and experiments with them reveal a possible neurophysiological explanation.

The perception of movement by a fly can be monitored by measuring its response to a moving, striped pattern: the insect tends to turn in the same

Mandyam V. Srinivasan is in the Centre for Visual Science, Research School of Biological Sciences, Australian National University, P. B. Box 475, Canberra, A. C. T. 0200, Australia