1D quasicrystals with the aid of curved spaces (H. Terrones, Mexico) and multifractals (G. Ananthakrishna) attracted attention. The recent advances in clusters and their description were summarized by Kumar, Italy. The last session at the end of the third day comprised nine short presentations, including an impromptu presentation by R. Prasad on the occurrence of amorphous lead in tubular fullerenes. All the participants

were impressed by the depth and breadth of presentations made in the area of quasicrystals and brought to focus the many significant contributions to the field from India. The diversity in the topics as well as the specializations of the participants was noted by many and aptly summarized by H. Terrones in the quotation from Rig-Veda: Let noble thoughts come to us from all sides.

The discussion meeting started with a brief address by V. Krishnan, Head Academic Programme of the Jawaharlal Nehru Centre and S. Ranganathan on behalf of the convenors (R. Chidambaram, S. Ranganathan, K. Chattopadhyay and P. Ramachandra Rao).

P. Ramachandra Rao, National Metallurgical Laboratory, Jamshedpur.

SCIENTIFIC CORRESPONDENCE

An epidemic outbreak of *Panonychus ulmi* (Koch) (Acari: Tetranychidae) in apple orchards of Himachal Pradesh, India

In July 1991 apple orchards in the wet temperate high hills of Shimla and Kulu regions of Himachal Pradesh presented a scorched look due to an epidemic outbreak of the European red mite, Panonychus ulmi (Koch) (Acari: Tetranychidae).

P. ulmi has been variously referred to in the literature as Paratetranychus pilosus (Canestrini and Fanzago) and Metatetranychus ulmi (Koch)¹. Commonly known as the European red mite, P. ulmi is found in many parts of the world. It is a major pest of apple, pear and plum in Europe, USA and Japan². In India, P. ulmi has however been recorded from peach, plum, apple, wheat, fig, hibiscus, tomato-seedlings and ivy from Jammu and Kashmir and from apricot in Himachal Pradesh, but has not been reported to cause any serious damage³.

The dry spell of June-August 1991 provided favourable conditions for the proliferation of *P. ulmi*. When mean daily temperatures ranged from 23 to 27°C, the European red mite completed

its life cycle in 5-7 days. First larvae from overwintering eggs, made their appearance in early April on warming up of weather which also coincided with the pink bud stage of the host apple trees. High mite counts (465 mites per apple leaf) were observed from orchards presenting a yellow and bronzed look. These were the orchards with higher insecticide score and conspicuous absence of the natural enemies of P. ulmi. Similar observations have been made in the apple orchards of Nova Scotia⁴. The mite-infested trees, during 1992, were observed to form fewer flower buds than normal trees, the fruits were smaller and there was unprecedented fruit drop.

The European red mite overwintered in the egg stage. Numerous overwintering eggs were laid by the females on roughened bark or crevices near the growing points. The number of overwintering eggs per 10 cm length of the twig ranged from 8 to 260. The deposition of the overwintering eggs was observed till the middle of October on the apple trees located between 2000 and 2300 m msl.

- Pritchard, A. E. and Baker, E. W., Hilgardia, 1952, 21 (9), 253-289.
- Jeppson, L. R., Keifer, H. H. and Baker, E. W., Mites Injurious to Economic Plants, University of California Press, California, 1975, pp. 615.
- 3. Prasao, " Int. J. Acarol., 1975, 1(2), 18-25.
- Hardman, J. M., Rogers, R. E. L., Nyrop,
 J. P. and Frish, T., J. Econ. Entomol.,
 1991, 84(2), 570-580.

Acknowledgements. I thank Dr G. P. Channa Basavanna, Professor of Entomology (Emeritus), Chief Advisor AICRP (Agricultural Acarology) and Dr B. Mallik, Acarologist, University of Agricultural Sciences, Bangalore, for kindly confirming the identity of the mite.

Received 14 December 1992; revised accepted 22 March 1993

R. KUMAR O. P. BHALLA

Department of Entomology Univ. of Horticulture and Forestry Solan 173 230, India