

- 14 Ashtekar, A., *Lectures on Nonperturbative Canonical Gravity*, World Scientific, Singapore, 1991, pp 334 (Advanced Series in Astrophysics and Cosmology, vol 6)
- 15 Ashtekar, A., New Variables for Classical and Quantum Gravity, *Phys Rev Lett*, 1986, 57, 2244-2247
- 16 Smolin, L., Experimental signatures of quantum gravity, CGPG-95-3-2, Mar 1995, e-Print Archive gr-qc/9503027.
- 17 Rovelli C and Smolin, L., Spin networks and quantum gravity, CGPG-95-4-4, April 1995, 42 pp, e-Print Archive. gr-qc/9505006
- 18 Rovelli, C and Smolin, L., Discreteness of area and volume in quantum gravity, CGPG-94-11-1, Nov 1994., 36 pp, e-Print Archive gr-qc/9411005.
- 19 Ashtekar, A., Lewandowski, J., Donald, M., Jose, M. and Thiemann, T., Quantization of diffeomorphism-invariant theories of connections with local degrees of freedom, UCSBTH-95-7, Apr 1995, 71 pp, e-Print Archive gr-qc/9504018.
- 20 Ashtekar, A and Lewandowski, J., Differential geometry on the space of connections via graphs and projective limits, CGPG-94-12-04, Dec 1994, 53 pp e-Print Archive hep-th/9412073.
21. Ashtekar, A., Lewandowski, J, Donald, M, Jose, M and Thiemann, T., Coherent state transforms for spaces of connections, CGPG-94-12-02, Dec 1994, 38 pp, e-Print Archive gr-qc/9412014
- 22 Cumrun, V and Edward, W., A one loop test of string duality, HUTP-95-A015, May 1995, 13 pp, e-Print Archive: hep-th/9505053.
23. Schwarz, J H., Classical duality symmetries in two dimensions, CALT-68-1994, May 1995, 13 pp., Presented at STRINGS 95: Future Perspectives in String Theory, Los Angeles, CA, 13-18 March 1995, e-Print Archive: hep-th/9505170.
24. Schwarz, J. H., String theory symmetries CALT-68-1984, March 1995, 12 pp, presented at International Workshop on Planck Scale Physics, Puri, India, 12-21 December 1994, e-Print Archive hep-th/9503127.
25. Schwarz, J. H., Evidence for nonperturbative string symmetries, CALT-68-1965, July 1994, 9 pp, Conference on Topology, Strings and Integrable Models (Satellite Colloquium to the

ICMP-11 18-23 Jul 1994), Paris, France, 25-28 July 1994, e-Print Archive hep-th/9411178

Notes

- 1 Ontology is the part of metaphysics concerned with the nature of existence.
2. Stephen Hawking's *A Brief History of Time. A Reader's Companion*, prepared by Gene Stone, Bantam Books, 1992, p125
3. In SR, a foliation is a one-parameter family of three-dimensional flat surfaces (hyperplanes) which, taken all together, fill out the entire spacetime. In GR, these surfaces are curved
- 4 Dust as a standard of space and time in canonical quantum gravity By J. David Brown (North Carolina State University), Karel V Kuchar (Utah U), GRQC-9409001, Aug 1994 57 pp e-Print Archive gr-qc@xxx.lanl.gov - 9409001
- 5 Dust is a basic type of matter which experiences only gravity, and no other interaction.
6. *Physics Today*, 1991, 44, 36, *Decoherence*, Wojciech Zurek
- 7 See for instance, *Extremal Black Holes and Elementary String States*, Ashoke Sen (ICTP, Trieste and Tata Inst) TIFR-TH-95-19, Apr 1995. 16 pp. e-Print Archive hep-th/9504147.
8. *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*, Steven Weinberg, New York, Wiley, 1972, xxviii, 657 pp., ill, 23 cm
- 9 A single-valued function $f(z)$ is holomorphic (or regular or analytic) in a region if it is differentiable at each point of the region
- 10 Superstring Theory, *Phys. Rept*, 1982, 89, 223.

ACKNOWLEDGEMENTS. This article was based on interviews with Chris Isham, Ed Witten and Abhay Ashtekar William Spence at Queen Mary and Westfield College in London helped review the final draft. It was the basis for a theory seminar presented at Melbourne University, Australia.

Faint star counts and the Milky Way structure

Devendra Ojha

The Milky Way Galaxy offers a unique opportunity for testing theories of Galaxy formation and evolution. Here we discuss how large surveys, both photometric and astrometric, of galactic stars are the keystones in investigations into such fundamental problems as the merging history and future of the Galaxy. This work features a sample survey plan to produce probes of stellar populations in the Milky Way. Objectives of this work are to trace the fine structure of our Galaxy through the statistical study of the stellar distributions according to their luminosity, colours and proper motions. This involves two steps: first, acquiring a new photometric and astrometric sample survey in various galactic directions; and second analysing the data using a model of population synthesis and determining the properties of populations in the Galaxy and constraints on the scenario of formation and evolution.

THE field of galactic structure is currently extremely active, due mostly to improvements in observational capabilities and the realization that our Galaxy is perhaps the one best-suited to test theories of Galaxy

formation. The concept of stellar populations has proved to be one of the most useful ideas of modern astronomy. Most simply, it is the idea that we can define a population as a set of stars that possess shared characteristics such as composition, age or kinematics, and that we can use the properties of various stellar populations to de-

Devendra Ojha is in the Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune 411 007, India

termine the structure and evolutionary history of the Galaxy. The first modern definition of stellar populations was given by Baade¹ who defined two stellar populations – the now famous populations I and II – based on colour-magnitude diagrams and on the brightness of resolved stars in M31 and M32. Population I is composed of stars found in the disk of our Galaxy and in the Magellanic Clouds; a signature of this population is the presence of highly luminous O- and B-type stars and open clusters. Population II is made up of stars with colour-magnitude diagrams like galactic globular clusters, which lack luminous blue stars; another signature of population II is the presence of short-period Cepheid variables. Now it became clear that population II was old and poor in metals, while population I was metal-rich and contains both young and old stars. The various doubts and worries about the population I/II scheme came to a head in the famous Vatican Conference² of 1957, during which Baade's original two populations were replaced with five: extreme population I, older population I, disk population, intermediate population II, and halo population II. Since the Vatican Conference, the field of stellar populations has been advanced significantly, and we now have a wealth of detailed information on populations throughout the Galaxy. One of the main developments is that we now have a good picture of the intermediate population (thick disk) in our Galaxy, i.e. a component with characteristics intermediate to those of the thin disk and halo populations³.

Studies of stellar populations often take as their starting point large-scale surveys, with selection criteria involving one or more observable variables. The physical location of the stars is arguably the most important constraint. With these points in mind, we have carried out a sample survey in UBV photometry and proper motions in various directions in the Galaxy. Large Schmidt telescope plates are well suited for measuring photometric magnitudes and positions for a large number of stars, because they have a large field ($5^\circ \times 5^\circ$) and go sufficiently faint. By choosing a reasonable number of fields in various galactic directions, we expected to get a representative sample of the spatial distribution of the different stellar populations, which have led to an understanding of the history of formation and evolution of the whole Galaxy. The kinematical studies of stellar populations are very important because they follow the galactic dynamics and retain the memory of the evolution of the whole potential. The results discussed here concern the kinematics of thin disk and thick disk populations of our Galaxy and the scenario of formation for the thick disk population.

Observations and data reduction

The chosen directions constitute a complete set of fields at high and intermediate latitudes:

- Direction towards the galactic anticentre⁴ ($l = 167^\circ$, $b = 47^\circ$).
- Direction towards galactic centre⁵ ($l = 3^\circ$, $b = 47^\circ$) near the globular cluster M5.
- Direction towards the galactic antirotation⁶ ($l = 278^\circ$, $b = 47^\circ$).
- Direction towards the North Galactic Pole⁷ ($l = 58^\circ$, $b = 80^\circ$).

The basic observational material for the survey is Schmidt plates. The plates used in our programme have been taken from Palomar, Tautenburg, European Southern Observatory and Observatoire de la Côte d'Azur (France) Schmidt telescopes. For each field the plates have been taken in UBV filters. In all we have a total of 58 plates for the various fields. Each plate has been scanned with the MAMA (Machine Automatique à Mesurer pour l'Astronomie) machine at Paris Observatory. A sampling step and pixel size of $10 \mu\text{m}$ has been used to scan the plates. The output of the MAMA machine is a catalogue of objects giving the positions X and Y , integrated density and the area of each object. The integrated densities are used to calibrate the plates photometrically, while a combination of various Schmidt plates of a given field is used for proper motion determination involving a time base of ~ 34 years. To calibrate the plates photometrically a number of photometric standards are required. The standards should cover the entire range of magnitudes and colours to be studied. To obtain photometric standards we have observed a number of subfields in each field using the CCD system attached to the 1-m Sampuranand telescope of the UP State Observatory at Nainital, India. A few standards have also been observed using the 1.2-m telescope of Observatoire de Haute-Provence in France. Our final photometric and astrometric survey contains 55,000 stars covering a 50 square degree field.

Kinematical analysis

The distance of each star in the catalogue was determined by estimating the absolute magnitude, which was obtained from a M_V vs $B-V$ relation of main-sequence stars of solar metallicity, taking into account the metallicity change as a function of the distance from the galactic plane. The cardinal components of the stellar space velocity (in km/s), U , V and W (where U is defined as positive in the direction of the galactic anticentre, V is positive in the direction of galactic rotation, and W is positive in the direction of NGP), were derived from proper motions, μ_l and μ_b (in arcsec per year) and line of sight distance d (in pc)⁸.

The major factors contributing to distance errors are the photometric errors in the colour of σ_{B-V} and magnitude σ_V . From our surveys, we expect the mean errors in

$\sigma_{R,V} = 0.1$ and $\sigma_V = 0.07$ for $V = 11-18$. From these values, our estimate of error of about 20% in distance seems to be realistic up to $z = 3$ kpc, where z is the distance above the galactic plane.

Kinematics of stellar populations

The kinematical properties of each stellar population are related to their spatial distributions. Scale height, velocity dispersions and asymmetric drift are linked by the Boltzmann equation. The proportion of each stellar population varies with the distance above the galactic plane and the selection of a stellar sample at a given distance allows one to optimize the proportion of one population. Since kinematical data allow one to improve this identification of populations, we have minimized bias from mutual contamination of each population at a given height by performing a kinematical separation of populations. To perform the kinematical separation, we have used a maximum-likelihood method (Stochastic Estimation Maximization (SEM) algorithm)⁹ in order to deconvolve the multivariate Gaussian distributions and estimate the corresponding parameters. The aim of the SEM algorithm is to resolve the finite mixture density estimation problem under the maximum-likelihood approach using a probabilistic teacher step. Through SEM one can obtain the number of components of the Gaussian mixture (without any assumption on this number), its mean values, dispersions and the percentage of each component with respect to the whole sample. We have used this method to separate the 2-D Gaussian velocity distributions (U, V) to identify the two components (thin disk and thick disk) of the Galaxy¹⁰. A multivariate discriminant analysis¹¹ is also used to distinguish the thick disk from other populations with the help of the model of population synthesis¹²⁻¹⁴.

Thin disk population

For the thin disk population, we observe a continuous increase of velocity dispersion with height^{4,5}. It can be explained by the fact that the thin disk is a mixture of disks with various scale heights and velocity dispersions, since the disk is not formed in isothermal components. This results in a vertical gradient of velocity dispersions, since the disk is not resolved in isothermal components. For the thin disk population, we obtain a measure of the kinematic gradients, which appear to be

$$\frac{\partial \ln \sigma_{U,W}^2}{\partial R} = -0.18 \pm 0.03 \text{ kpc}^{-1}$$

and

$$\frac{\partial \ln \sigma_V^2}{\partial R} = -0.05 \pm 0.02 \text{ kpc}^{-1}.$$

R is the galactocentric distance projected upon the galactic plane. Our results^{10,11} confirm that the thin disk has a relatively short scale length of -2.3 ± 0.5 kpc and scale height of -258 ± 50 pc.

Thick disk population and the merging history of the Milky Way

The thick disk population has been revisited under the light of these new data. We used also other photometric and astrometric sample survey in complementing direction, like the pole⁷.

The sample of stars in two fields (galactic centre and anticentre) have been divided into 6 or 7 bins of distance, and in each bin of distance a fit has been performed with a SEM algorithm to identify the Gaussian (U, V) components corresponding to the different populations^{4,5}. The thick disk population has been identified as a discrete and distinct component. In our present study¹¹, the thick disk population appears to be isothermal, since the kinematical characteristics are nearly constant along the line of sight and so that there is no vertical gradient. For this reason, the velocity characteristics for this population are much better defined. For the thick disk population, the kinematic radial gradients appear to be almost zero:

$$\frac{\partial \ln \sigma_{U+W}^2}{\partial R} = -0.008 \pm 0.01 \text{ kpc}^{-1}$$

and

$$\frac{\partial \ln \sigma_V^2}{\partial R} = +0.02 \pm 0.02 \text{ kpc}^{-1}.$$

By combining the kinematical results deduced from our four surveys (galactic centre, anticentre, antirotation, and NGP), we have derived the velocity ellipsoid of the thick disk population¹¹. The mean kinematic parameters are: $(\sigma_U, \sigma_V, \sigma_W, V_{\text{lag}}) = (72 \pm 4, 54 \pm 3, 40 \pm 3, 40 \pm 10)$ km/s. The most probable values of scale height and local normalization for the thick disk component are determined to be $h_z = 759 \pm 50$ pc and $A_{\text{thick}} = 7.4_{-1.5}^{+2.5}\%$ of the disk. The ratio of the number of thick disk stars in the galactic centre region to that in the centre region yields $h_R = 2.8-3.7$ kpc for the scale length of thick disk¹¹.

These properties have been compared with several scenarios of formation. 'Top-down' models, where the thick disk forms through a dissipational collapse, after the halo formation and before the thin disk has completely settled down, are in contradiction with the data, since they infer a continuity in the characteristics of the thick disk and thin disk and large kinematical vertical gradients. Alternatively, 'bottom-up' models suppose a formation of a thick disk after the complete collapse of the thin disk. We can eliminate those where the thick

disk formed from a secular kinematic diffusion of thin disk stars (which would produce a continuity between thin and thick disks). It remains a scenario well in agreement with the present data: the thick disk could have been formed from the dynamical heating of the thin disk during the sink of a small galaxy into the Milky Way. This merging event has to happen at the beginning of the thin disk life time so that the gas can cool again and form stars in the long-lasting thin galactic disk that we see now. This violent bottom-up scenario leaves two important observational signatures. First the thick disk is a separate population distinct from the thin disk and the halo. Second, no gradient can be generated in the thick disk by the event, although a preexisting gradient may survive the merger.

Conclusions

The thick disk population is found from all points of view (density laws, kinematics and metallicities) as a population well separated from the thin disk. It shows no gradient either on abundance³ or on kinematics. The results emerging from the present study of the correlations between photometry and kinematics give a mounting evidence that the thick disk of the Galaxy is most likely a sequel of a dwarf satellite galaxy merging in the Milky Way disk during the early epoch of this disk. This interpretation emerged from an accurate characterization of the thick disk properties: the scale height and the scale length of this component have been established; its rotation and velocity dispersion turned

out to be quite distinct from both the disk itself and the halo. The local density of the thick disk component is twice what was previously assumed and the stellar colours do not reflect any significant chemical gradient.

- 1 Baade, W, *Astrophys. J.*, 1944, **100**, 137–146.
- 2 Blaauw, A, in *Galactic Structure* (eds Blaauw, A. and Schmidt, M), Univ. of Chicago Press, Chicago, 1965, pp 435–453
- 3 Robin, A. C., Haywood, M, Crézé, M, Ojha, D. K. and Bienaymé, O, *Astron. Astrophys.*, 1995 (in press)
- 4 Ojha, D. K., Bienaymé, O, Robin, A. C. and Mohan, V, *Astron Astrophys*, 1994, **284**, 810–826.
- 5 Ojha, D. K., Bienaymé, O., Robin, A. C. and Mohan, V, *Astron Astrophys*, 1994, **290**, 771–784
- 6 Ojha, D. K., Bienaymé, O., Mohan, V., Haywood, M and Robin, A. C., *Astron Astrophys*, 1995 (in preparation)
- 7 Soubiran, C, *Astron. Astrophys*, 1992, **259**, 394–403.
- 8 Murray, C. A., *Vectorial Astronomy*, Adam Hilger, Bristol, 1983, pp 271–311.
- 9 Celeux, G and Diebolt, J., *Rev. Statist. Appl.*, 1986, **34**, 35–52.
- 10 Ojha, D. K., Ph D thesis, Strasbourg University, France, 1994, pp 75–83
- 11 Ojha, D. K., Bienaymé, O., Robin, A. C., Crézé, M and Mohan, V, *Astron Astrophys.*, 1995 (submitted)
- 12 Robin, A. C. and Crézé, M, *Astron Astrophys*, 1986, **157**, 71–90
- 13 Bienaymé, O., Robin, A. C. and Crézé, M., *Astron Astrophys*, 1987, **180**, 94–110.
- 14 Haywood, M, Ph D thesis, Paris Observatory, France, 1994, pp 28–69.

ACKNOWLEDGEMENTS I thank Drs O. Bienaymé, A. C. Robin, M. Crézé, and V. Mohan for useful suggestions and helpful discussions during the course of this work. Financial assistance from IFCPAR, New Delhi, is also thankfully acknowledged.

REVIEW ARTICLE

A unique and remarkable binary pulsar

Simon Johnston

Research Centre for Theoretical Astrophysics, University of Sydney, NSW 2006, Australia

PSR B1259–63 is one of the most remarkable pulsars ever discovered. It is the only pulsar known to be orbiting a main-sequence Be star and forms the ‘missing link’ in the evolutionary scenario of binary stars. Its orbit is highly elliptical and every 3.5 yr it approaches to within 0.5 AU of its companion star. Its pulse profile is unique among pulsars and shows that the opening angle of its emission cone exceeds 180°. Observations made around the time of closest approach of the two

stars show changes in the pulsar’s dispersion measure, rotation measure and fractional linear polarization on very short timescales. The pulsar is observed to spin down during the periastron passage. The companion star to the pulsar has been monitored at optical and UV wavelengths. The system is a radio and X-ray transient and has been detected at γ -ray energies, with the high-energy emission probably occurring in a shock front formed by the collision of the two stellar winds.

In this review paper, I will discuss the remarkable binary pulsar PSR B1259–63. I will start with a general introduction into pulsars and binary systems before moving on to the specifics of the system containing PSR

B1259–63. There are an increasing number of excellent conference proceedings and review articles on pulsars in general with varying degrees of complexity. Interested readers should consult refs. 1–11 and the references therein.