

4. Usher, M. B., Wildlife conservation evaluation; Attributes, criteria and values. in *Wildlife Conservation Evaluation* (ed. Usher, M. B.), Chapman and Hall, 1986, pp. 3–44.
5. Huston, M. A., *Biological Diversity: The Coexistence of Species in Changing Landscapes*, Cambridge University Press, 1994.
6. Thothathri Banerjee, S. P., Hazra, P. K. and Pal, G. D., On a collection of *Psilotum* from the Great Nicobar Island. *Bull. Bot. Surv. India*, 1970, **12**, 279–281.
7. Rao, P. S. N., A distributional note on *Psilotum* Sw. (Psilotaceae) in Andaman and Nicobar Island. *J. Andaman Sci. Assoc., Port Blair*, 1990, **6**, 155–156.
8. Champion, H. G. and Seth, S. K., *A Revised Survey of the Forest Types in India*, Govt. of India Press, Nasik, 1968.

ACKNOWLEDGEMENTS. We thank Dr P. K. Hajra, ex-director Botanical Survey of India for identification of the specimen under study and also for his constant encouragement and support. We are all thankful to Shri Yesudas, Assistant Conservator of Forests, Great Nicobar for his timely support. Thanks are also due to Shri Rishikesh, Department of Science and Technology, Port Blair for providing necessary support.

Received 27 November 2002; revised accepted 10 February 2003

## Phenotypic plasticity and plant invasiveness: Case study of congress grass

C. Annapurna and J. S. Singh\*

Department of Botany, Banaras Hindu University, Varanasi 221 005, India

**This paper highlights the phenotypic variation of congress grass (*Parthenium hysterophorus*) to varying soil texture. Plants demonstrated significant differences in some traits among the different soil types and these phenotypically plastic traits, in combination, contribute to invasiveness of the species, allowing for the range expansion observed in different habitats.**

MANY organisms can respond with considerable flexibility to a changing environment, generating a range of characteristics depending on the environment in which the developing organism finds itself<sup>1</sup>. A new species, when introduced into an alien environment, must become established within the constraints of physical site characteristics and competition with resident vegetation<sup>2</sup>. The ability of an organism to express different phenotypes in response to environmental cues, i.e. phenotypic plasticity<sup>3</sup>, is generally interpreted as an adaptation for dealing with a variable environment. The primary benefit of being phenotypically plastic as an invading taxa lies in

the ability of plasticity for traits that contribute directly to fitness<sup>4</sup> and broaden the niche width of the population, and therefore, its range of potentially available resources<sup>5</sup>.

The success of an alien species depends on the degree of invasiveness, i.e. the potentiality to establish and spread<sup>6</sup>. A few simple biological attributes can be strong predictors of potential invasiveness of a species<sup>6,7</sup>. Plant height, relative growth rate (RGR) and seed mass, as indicators of both the establishment and regenerative phases of the life cycle<sup>8</sup> are the most relevant traits that address competitive ability<sup>9–11</sup>.

The nature of soil, associated with soil attributes such as texture, organic matter, pH and bulk density<sup>12</sup>, is aptly known to be a potent determinant of plant adaptation and distribution<sup>13</sup>. Soil texture is a surrogate index of soil quality. It influences organic matter accumulation<sup>14</sup>, distribution of soil N (ref. 15) in association with topography and also independently<sup>16</sup>, and the dynamics of soil water<sup>17</sup> that most frequently limits the biological processes in semi-arid regions<sup>18</sup>.

We hypothesize that the colonizing ability of an invader is a function of plasticity for fitness-related traits, enabling it to cope with and perhaps to benefit from habitat conditions. In order to test this hypothesis, we evaluated the response of a well-known invader species, *Parthenium hysterophorus* (congress grass; Asteraceae) in terms of plant height, RGR and seed mass, when cultured in soils differing in texture. *P. hysterophorus*, native of tropical America, is a rapidly colonizing weed and a health hazard<sup>19</sup>, especially in the tropics. In India, it was first recorded from Pune, Maharashtra<sup>20</sup>. During the last 30 years, it has spread alarmingly to virtually every state in the Indian territory, occupying vast areas of waste- and cultivated lands.

A pot culture experiment in a completely randomized design, was established in the Botanical Garden, Banaras Hindu University (BHU), Varanasi (25°18'N, 83°03'E, 129 m asl), India. The proportions of clay and sand were varied by thoroughly mixing the S<sub>1</sub> soil (collected from subsoil layer, below 25 cm of a pond) and the S<sub>5</sub> soil, nearly pure sand (collected from a river bank) to create homogenous soil mixtures (in 3 : 1, S<sub>2</sub> and 1 : 3, S<sub>3</sub>). An alluvial garden soil (Inceptisol) was the fifth soil type (S<sub>4</sub>). Sand, silt and clay fractions in the soil mixtures were determined by the hydrometer method<sup>21</sup> (Table 1). Thus five treatments reflecting a gradient of decreasing clay content (%), 14 (S<sub>1</sub>), 10 (S<sub>2</sub>), 3 (S<sub>3</sub>), 2 (S<sub>4</sub>), and 0 (S<sub>5</sub>), were created.

For seed germination trial, fresh *P. hysterophorus* achenes (hereafter referred to as seeds) were collected from a single local population of plants of uniform height. The seeds were stored for six weeks in the dark at room temperature until the experiments commenced. On 25 May 2000, 45 seeds in each treatment (three replicates per treatment) were placed into earthen pots (2.87 l) filled with respective soil mixtures, moistened to field

\*For correspondence. (e-mail: jssingh@bhu.ac.in)

## RESEARCH COMMUNICATIONS

capacity. Per cent germination was calculated on the daily observations (till no more seed germinated). The length of seedlings was measured ten days after the germination.

On 21 June 2000, two-week-old seedlings (with a biomass equivalent to 40 mg dry wt per seedling), selected for uniformity, were transplanted (one per pot) into 15 earthen pots (11.22 l) per treatment containing the respective soil mixtures. The pots were irrigated every alternate day. To determine the influence of treatments on *P. hysterophorus*, three plants (per treatment) were destructively harvested at vegetative stage (29 days after transplantation) and three (per treatment) at mature stage (during flowering prior to achene dehiscence). At each harvest, after measuring the height, plant parts were separated and oven-dried at 80°C to constant weight. RGR ( $\text{mg g}^{-1} \text{day}^{-1}$ ) was calculated as follows<sup>22</sup>:

$$\text{RGR} = \frac{\ln(W_2) - \ln(W_1)}{T_2 - T_1},$$

where  $W_1$  is the dry weight of two-week-old (i.e.  $T_1$ ) seedlings and  $W_2$  is the dry weight at 29 days after transplantation in the case of juveniles, and that at the flowering stage prior to achene dehiscence in the case of post juveniles (i.e.  $T_2$ ), and  $T_2 - T_1$  is the sampling interval in days.

Reproductive growth was studied by measuring seed production (mean number of seeds per capitulum  $\times$  the number of capitula per plant) and mean seed dry mass (80°C), in each of the remaining nine individuals (out of 15) per treatment (except  $S_1$ , where data were based on merely three plants which survived till seed production). Reproductive effort (RE) was estimated as the ratio of number of seeds per gram above-ground dry weight<sup>23</sup>.

Effect of treatment on growth variables was analysed by one-way ANOVA. Differences between treatment means were tested by Tukey's HSD test (at  $P < 0.05$ ). Linear regression was used wherever necessary. All statistical analyses were conducted using the SPSS statistical package (SPSS Inc., Chicago, USA).


**Table 1.** Textural properties of soil mixtures

Treatment	Proportion (%)		
	Clay	Silt	Sand
$S_1$	14a	62a	24a
$S_2$	10b	62a	28a
$S_3$	3c	21b	76b
$S_4$	2c	73c	25a
$S_5$	0d	1d	99c

Values in a column with different letters are significantly different from each other according to Tukey's HSD test at  $P < 0.05$ .


Seed germination ( $F_{4,10} = 90.22$ ;  $P < 0.001$ ) and seedling length ( $F_{4,10} = 7.04$ ;  $P = 0.006$ ) were significantly affected by the clay content. Germination was greatest (62%) in  $S_4$  (i.e. 2% clay) and was significantly reduced in soils above and below this clay content (Figure 1a). The lowest germination (9%) occurred in  $S_1$ . However, germination was 64 and 39% lower, respectively in  $S_2$  and  $S_5$ , compared to  $S_4$  treatment. Further, differences in seedling length were not significant among treatments except for  $S_5$ , where it was only half as that in  $S_4$  (Figure 1b).

*P. hysterophorus* exhibited substantial phenotypic variability in plant traits in response to soil quality (see ref.



**Figure 1.** Effect of clay content on per cent seed germination (a) and seedling length after 10 days (b) of *P. hysterophorus*.

24 for additional data). Trends in plant height and RGR were similar at vegetative and reproductive stages (Figure 2a and b), where the peak was attained in S<sub>3</sub> treatment. Increased clay content (i.e. >3%) led to consistent decline in these estimates. ANOVA indicated that plant height differed between the vegetative and reproductive stages ( $F_{1,20} = 12174.33$ ;  $P < 0.001$ ) and among treatments ( $F_{4,20} = 324.06$ ;  $P < 0.001$ ). RGR also differed between the vegetative and post-vegetative stages ( $F_{1,20} = 15048.73$ ;  $P < 0.001$ ) and was significantly affected by treatment ( $F_{4,20} = 1150.01$ ;  $P < 0.001$ ). The stage  $\times$  treatment interaction was highly significant, indicating differential responses of the variables. Juveniles were 52 and 39% shorter, respectively in S<sub>1</sub> and S<sub>5</sub> than those in S<sub>3</sub>.



**Figure 2.** Effect of clay content on plant height (a) and relative growth rate (RGR) (b) of *P. hysterophorus*.

Similarly, RGR during the vegetative stage was 35 and 27% lower in S<sub>1</sub> and S<sub>5</sub> treatments respectively, compared to S<sub>3</sub>. Plant height at mature stage in S<sub>1</sub> and S<sub>5</sub> was respectively 30 and 20% lower than in S<sub>3</sub>. Correspondingly, post-juvenile plants of S<sub>1</sub> and S<sub>5</sub> had 56 and 39% lower RGR respectively, compared to those grown on S<sub>3</sub>.

HSD analyses showed that the above-mentioned traits differed significantly among treatments, except that plant height in juveniles did not vary among S<sub>2</sub>, S<sub>4</sub> and S<sub>5</sub>, and in adults between S<sub>1</sub> and S<sub>2</sub>, and S<sub>3</sub> and S<sub>4</sub>. RGR did not differ significantly between S<sub>2</sub> and S<sub>4</sub> treatments in juveniles and that between S<sub>3</sub> and S<sub>4</sub> in post-juvenile plants. However, RGR was 14% lower ( $P < 0.05$ ) in S<sub>4</sub> compared to S<sub>3</sub> during the vegetative stage. Invariably higher RGR in juveniles than mature plants suggested that greater pace of growth during early phases of the life cycle would confer an advantage to the establishing plant.

There were significant treatment effects on plant traits for RE ( $F_{4,10} = 16.71$ ;  $P < 0.001$ ) and seed mass ( $F_{4,10} = 19.03$ ;  $P < 0.001$ ). RE increased up to S<sub>4</sub> and then declined significantly as clay content increased (Figure 3a). In contrast, with increasing clay content, seed mass continuously increased (Figure 3b). These traits were 50% lower in S<sub>5</sub> compared to S<sub>4</sub>.

According to Bazzaz<sup>25</sup>, availability of soil resources, especially nutrients, critically influenced plant growth, population dynamics and competitive interactions. We expect that the amount of plant-available N could be low in soils high in clay content where restricted aeration<sup>26</sup> may increase denitrification losses<sup>27</sup> and reduce mineralization of organic N (ref. 28). Siemens *et al.*<sup>29</sup> reported that limited resources (e.g. soil nutrients, water) can directly inhibit the rate of growth. Hence, low RGR exhibited on clay-rich soils (S<sub>1</sub> and S<sub>2</sub> treatments) suggests that *P. hysterophorus* would maintain slower growth rates under nutrient-poor habitats as a survival strategy. Further, inadequate contact of roots with the soil in sands<sup>30</sup> could limit the uptake of water and nutrients, which in turn, appears to reduce the seedling length and growth rates in *P. hysterophorus*.

On the other hand, high RGR observed in S<sub>3</sub> and S<sub>4</sub> treatments would enable the plant to pre-empt growth resources. The fact that height is an indicator of relative size may explain the greater ability of larger plants to extract resources<sup>31</sup> from these soils. Consequently, *P. hysterophorus* can take advantage of resource abundance and grow more rapidly than its neighbouring plants. This trade-off is thought to be a necessary condition for the persistence of parthenium weed across fertility gradients.

Our observation that RE decreased with high clay content (>3%) agrees with the prediction that in environments where juvenile mortality is high, RE should be low<sup>32</sup>. In the S<sub>1</sub> treatment (with 14% clay), two-thirds of transplanted plants died during the vegetative stage (data not shown). Further, juveniles established successfully with comparatively higher RE on coarser soils. Plasticity


in terms of texture-induced variability in RE may suggest that *P. hysterophorus* could colonize under conditions of both high density-independent mortality (e.g. in open, disturbed habitats) and density-dependent regulation (e.g. in closed, less disturbed communities). The plant height in adults was positively related to RE ( $r^2 = 0.70$ ;  $P < 0.001$ ) indicating the influence of plant size on seed production and proportion of energy allocated to seeds. Thus, large plants seem to be at a reproductive advantage<sup>33</sup>.

Light seeds are more dispersible<sup>34</sup> compared to heavier seeds<sup>35</sup> and production of numerous light seeds in *P. hysterophorus* grown on coarse-textured soils may thus facilitate colonization of new sites. On the other hand, the less dispersible heavier seeds produced in clay-enriched

**Table 2.** Selected habitat factors and performance of *P. hysterophorus* in two field microsites

	Shade and moist	Open and moist
Clay content (%)	1.4a	2.1b
Sand content (%)	58.0a	53.0b
Soil C (mg g <sup>-1</sup> )	10.6a	6.7b
Soil N (mg g <sup>-1</sup> )	0.7a	0.6b
Plant height (cm)	119.5a	96.0b
Number of seeds (plant <sup>-1</sup> )	29367.0a	19254.0b
Seed mass (mg seed <sup>-1</sup> )	0.31a	0.5b
Density (plant m <sup>-2</sup> )	970.3a	866.3b

Values in a row with different letters are significantly different from each other according to 't' test at  $P < 0.05$ .



**Figure 3.** Effect of clay content on reproductive effort (RE) (a) and seed mass (b) of *P. hysterophorus*.

soil will likely form a persistent seed bank as argued by Navie *et al.*<sup>36</sup>. We found that soils high in clay content reduced the germination of *P. hysterophorus* seeds, probably because of restricted aeration<sup>26</sup>. The production of larger seeds (with greater reserves), although fewer in number, in clay-rich soils would confer an added benefit to this invasive species. In frequently disturbed areas where chances of seed burial are more<sup>36</sup>, larger seeds may ensure the long-term persistence of *P. hysterophorus* through emergence from greater depths of burial<sup>37</sup>. These observations support that the availability of seed reserves can have important impact on seedling establishment<sup>11</sup> and thus, persistence or invasiveness of a species.

This study has revealed that plasticity in plant traits due to soil quality led to two contrasting strategies which make *P. hysterophorus* a successful invader. The first resulted in tall, fast-growing competitors with small seed mass, appropriate for a rapid population expansion. The second strategy yielded short plants with high seed mass for persistence in an otherwise less favourable habitat leading to a slow build-up of population, with a gradual increase in the size of the seed bank.

*P. hysterophorus* has been reported from a wide variety of habitats<sup>19</sup>. Because of the inherent plasticity, this weed can rely on phenotypically varying plant traits to exploit the available 'invasion windows' (conditions suitable for establishment, growth, and proliferation of invaders, see ref. 38). We believe that the ubiquity of heterogeneity in natural habitats<sup>39</sup> might create a mosaic of microsites, with populations of relatively taller plants growing in the more favourable spots and those of smaller plants in less favourable ones. This is illustrated in Table 2 from two habitat microsites located within the BHU campus. Management practices should focus on limiting seed dispersal from taller populations and preventing the build-up of persistent seed bank in shorter populations in clay-rich soils.

We may point out however, that in addition to variable soil conditions, the presence of other species in natural habitats may also elicit a plastic response by influencing

the growth performance of *P. hysterophorus* through competitive and other indirect effects.

1. Dusheck, J., *Nature*, 2002, **418**, 578–579.
2. Parshall, T., *Ecology*, 2002, **83**, 1386–1398.
3. Sultan, S. E., *Evol. Biol.*, 1987, **21**, 127–176.
4. Travis, J., in *Ecological Genetics* (ed. Real, L.), Princeton University Press, Princeton, New Jersey, USA, 1994, pp. 171–204.
5. Sultan, S. E. and Bazzaz, F. A., *Evolution*, 1993, **47**, 1050–1071.
6. Rejmanek, M. and Richardson, D. M., *Ecology*, 1996, **77**, 1655–1661.
7. Rejmanek, M., *Aust. Ecol.*, 2000, **25**, 497–506.
8. Hodgkinson, D. J. and Thompson, K., *J. Appl. Ecol.*, 1997, **34**, 1484–1496.
9. Westoby, M., *Plant Soil*, 1998, **199**, 213–227.
10. Hodgson, J. G., Wilson, P. J., Hunt, R., Grime, J. P. and Thompson, K., *Oikos*, 1999, **85**, 282–296.
11. Leishman, M. R., *New Phytol.*, 1999, **141**, 487–496.
12. Gregorich, E. G., Carter, M. R., Angers, D. A., Monreal, C. M. and Ellert, B. H., *Can. J. Soil Sci.*, 1994, **74**, 367–385.
13. Epstein, E., *Mineral Nutrition of Plants: Principles and Perspectives*, John Wiley, New York, 1972.
14. Hassink, J., *Soil Sci. Soc. Am. J.*, 1996, **60**, 487–491.
15. Hook, P. B. and Burke, I. C., *Ecology*, 2000, **81**, 2686–2703.
16. Raghubanshi, A. S., *Soil Biol. Biochem.*, 1992, **24**, 145–150.
17. Lauenroth, W. K. and Milchunas, D. G., in *Natural Grasslands I: Introduction and Western Hemisphere* (ed. Coupland, R. T.), Elsevier, New York, 1992, pp. 183–226.
18. Singh, J. S., Milchunas, D. G. and Lauenroth, W. K., *Plant Ecol.*, 1998, **134**, 77–89.
19. Navie, S. C., McFadyen, R. E., Panetta, F. D. and Adkins, S. W., *Plant Prot. Q.*, 1996, **11**, 76–88.
20. Rao, R. S., *J. Bombay Nat. Hist. Soc.*, 1956, **54**, 218–220.
21. Bowles, J. E., *Engineering Properties of Soils and their Measurement*, McGraw-Hill International Editions, Civil Engineering Series, McGraw-Hill, Singapore, 1988, 3rd edn.
22. Evans, G. C., *The Quantitative Analysis of Plant Growth*, University of California Press, Berkeley, 1972.
23. Regehr, D. L. and Bazzaz, F. A., *J. Ecol.*, 1979, **67**, 923–933.
24. Annapurna, C. and Singh, J. S., *Weed Res.*, 2003, **43**, 190–198.
25. Bazzaz, F. A., *Plants in Changing Environments, Linking Physiological, Population, and Community Ecology*, Cambridge University Press, Cambridge, 1996.
26. Brady, N. C., *The Nature and Properties of Soils*, Macmillan, New York, USA, 1990, 10th edn.
27. Hansen, S., Maehlum, J. E. and Bakken, L. R., *Soil Biol. Biochem.*, 1993, **25**, 621–630.
28. Eijsackers, H. and Zehnder, A. J. B., *Biogeochemistry*, 1990, **11**, 153–174.
29. Siemens, D. H., Garner, S. H., Mitchell-Olds, T. and Callaway, R. M., *Ecology*, 2002, **83**, 505–517.
30. Passioura, J. B., *Aust. J. Soil Res.*, 1991, **29**, 717–728.
31. Niklas, K. J., *Evol. Trends Plants*, 1993, **7**, 42–48.
32. Bazzaz, F. A. and Reekie, E. G., in *Studies on Plant Demography: A Festschrift for John L. Harper* (ed. White, J.), Academic Press, London, 1985, pp. 373–387.
33. Ollerton, J. and Lack, A., *Ecology*, 1998, **139**, 35–47.
34. Weiher, E., van der Werf, A., Thompson, K., Roderick, M., Garnier, E. and Eriksson, O., *J. Veg. Sci.*, 1999, **10**, 609–620.
35. Fenner, M., *Seed Ecology*, Chapman and Hall, London, 1985.
36. Navie, S. C., Panetta, F. D., McFadyen, R. E. and Adkins, S. W., *Weed Res.*, 1998, **38**, 335–341.
37. Baskin, C. C. and Baskin, J. M., *Seeds: Ecology, Biogeography, and Evolution of Dormancy and Germination*, Academic Press, New York, 1998.

38. Johnstone, I. M., *Biol. Rev.*, 1986, **61**, 369–394.
39. Jackson, R. B. and Caldwell, M. M., *Ecology*, 1993, **84**, 891–903.

ACKNOWLEDGEMENTS. C.A. is thankful to CSIR, New Delhi for the award of a senior research fellowship.

Received 23 December 2002; revised accepted 11 March 2003

## Protective effect of vitamin A, ascorbic acid and $\alpha$ -tocopherol on 2,4-dimethylaminoazobenzene-induced hepatoma in rats

A. Antony Joseph Velanganni<sup>†</sup> and C. Balasundaram<sup>#,\*</sup>

<sup>†</sup>Department of Biochemistry, JJ College of Arts and Science, Pudukkottai 622 404, India

<sup>#</sup>Department of Animal Science, Bharathidasan University, Tiruchirappalli 620 024, India

**The chemoprotective effect of antioxidant vitamins (A, C and E) on chemically-induced hepatoma by 2,4-dimethylaminoazobenzene (DAB) in male albino rats has been studied. Group-I animals which were fed *ad libitum* with normal rat feed and water served as control. Each animal of the groups II–V received DAB injection intraperitoneally at a concentration of 20 mg/kg body wt once in a week. Immediately after administering the carcinogen, vitamins A (10,000 IU/kg body wt), C (250 mg/kg body wt) and E (400 mg/kg body wt) were given orally to groups III, IV and V, respectively and the experiment continued for a period of two months. Oral administration of antioxidant vitamins has a protective effect on the incidence of liver tumour monitored on the basis of liver weight, histological studies and enzymatic analysis. Administration of vitamins effectively protects the hyperplasia of the liver parenchymal cells and prevents damage of the nuclear envelope. Elevated levels of serum  $\alpha$ -glutamyl transpeptidase, acetylcholinesterase, GSH, ALP and bilirubin in the DAB-administered group were significantly reduced by administration of each vitamin.**

DIET is the second major cause for cancer and induces cancer of colon, breast, stomach, liver, etc. Certain foods like high-fiber low-fat diets and fresh fruits and vegetables provide protection against cancer<sup>1</sup>. Fresh fruits and vegetables are a rich source of antioxidant vitamins like A, C and E that prevent cellular damage associated with cancer incidence. The scavenging capacity of the antioxidant vitamins prevents oxidative damage by neutraliz-

\*For correspondence. (e-mail: bal213@rediffmail.com)