

Regulation of DNA topology in mycobacteria

V. Nagaraja

Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore 560 012 and Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore 560 064, India

DNA topoisomerases catalyse essential DNA transaction processes in order to attain the balanced topology of the genome. Contrasting activities of DNA topoisomerase I and DNA gyrase result in the maintenance of topological homeostasis. The regulation of expression of different topoisomerases ensure steady state optimum level expression of the enzymes. Many aspects of their organization and regulation seem to be different in mycobacteria when compared with that of *Escherichia coli*. Here we present several aspects of the regulation of mycobacterial topoisomerases and discuss the significance.

Introduction

DNA topoisomerases have evolved to catalyse the topological alterations in DNA in order to ensure that DNA-transaction processes are completed without topological interruptions. Hence the enzymes play essential roles during replication, transcription, recombination, repair and chromosome segregation¹. The genomes of bacteria are normally maintained in negative supercoiled state. Since all processes that involve DNA as a substrate either need to melt, bend or distort DNA; negative supercoiling modulates these cellular processes. For example, supercoiling influences recombination at two levels independently. First, supercoiling enhances recombination because the plectonemic winding of DNA facilitates the juxtapositioning of recombination sites and limits the extent of diffusion required for the sites to collide². Second, the extent of supercoiling of the substrate determines the complexity of the product(s) since recombination converts the supercoil nodes into nodes of catenation or knots, depending on the relative orientation of the sites³. In a complementary manner, various DNA transactions alter the topology of DNA. The most obvious of these being the generation of catenated daughter duplexes after replication and activities of DNA tracking machineries. In eubacteria, the principal enzymes that influence the vital processes are topoisomerase I and DNA gyrase with substantial contribution from topoisomerase IV when present (see later section). Hence, the regulation of their expression and activities is an important determinant in the maintenance of balanced topological state and the global supercoiling of DNA is thus dependent on the balance of activities of various topoisomerases.

Amongst the four topoisomerases found in *E. coli*, topoisomerase I and III belong to type IA group while DNA gyrase and topoisomerase IV are type II enzymes⁴. The key enzyme in all bacteria catalysing the formation of negatively supercoiled DNA in an ATP dependent reaction is DNA gyrase. The enzyme, encoded by *gyrA* and *gyrB*, is a heterotetrameric protein⁵. By virtue of its indispensability, the enzyme has been and continues to be a favourite drug target. As a consequence, several inhibitors and poisons, both natural and synthetic, have been characterized⁶. The second major player in influencing global topology is DNA topoisomerase I. In contrast to DNA gyrase, it comprises of a single polypeptide, encoded by *topA* gene. The enzyme catalyses the conversion of negatively supercoiled DNA into relaxed form in an ATP independent reaction⁴.

Unlike *E. coli*, where four topoisomerases have been characterized, mycobacteria and many other organisms do not encode the full complement of topoisomerases. This was evident during our efforts to clone the genes for DNA gyrase from both *M. smegmatis* and *M. tuberculosis*^{7,8}. Efforts to clone genes for other topoisomerases such as topoisomerase IV were unsuccessful hinting at the possibility of absence of these genes in *M. tuberculosis*. Genome sequencing efforts subsequently confirmed the presence of only single topoisomerase I and DNA gyrase in *M. tuberculosis*, while some other species such as *M. smegmatis*, *M. bovis* appear to have genes encoding for additional topoisomerases⁹⁻¹¹. Amongst the two type IA enzymes found in *E. coli* and other bacteria, only topoisomerase I is present in mycobacteria¹².

Figure 1 depicts the organization of genes encoding DNA gyrase in *E. coli*, *M. smegmatis* and *M. tuberculosis*. Notably, *gyrB* and *gyrA* in *E. coli* are located far apart in the circular chromosome but present next to each other in both the species of mycobacteria. Furthermore, significant additional differences are observed in their primary sequences¹³. The genetic linkage between the *gyr* genes seems to correlate with the size of the *gyrB* gene. Species in which the genes are present far apart have 165 amino acids extra in the C-terminal half of *GyrB*¹⁴ and this insertion appears to be involved in DNA binding¹⁵. Our work over the last decade has revealed some of the distinctive characteristics of mycobacterial topoisomerase organization, function and regulation. In the following sections, the salient features of topoisomerase regulation in mycobacteria are presented and compared with that of *E. coli*.

*For correspondence. (e-mail: vraj@mcbl.iisc.ernet.in)

Topology and transcription

Topological organization of DNA is known to have important influence in regulation of gene expression. Movement of RNA polymerase along the helical axis results in an increase in twist ahead of the tracking machinery and decrease behind, introducing positive and negative supercoils respectively. This is termed as twin domain of supercoiling¹⁶. As a result, DNA gyrase and topoisomerase I have to function ahead and behind the transcription bubble to remove positive and negative supercoils respectively (Figure 2). The twin domain model of supercoiling has several biological implications. (i) DNA transactions may prove to be a major determinant of local DNA topology; (ii) transcription of adjacent genes could significantly influence expression of a particular gene; (iii) Most importantly, for the first time, there appeared to be a necessity to expect efficiency in topoisomerases. Supercoiling influences transcription of many genes in the cell^{17–19}, modulating by several ways. Directly, it can realign promoter elements or facilitate open complex formation. Indirectly, it can stabilize loops, bends or other non-B-DNA structures in DNA.

In majority of the promoters, negative supercoiling facilitates isomerization of closed complex to open complex. However, failure to remove negative supercoils generated behind the transcription elongation complex would lead to the accumulation of R-loops and as a consequence, inhibition of transcription²⁰. In addition, study of regulation of topoisomerase expression in mycobacteria is important especially since the genome lacks full complement of topoisomerases. Furthermore, in many pathogenic bacteria, expression of virulence genes is

dependent on topological status of the genome^{21,22}. As sensor of supercoils, the topoisomerases influence the specific gene expression.

Transcription of topoisomerases and regulation of topology

The net supercoiling of intracellular DNA is maintained by the relaxation activities of DNA topoisomerases I and IV opposing the supercoiling activity of DNA gyrase. Thus, by modulating the expression of any one of these genes, the cell can bring about rapid changes in supercoiling as well as compensate for sudden changes in supercoiling. As the sole supercoiling activity in the cell, DNA gyrase faces the daunting task of opposing the relaxation activities of both topoisomerases I and IV²³ and regulates its expression by a unique mechanism termed relaxation-stimulated transcription (RST). In general, transcription of most genes is induced by increased negative supercoiling. In contrast, negative supercoiling represses transcription of the gyrase genes in *E. coli*²⁴. Increased gyrase levels lead to an increase in supercoiling, which, in turn, represses the expression of gyrase and allows other topoisomerases to bring the topology of the DNA back to its optimum state. Following observations led to the discovery of RST in *E. coli*. Cell-free transcription showed that transcription was dependent directly on the DNA topology, being maximal on a relaxed template²⁴. Deletion analysis of the promoter regions of both *gyrA* and *gyrB* genes defined a short region around the transcriptional start site, including the –10 region, that is necessary and sufficient for conferring RST to a reporter gene^{25,26}. Extensive mutagenesis of the




Figure 1. Transcriptional organization and regulation of *gyr* genes in *E. coli* and two species of mycobacteria. In *E. coli*, the genes are transcribed independently. A strong promoter drives the transcription of *gyrBA* operon in both *M. smegmatis* and *M. tuberculosis*. The vertical arrows and their thickness depict the promoters and their strength. tss; transcription start site.

gyrA promoter showed that the -10 region is responsible for both promoter strength and supercoil-sensitive behaviour²⁷. Paradoxically, the promoter region harbours a sequence that matches the *E. coli* consensus for extended -10 promoters²⁸. Since most extended -10 promoters do not show RST, it appears unlikely that the sequence of the -10 region alone is responsible for RST. Our recent analysis suggests that DNA curvature around the transcription start point plays a role in RST in *E. coli*²⁹.

The *gyr* operon in *M. smegmatis* is induced by novobiocin at the transcriptional level. However, unlike *E. coli*, minimal promoter of *gyr* operon do not confer RST in this case³⁰. The presence of a strong CHPS (cruciform/hairpin potential sequence) with an 8 base pair stem and a 4 base loop in the 5' untranslated region suggested a potential mechanism for RST in *M. smegmatis* and found to have a positive effect on promoter activity but is not sufficient for the operon to respond to novobiocin. The induction of the genomic copy and the results with the constructs harbouring upstream and downstream DNA sequences show the essential role played by promoter distal elements. DNA elements that are present 600 bp downstream of the promoter are necessary for RST to occur in the plasmid context³⁰. This suggests the involvement of long range interactions and formation of repressor loops which could either prevent the binding of the polymerase to the promoter or prevent its release. Since such repressor loops are stabilized by negative supercoiling^{31–33}, the repression would occur in a supercoil sensitive manner. However, the downstream element along with the minimal promoter region is not sufficient to respond to novobiocin. Sequential deletions of both upstream and downstream distal regions reveal interesting novel features of the regulation. Therefore, RST appears

to operate in *M. smegmatis* by some sort of de-repression rather than direct induction and a distal element has a negative influence on transcription³⁰.

A comparison of the expression of DNA gyrase in *M. smegmatis* and *M. tuberculosis* reveals an amalgamation of conserved and divergent features (Figure 1). In addition to conserved genomic arrangement and dicistronic organization, the primary promoter in *M. tuberculosis*, P_{B1} , is located upstream of the *gyrB* gene at a position similar to that of the *M. smegmatis gyr* promoter³⁴. Furthermore, the promoter region of P_{B1} shows extensive conservation with P_{gyr} , the promoter driving the *gyr* genes of *M. smegmatis*, indicating the evolutionary relationship. Apart from the primary promoter, the *gyr* locus in *M. tuberculosis* employs at least three other promoters³⁴. These additional promoters are weak and appear to play a regulatory role. P_A , the internal promoter for *gyrA* is 70-fold weaker than P_{B1} in exponentially growing *M. tuberculosis*, possibly employs an *M. tuberculosis*-specific sigma factor. Moreover, P_A may be induced under specific conditions which require the production of excess GyrA. Induction of GyrA alone in *E. coli* in response to treatment with GyrA inhibitors has been demonstrated earlier³⁵. The other weak promoter, P_R , is divergently oriented and almost completely overlaps P_{B1} . Therefore, the binding of RNA polymerase to one of them would prevent binding in the opposite orientation. There are no identifiable coding sequences upstream of *gyrB* that P_R could be involved in transcribing, suggesting the function of P_R to be regulatory. Overlapping, mutually exclusive promoters are one of the mechanisms for regulating gene expression³⁶. Recruitment of the polymerase to P_R would decrease expression of DNA gyrase by reducing transcription initiation at P_{B1} . In the converse scenario, as in relaxation of the template, P_R is repressed and P_{B1} gets induced to almost the same extent.

These studies highlight the importance of regulation of constitutively expressed, housekeeping, essential functions. While RST is a convenient mechanism to attain steady state levels of the enzyme, the complete molecular details of its operation vary and are not yet understood. Analysis of the promoter region of *M. smegmatis* and *M. tuberculosis* reveals a distinct lack of any axial distortion upstream of the $+1$ start site unlike *E. coli*²⁹. To further substantiate the demarcation, all known gyrase promoters were analysed for the presence of curvature in the vicinity of the -10 region ($+5$ nucleotides). It is noteworthy that roughly half of these show a significant curvature in this region while others do not²⁹. Interestingly, both the position and the extent of curvature are conserved between the *E. coli* and *Klebsiella pneumoniae gyrA* promoters²⁹. How a promoter distal (downstream) element located within the ORF contributes for RST in *M. smegmatis* is not clear. Although it appears that trans factor/s could be involved in promoting the long range interactions, the molecular mechanism is still elusive. *M. tuberculosis*




Figure 2. Twin domain of supercoiling. During transcription, RNA polymerase movement results in positive (+) in front and negative (–) supercoils behind the tracking machine. Action of the topoisomerases relieve the excessive supercoiling ensuring the further movement of RNA polymerase.

genome unlike that of *M. smegmatis*, has its own variant mechanism of RST. The RST response in case of *M. tuberculosis* is extremely slow. The gyrases from both the species are very similar (about 90% identity) and hence the difference in the response is not likely due to catalytic properties. On the other hand, rate of transcription is varied between the species. *M. tuberculosis* RNA polymerase is at least 3–5 times slower than *M. smegmatis* enzyme³⁷. Lower transcriptional rates in conjunction with yet unknown features may contribute for the difference in the response.

In contrast to *gyrB* and *gyrA* genes which are transcribed by single promoters, topoisomerase I expression appears to be regulated by multiple promoters in *E. coli*^{35,38}. Two of them seem to be dependent on sigma-70, the major sigma factor of *E. coli* RNA polymerase. Amongst the others, one promoter is active during heat shock response while the other directs transcription at stationary phase of the growth^{38–40}. As a consequence, the overall topoisomerase I expression is the net result of combined activation of various promoters³⁸. Although the underlying mechanisms of regulation of *topA* and *gyr* genes by supercoiling are not completely understood, these represent an efficient homeostatic mechanism for the maintenance of supercoiling within physiological limits. For example, when global supercoiling goes down, *topA* is repressed while *gyr* genes are induced, compensating for the deficit in supercoiling. Instead of transfactor/s, DNA topology, the substrate/product for topoisomerases, directly modulates the levels of the enzymes in a manner reminiscent of product mediated inhibition in metabolic pathways.

Post-transcriptional regulation: mRNA stability

The half-life of the bulk of the mRNA in *E. coli* is 2.4 min at 37°C⁴¹. This short half-life could reflect the fastgrowing nature of *E. coli*, possibly facilitating rapid adaptation to environmental changes⁴². Thus, one would expect mycobacteria and other slow-growing organisms would have more stable messages. In addition, the regulation of degradation of these messages would be different. Based

on this hypothesis, stability of the DNA gyrase mRNA in *M. smegmatis* was analysed⁴³. A secondary structure near the 5' end of mRNA that protects the message against degradation was identified (Figure 3). The stabilization effect is significantly pronounced in nutrient-deprived conditions. In addition to the transcriptional regulation discussed earlier, the nutrient-dependent stabilization of the gyrase message, represents a second, hitherto unexplored, level of regulation of the *gyr* genes in any organism. While, in general, stabilization of a housekeeping message would be important for slow-growing organisms like mycobacteria, in the specific case of DNA gyrase, it probably has additional significance due to the operonic arrangement and for all known biological functions both proteins are required in equimolar amounts. Since genes present downstream in an operon are usually under-represented at the protein level, it would be useful for the organism to evolve methods to prevent this discrepancy. The mycobacterial *gyr* operon attempts to circumvent this problem by subtle changes in its primary sequence (see later section). In such a context, the presence of a stabilizing secondary structure is probably an additional mechanism to ensure that the downstream message is maintained long enough to be translated efficiently. Thus, *M. smegmatis* appears to rely on two distinct sensors: a promoter-proximal sensor for nutrient levels and a promoter-distal sensor for DNA topology. Although not experimentally verified for its function, similar secondary structure is found upstream of *gyrBA* operon in *M. tuberculosis*.

Enhanced stability of mRNA upon starvation has been reported in many organisms^{42,44,45} which would allow the cells to utilize the already synthesised messages to their fullest and conserve resources when they are scarce. This is arguably more important for organisms like mycobacteria that grow slowly even under nutrient-rich conditions. In these organisms, the lower rate of transcription elongation is probably compensated for by enhanced stability of the message. Secondary structures at or near the 5' end of the mRNA in *E. coli* are believed to function by preventing access of RNase E⁴⁶. Furthermore, cleavage by RNase E appears to be the primary rate




Figure 3. Translational regulation of *gyr* mRNA. The stabilizing stem loop structure is shown at 5' end of the *gyrBA* mRNA. The SD and start codon sequences of the individual cistrons are shown which suggest translational optimization. The dots between SD sequence and start codon represent the actual number of nucleotides.

determining step in the degradation of most messages in *E. coli*⁴⁷. Genomes of *M. tuberculosis*¹⁰ and *Mycobacterium leprae*⁴⁸ encode for a homologue of this enzyme. Genome-wide analysis of the distribution of secondary structures indicates that genes in slow-growing organisms like *M. tuberculosis* are more likely to have a strong secondary structure ahead of them than those in fast-growing organisms like *E. coli*⁴⁹. This probably protects a majority of the messages against the degradative activities. Such a strategy would make economic sense for a slow growing organism that does not necessarily need to respond to environmental changes rapidly^{29,34,50}.

Post-translational regulation

The *gyrBA* dicistron in both *M. smegmatis* and *M. tuberculosis* exhibits additional interesting regulatory features (Figure 3). The *gyrB* has a near perfect Shine-Dalgarno (SD) sequence upstream of a weak start codon while *gyrA* has relatively weak SD sequence and an efficient start codon. Such an arrangement suggests a translational regulation that could facilitate the production of equimolar amounts of the two subunits that constitute the heterotetrameric functional holoenzyme.

Notably, there is a remarkable difference in the activities of DNA gyrase from *E. coli* and mycobacteria. *M. smegmatis* DNA gyrase has 3–5 fold weaker ATP hydrolyzing activity compared to that of *E. coli*⁵¹. However, the enzyme is a potent decatenase suggesting a more important role during segregation of daughter chromosomes. Since *M. tuberculosis* genome encodes only for topoisomerase I and DNA gyrase¹⁰, the DNA gyrase is likely to possess strong decatenase activity to take care of added responsibility of daughter genome segregation.

Additional post translational measures seem to play a role in the expression of functional gyrase in mycobacteria. The mycobacterial recombinant DNA gyrase expressed in *E. coli* do not show high specific activity when analysed for supercoiling or ATPase activities in contrast to the enzymes isolated from wild type cells or over-expressed in mycobacteria (unpublished). This suggests a role for post-translational modification of the enzyme. Furthermore, the presence of two *gyrB* in *M. smegmatis* raises interesting possibilities regarding the intracellular functions¹¹. The GyrB encoded by *gyrBA* operon is known to be associated with GyrA in a tetrameric holoenzyme that carries out the DNA supercoiling reaction⁵¹. From our comparative analysis it appears that the additional *gyrB*, termed as orphan *gyrB* is a functional allele and hence retained in the genome at a different location. Considering the difference in the growth rates of *M. smegmatis* and *M. tuberculosis*, the orphan GyrB could be contributing to the higher levels of enzymatic activity required during exponential growth phase. Alternatively, it is expressed differentially under certain conditions as an immediate requirement for cellular function.

Another point to be noted is that GyrB is intrinsically less stable than GyrA in *E. coli*⁵² and also in *M. smegmatis* (unpublished results).

Conclusions

Diverse topoisomerases influence the topological state of the genome. Although topoisomerases are essential house-keeping functions, the fine tuning of their expression is important in order to maintain the balanced topological state. Our analysis of regulation of gyrase expression between two species of mycobacteria has revealed an amalgamation of several concepts with important species specific differences. The conserved features include dicistronic organization, mycobacteria specific promoters, RNA stability etc. However, autoregulation of transcription appears to have species specific variation. In *M. smegmatis* promoter distal downstream elements and possibly transactors have a role in RST while in *M. tuberculosis*, overlapping mutually exclusive divergently organized promoters regulate the process. The organization as an operon in order to assemble heterotetrameric enzyme rapidly and extraordinary stability of the dicistronic mRNA are some of the measures taken by these group of bacteria to compensate for slower growth rates. The studies on regulation of topoisomerase I expression now underway would reveal other facets of regulation which contribute to attain cellular homeostasis in DNA topology.

1. Gellert, M., DNA topoisomerases. *Annu. Rev. Biochem.*, 1981, **50**, 879–910.
2. Droge, P. Protein tracking-induced supercoiling of DNA: a tool to regulate DNA transcription *in vivo*? *Bioassays*, 1994, **16**, 91–99.
3. Bliska, J. B. and Cozzarelli, N. R., Use of site-specific recombination as a probe of DNA structure and metabolism *in vivo*. *J. Mol. Biol.*, 1987, **194**, 205–218.
4. Champoux, J. J., DNA topoisomerases: structure, function and mechanism. *Annu. Rev. Biochem.*, 2001, **70**, 369–413.
5. Reece, R. J. and Maxwell, A., DNA gyrase: structure and function. *Crit. Rev. Biochem. Mol. Biol.*, 1991, **26**, 335–375.
6. Maxwell, A., DNA gyrase as a drug target. *Trends Microbiol.*, 1997, **5**, 102–109.
7. Madhusudan, K., Ramesh, V. and Nagaraja, V., Molecular cloning of *gyrA* and *gyrB* genes of *Mycobacterium tuberculosis*: analysis of nucleotide sequence. *Biochem. Mol. Biol. Int.*, 1994, **33**, 651–660.
8. Madhusudan, K. and Nagaraja, V., *Mycobacterium smegmatis* DNA gyrase: cloning and overexpression in *Escherichia coli*. *Microbiology*, 1995, **141**, 3029–3037.
9. Brosch, R., Gordon, S. V., Buchrieser, C., Pym, A. S., Garnier, T. and Cole, S. T., Comparative genomics uncovers large tandem chromosomal duplications in *Mycobacterium bovis* BC Pasteur. *Yeast*, 2000, **17**, 111–123.
10. Cole, S. T. *et al.*, Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature*, 1998, **393**, 537–544.
11. Jain, P. and Nagaraja, V., An orphan *gyrB* in the *Mycobacterium smegmatis* genome uncovered by comparative genomics. *J. Genet.*, 2002, **81**, 105–110.
12. Nagaraja, V., Sikder, D. and Jain, P., DNA topoisomerase I from mycobacteria – a potential drug target. *Curr. Pharm. Des.*, 2002, **8**, 1995–2007.

13. Madhusudan, K. and Nagaraja, V., Alignment and phylogenetic analysis of type II DNA topoisomerases., *J. Biosci.*, 1996, **21**, 613–629.
14. Huang, W. M., Type II DNA topoisomerase genes. *Adv. Pharmacol.*, 1994, **29**, 201–225.
15. Chatterji, M., Unniraman, S., Maxwell, A. and Nagaraja, V., The additional 165 amino acids in the B protein of *Escherichia coli* DNA gyrase have an important role in DNA binding. *J. Biol. Chem.*, 2000, **275**, 22888–22894.
16. Liu, L. F. and Wang, J. C., Supercoiling of the DNA template during transcription. *Proc. Natl. Acad. Sci. USA*, 1987, **84**, 7024–7027.
17. Kreuzer, K. N. and Cozzarelli, N. R., *Escherichia coli* mutants thermosensitive for deoxyribonucleic acid gyrase subunit A: effects on deoxyribonucleic acid replication, transcription, and bacteriophage growth. *J. Bacteriol.*, 1979, **140**, 424–435.
18. Sanzey, B., Modulation of gene expression by drugs affecting deoxyribonucleic acid gyrase. *J. Bacteriol.*, 1979, **138**, 40–47.
19. Yang, H. L., Heller, K., Gellert, M. and Zubay, G., Differential sensitivity of gene expression by drugs affecting deoxyribonucleic acid gyrase. *Proc. Natl. Acad. USA*, 1979, **76**, 3304–3308.
20. Phoenix, P., Raymond, M. A., Masse, E. and Drolet, M., Role of DNA topoisomerases in the regulation of R-loop formation *in vitro*. *J. Biol. Chem.*, 1997, **272**, 1473–1479.
21. Dorman, C. J., Fleming Lecture: DNA topology and the global control of bacterial gene expression: implications for the regulation of virulence gene expression. *Microbiology*, 1995, **141**, 1271–1280.
22. Wang, J. C. and Lynch, A. S., Transcription and DNA supercoiling. *Curr. Opin. Genet. Dev.*, 1993, **3**, 764–768.
23. Zechiedrich, E. L., Khodursky, A. B., Bachellier, S., Schneider, R., Chen, D., Lilley, D. M. and Cozzarelli, N. R., Roles of topoisomerases in maintaining steady-state DNA supercoiling in *Escherichia coli*. *J. Biol. Chem.*, 2000, **275**, 8103–8113.
24. Menzel, R. and Gellert, M., Regulation of the genes for *E. coli* DNA gyrase: homeostatic control of DNA supercoiling. *Cell*, 1983, **34**, 105–113.
25. Menzel, R. and Gellert, M., Fusions of the *Escherichia coli* *gyrA* and *gyrB* control regions to the galactokinase gene are inducible by coumermycin treatment. *J. Bacteriol.*, 1987, **169**, 1272–1278.
26. Menzel, R. and Gellert, M., Modulation of transcription by DNA supercoiling: a deletion analysis of the *Escherichia coli* *gyrA* and *gyrB* promoters. *Proc. Natl. Acad. Sci. USA*, 1987, **84**, 4185–4189.
27. Straney, R., Krah, R. and Menzel, R., Mutations in the –10 TATAAT sequence of the *gyrA* promoter affect both promoter strength and sensitivity to DNA supercoiling. *J. Bacteriol.*, 1994, **176**, 5999–6006.
28. Keilty, S. and Rosenberg, M., Constitutive function of a positively regulated promoter reveals new sequences essential for activity. *J. Biol. Chem.*, 1987, **262**, 6389–6395.
29. Unniraman, S. and Nagaraja, V., Axial distortion as a sensor of supercoil changes: a molecular model for the homeostatic regulation of DNA gyrase. *J. Genet.*, 2001, **80**, 119–124.
30. Unniraman, S. and Nagaraja, V., Regulation of DNA gyrase operon in *Mycobacterium smegmatis*: a distinct mechanism of relaxation stimulated transcription. *Genes Cells*, 1999, **4**, 697–706.
31. Borowiec, J. A., Zhang, L., Sasse-Dwight, S. and Gralla, J. D., DNA supercoiling promotes formation of a bent repression loop in lac DNA. *J. Mol. Biol.*, 1987, **196**, 101–111.
32. Choy, H. E., Park, S. W., Parrack, P. and Adhya, S., Transcription regulation by inflexibility of promoter DNA in a looped complex. *Proc. Natl. Acad. Sci. USA*, 1995, **92**, 7327–7331.
33. Lewis, D. E., Geanacopoulos, M. and Adhya, S., Role of HU and DNA supercoiling in transcription repression, specialized nucleoprotein repression complex at gal promoters in *Escherichia coli*. *Mol. Microbiol.*, 1999, **31**, 451–461.
34. Unniraman, S., Chatterji, M. and Nagaraja, V., DNA gyrase genes in *Mycobacterium tuberculosis*: a single operon driven by multiple promoters. *J. Bacteriol.*, 2002, **184**, 5499–5456.
35. Neumann, S. and Quinones, A., Discoordinate gene expression of *gyrA* and *gyrB* in response to DNA gyrase inhibition in *Escherichia coli*. *J. Basic Microbiol.*, 1997, **37**, 53–59.
36. Nagaraja, V., Control of transcription initiation. *J. Biosci.*, 1993, **18**, 13–25.
37. Harshey, R. M. and Ramakrishnan, T., Rate of RNA chain growth in *Mycobacterium tuberculosis* H₃₇R_v. *J. Bacteriol.*, 1977, **129**, 616–622.
38. Rui, S. and Tse-Dinh, Y.-C., Topoisomerase function during bacterial responses to environmental challenge. *Front. Biosci.*, 2003, **8**, 256–263.
39. Leslie, S. A., Jovanovich, S. B., Tse-Dinh, Y.-C. and Burgess, R. R., Identification of a heat shock promoter in the *top A* gene of *Escherichia coli*. *J. Bacteriol.*, 1990, **172**, 6871–6874.
40. Qi, H., Menzel, R. and Tse-Dinh, Y.-C., Regulation of *Escherichia coli* *top A* gene transcription: involvement of a sigma-S dependent promoter. *J. Mol. Biol.*, 1997, **267**, 481–489.
41. Regnier, P. and Arraiano, C. M., Degradation of mRNA in bacteria: emergence of ubiquitous features. *BioEssays*, 2000, **22**, 235–244.
42. Takayama, K. and Kjelleberg, S., The role of RNA stability during bacterial stress responses and starvation. *Environ. Microbiol.*, 2000, **2**, 355–365.
43. Unniraman, S., Chatterji, M. and Nagaraja, V., A hairpin near the 5' end stabilizes the DNA gyrase mRNA in *Mycobacterium smegmatis*. *Nucl. Acids Res.*, 2002, **30**, 5376–5381.
44. Condon, C., Putzer, H. and Grunberg-Manago, M., Processing of the leader mRNA plays a major role in the induction of *thrS* expression following threonine starvation in *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA*, 1996, **93**, 6992–6997.
45. Thorne, S. H. and Willimans, H. D., Adaptation to nutrient starvation in *Rhizobium leguminosarum* by *phaseoli*: analysis of survival, stress resistance and changes in macromolecular synthesis during entry to and exit from stationary phase. *J. Bacteriol.*, 1997, **179**, 6894–6901.
46. Hansen, M. J., Chen, L. H., Fejzo, M. L. and Belasco, J. G., The *ompA* 5' untranslated region impedes a major pathway for mRNA degradation in *Escherichia coli*. *Mol. Microbiol.*, 1994, **12**, 707–716.
47. Jain, C., Deana, A. and Belasco, J. G., Consequences of RNase E scarcity in *Escherichia coli*. *Mol. Microbiol.*, 2002, **43**, 1053–1064.
48. Cole, S. T. *et al.*, Massive gene decay in leprosy bacillus. *Nature*, 2001, **409**, 1007–1011.
49. Unniraman, S., Prakash, R. and Nagaraja, V., Alternate paradigm for intrinsic transcription termination in eubacteria. *J. Biol. Chem.*, 2001, **276**, 41850–41855.
50. Papavinasandaram, K. G. *et al.*, Slow induction of RecA by DNA damage in *Mycobacterium tuberculosis*. *Microbiology*, 2001, **147**, 3271–3279.
51. Manjunatha, U. H., Dalal, M., Chatterji, M., Radha, D. R., Visweswariah, S. S. and Nagaraja, V., Functional characterization of mycobacterial DNA gyrase: an efficient decatenase. *Nucl. Acids Res.*, 2002, **30**, 2144–2153.
52. Higgins, N. P., Peebles, C. L., Sugino, A. and Cozzarelli, N. R., Purification of subunits of *Escherichia coli* DNA gyrase and reconstitution of enzymatic activity. *Proc. Natl. Acad. Sci. USA*, 1978, **75**, 1773–1777.

ACKNOWLEDGEMENTS. I would like to thank my colleagues especially S. Unniraman, U. H. Manjunatha, D. R. Radha, M. Chatterjee and P. Jain who carried out most of the experiments. H. V. Jayashree and J. Robert have helped in the preparation of the manuscript. Research in my laboratory for this work has been supported by extramural grants from Council for Scientific and Industrial Research, Department of Science and Technology, Indian Council of Medical Research, Government of India. Part of the infrastructural support has come from ICMR Centre for Advanced Study in Molecular Medical Microbiology to the department of Microbiology and Cell Biology, Indian Institute of Science.