

Table 1. (Contd...)

Seaweed	Pre-monsoon	Monsoon	Post-monsoon
<i>H. valentiae</i> (Turner) Montagne	++	–	+++
<i>Sarconema filiforme</i> (Sonder) Kylin	+	+	++
<i>S. scinaoides</i> Børgesen	–	–	+
<i>Champia compressa</i> Harvey	–	–	++
<i>C. globulifera</i> Børgesen	–	–	+
<i>Centroceras clavulatum</i> (C. Agardh) Montagne	+	–	++
<i>Ceramium rubrum</i> Auctorum	+	+	++
<i>Wrangelia argus</i> (Montagne) Montagne	++	–	–
<i>Caloglossa leprieurii</i> (Montagne) G. Martens	++	++	++
<i>Bostrychia tenella</i> (Lamouroux) J. Agardh	++	++	++
<i>Chondria dasyphylla</i> (Woodward) J. Agardh	+	++	++
<i>Laurencia papillosa</i> (C. Agardh) Greville	+	–	–
<i>Polysiphonia platycarpa</i> Børgesen	+	–	+

+++ Dominant; ++ Common; + Sparse; – Absent.

exhibiting live seaweeds, which could boost the ecotourism industry, and ultimately lead to the conservation of these species.

- http://bhagirath.faihtweb.com/daman_diu.html
- Sreenivasa Rao, P. and Trivedi, M. K., In *Taxonomy of Algae* (eds Desikachary, T. V. and Raja Rao, V. N.), University of Madras Publication, 1980, pp. 255–260.
- Srinivasan, K. S., In *Phycologia Indica: Icones of Indian Marine Algae, Vol. – I*, BSI Publication, Calcutta, 1969, p. 52.
- Srinivasan, K. S., In *Phycologia Indica: Icones of Indian Marine Algae, Vol. – II*, BSI Publication, Calcutta, 1973, p. 60.
- Mantri, V. A., Joshi, H. V. and Zaidi, S. H., *Seaweed Res. Util.*, 2004, **26**, 231–236.

- May, R. M., *Science*, 1988, **241**, 1441–1449.
- Taylor, W. R., In *Algae of the Eastern Tropical and Subtropical Coasts of the Americas*, University of Michigan Publication, 1960, p. 662.
- Desikachary, T. V., Krishnamurthy, V., and Balkrishnan, M. S., In *Rhodophyta – II. Taxonomy Part – II A*, Madras Science Foundation Publication, 1990, p. 279.
- Mishra, J. N., In *Phaeophyceae in India*, Indian Council of Agricultural Research, New Delhi, 1967, p. 203.
- Krishnamurthy, V., *Algae of India and Neighboring Countries – Chlorophycota*, Oxford and IBH, 2000, p. 210.
- Singh, J. S., *Curr. Sci.* 2002, **82**, 638–647.
- Sahoo, D., Nivedita and Debasish, In *Seaweeds of Indian Coast*, A.P.H. Publishing Corporation, New Delhi, 2001, p. 283.

ACKNOWLEDGEMENTS. We thank the Director, CSMCRI, Bhavnagar for encouragement. Thanks are also due to the Head, Marine Algae and Marine Environment Discipline, CSMCRI for providing facilities.

Received 23 December 2004; revised accepted 16 October 2005

VAIBHAV A. MANTRI
P. V. SUBBA RAO*

*Marine Algae and Marine Environment
Discipline,
Central Salt and Marine Chemicals
Research Institute,
Bhavnagar 364 002, India
*For correspondence.
e-mail: pvsubarao@csmcricri.org*

A new *Orontobryum* Mitt. ex Fleisch. from Darjeeling, eastern Himalayas

Genus *Orontobryum* Mitt. ex Fleisch. was established by Fleischer¹ from Sikkim, eastern Himalayas with its type species *Orontobryum hookeri* (Mitt.) Fleisch. At the time of institution of this genus, Fleischer was little reluctant whether it should be separated as new genus because of its close affinity with genus *Macrotamnium* Fleisch. belonging to family Hylocomiaceae (order Hypnobryales), which was also the view of Brotherus². However, Brotherus³ revised his opinion and placed the *Orontobryum* in the family Hookeriaceae (order Hookeriales). The status of the genus given by Brotherus³

has been followed by Gangulee⁴ and the same is maintained during the present study. Genus *Orontobryum* was earlier represented in India by two species, i.e. *O. hookeri* (Mitt.) Fleisch. and *O. recurvulum* Gangulee known from Sikkim, and Sikkim and Bhutan respectively, and is endemic to eastern Himalayas. During the recent study on the bryophytes of Rim-bick, Darjeeling some specimens approaching to genus *Orontobryum* have been encountered. A critical and comparative study has revealed that these specimens are clearly distinctive from the hitherto known species of the genus. Hence it is

described here as *Orontobryum darjeelingensis* sp. nov., which is new to the science.

Orontobryum darjeelingensis Nath, Asthana & Sahu sp. nov. (Figure 1)

Folia late cordata ± 1 mm longa et ± 1 mm lata, margineum serrata–dentata ad basa. Ramorum folia ovata–oblonga 0.064–0.7 mm longa et ± 0.35 mm lata, margineum dentata ad mediana. Foliorum cellulae apica 32–40 \times 8–12 μ m, mediana 40–44 \times 4 μ m, cellulae alares 30–32 \times 25–37.5 \times 12.5–20 μ m, pale-light brown. Ramorum foliorum cellulae apica 34–40 \times 4–7 μ m, mediana 40–68 \times 4 μ m,




Figure 1. *Orontobryum darjeelingensis* sp. nov. 1, Plant (diagrammatic); 2, Plant (a portion); 3–5, Stem leaves; 6, 7, Branch leaves; 8, Apical cells (stem leaf); 9, Marginal cells (stem leaf); 10, Median cells (stem leaf); 11, Alar cells (branch leaf); 12, 13, Alar cells (stem leaf).

cellulae alares $27.5\text{--}35 \times 12\text{--}16 \mu\text{m}$, coloratae, cellulae ad juncturum caule parviores.

Plants medium to robust, yellowish-brown, pinnately branched, branches slender. Stem brown to reddish-brown. Leaves in several rows, sparsely arranged, recurved at base, acute at apex. Stem leaves larger, widely cordate with acute apex, ± 1 mm long and ± 1 mm wide, margin serrate-dentate nearly up to base; costa double, short and indistinct; leaf cells at apex $32\text{--}40 \times 8\text{--}12 \mu\text{m}$, at middle $40\text{--}44 \times 4 \mu\text{m}$, alar cells few and well differentiated, yellow to yellowish-brown, $32.5\text{--}37.5 \times 12.5\text{--}20 \mu\text{m}$. Branch leaves smaller, $0.64\text{--}0.7$ mm long and 0.35 mm

wide, ovate-oblong (cordate) with acute apex, margin dentate, dentitions $10\text{--}12.5$ (~ 15) μm high up to middle, costa double, short, not distinct. Branch leaf cells at apex $34\text{--}40 \times 4\text{--}7 \mu\text{m}$, at middle $40\text{--}68 \times 4 \mu\text{m}$, alar cells yellowish brown, $27.5\text{--}35 \times 12\text{--}16 \mu\text{m}$, leaf cells somewhat pointed at ends. Fertile plants not seen.

Plants grow on stem bark under moist and shady conditions at Forest Rest House, Rimbick, district Darjeeling at an altitude of ca. 2286 m and temperature range 1.5 (min) to 15°C (max).

Specimens examined: *Orontobryum darjeelingensis* sp. nov. India: eastern Himalayas, Rimbick (alt. ca. 2286 m),

16.04.1965, Leg. S. Chandra, HOLOTYPE – 202016 A (LWG).

Orontobryum hookeri (Mitt.) Fleish. ex Broth. India: Sikkim Himalaya. Ratong river, Leg. J.D. Hooker 819 (W. 336) (Isosyntype).

Orontobryum recurvulum Gangulee. Bhutan: Khagumpa (alt. ca 6000 ft), Herb. of the Late East India Company No. 750, Hb. Griffith 130 (Isotype).

A critical and comparative study with the type specimens of allied Indian species of genus *Orontobryum* has revealed that plant specimens of *O. darjeelingensis* sp. nov. closely approach *O. recurvulum* Gangulee in plant habit, widely

cordate stem leaves, branch leaves with serrate–dentate margin and well differentiated, yellow–light-brown alar cells; but the former can be distinctly recognized from the latter in having sparsely arranged leaves on stem, widely cordate leaf shape (subtriangular to triangular in *O. recurvulum*), serrate–dentate margin nearly up to the base in leaves, larger apical cells, viz. $32\text{--}40 \times 8\text{--}12 \mu\text{m}$ ($25\text{--}27.5 \times 5\text{--}7 \mu\text{m}$ in *O. recurvulum*) in stem leaves and $34\text{--}40 \times 4\text{--}7 \mu\text{m}$ in branch leaves ($20\text{--}22.5 \times 7.5\text{--}10 \mu\text{m}$ in *O. recurvulum*), smaller dentitions ($10\text{--}12.5 \mu\text{m}$ high) at branch leaf margins compared to *O. recurvulum* ($15\text{--}17.5 \mu\text{m}$ high dentitions) and comparatively light coloured, larger alar cells in branch leaves ($27.5\text{--}35 \times 12.5\text{--}15 \mu\text{m}$ in *O. darjeelingensis* and $20\text{--}25 \times 7.5\text{--}10 \mu\text{m}$ in *O. recurvulum*). *O. hookeri* can be well differentiated from *O. darjeelingensis* sp. nov. by closely arranged leaves on stem, ovate to oblong stem leaves ($\pm 0.9 \text{ mm} \times 0.64 \text{ mm}$) with serrate–spi-

nose margin restricted to apex, larger branch leaves ($0.83 \times 0.57 \text{ mm}$), narrower apical leaf cells and smaller inflated alar cells ($\pm 27 \times 15 \mu\text{m}$). A comparative study with isosytype and isotype specimens (obtained on loan from British Museum) of *O. hookeri* and *O. recurvulum* with specimens collected from Rimbick showed that the latter is clearly distinct from earlier known species. Hence it has been designated as *Orontobryum darjeelingensis* sp. nov.

1. Fleischer, M., *Hedwigia*, 1922, **63**, 209–216.
2. Brotherus, V. F., *Pflanzen Familien*, 1908, 1st edn.
3. Brotherus, V. F., *Sber Akad. Wiss. Wien Math. Natur. wiss. Kl. Abt. I*, 1922, **131**, 208–222.
4. Gangulee, H. C., *Mosses of Eastern India and Adjacent Regions*, 1976, **2**, 831–1546.

ACKNOWLEDGEMENTS. We are grateful to Dr P. Pushpangadan, Director, National

Botanical Research Institute, Lucknow for encouragement and providing the facilities. Our sincere thanks are also due to the Keeper of Botany, British Museum (Natural History), London, for providing authentic specimens of genus *Orontobryum* on loan for the present study.

Received 4 July 2005; revised accepted 29 August 2005

VIRENDRA NATH*
A. K. ASTHANA
VINAY SAHU

*Bryology Laboratory,
National Botanical Research Institute,
Lucknow 226 001, India*

**For correspondence.
e-mail: drvirendranath2001@rediffmail.com*