

4. Banasco, P., Fuente, L. De La, Gaultieri, G., Noya, F. and Arias, A., Fluorescent *Pseudomonas* spp. as biocontrol agents against forage legume root pathogenic fungi. *Soil Biol. Biochem.*, 1998, **10**, 1317–1323.
5. Dileep, C., Kumar, B. S. D. and Dube, H. C., Promotion of plant growth and yield by two rhizoplane fluorescent *Pseudomonas*. *Indian J. Exp. Biol.*, 1998, **36**, 399–402.
6. Pierson, E. A. and Weller, D. M., Use of mixture of fluorescent *Pseudomonas* to suppress take-all and improve growth of wheat. *Phytopathology*, 1994, **84**, 940–947.
7. Yeole, R. D. and Dube, H. C., Increased plant growth and yield through seed bacterization. *Indian Phytopathol.*, 1997, **50**, 316–319.
8. Kraus, J. and Loper, J., Characterization of genomic region required for production of antibiotic pyoluteorin by the biological control agent *Pseudomonas fluorescens* Pf-5. *Appl. Environ. Microbiol.*, 1995, **61**, 849–854.
9. Rodriguez, H. and Fraga, R., Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol. Adv.*, 1999, **17**, 319–339.
10. Seong, K. Y. and Shin, P. G., Effect of siderophore on biological control of plant pathogens and promotion of plant growth by *Pseudomonas fluorescens* ps88. *Agric. Chem. Biotechnol.*, 1996, **39**, 20–24.
11. Hagedorn, D. J. (ed.), *Compendium of Pea Diseases*, APS Press, 1984, p. 57.
12. King, E. O., Ward, M. K. and Raney, D. E., Two simple media for the demonstration of pyocanin and fluorescin. *J. Clin. Lab. Med.*, 1954, **44**, 301–307.
13. Brown, M. E., Seed and root bacterization. *Annu. Rev. Phytopathol.*, 1974, **12**, 181–197.
14. O'Sullivan and O'Gara, F., Traits of fluorescent *Pseudomonas* spp. involved in suppression of plant pathogens. *Microbiol. Rev.*, 1992, **56**, 662–672.
15. Weller, D. M., Biological control of soil-borne plant pathogens in the rhizosphere with bacteria. *Annu. Rev. Phytopathol.*, 1998, **26**, 379–407.
16. Nautiyal, C. S., An efficient microbial growth medium for screening of phosphate solubilizing microorganisms. *FEMS Microbiol. Lett.*, 1999, **170**, 265–270.
17. Bakker, A. W. and Schippers, B., Microbial cyanide production in the rhizosphere in relation to potato yield reduction and *Pseudomonas* spp. mediated plant growth stimulation. *Soil Biol. Biochem.*, 1987, **19**, 451–457.
18. Leisinger, T. and Mangraff, F., Secondary metabolites of the fluorescent pseudomonas. *Microbiol. Rev.*, 1979, **43**, 422–442.
19. Mazzola, M. and Cook, R. J., Effects of fungal root pathogens on the population dynamics of biocontrol strains of fluorescent *Pseudomonas* in wheat rhizosphere. *Appl. Environ. Microbiol.*, 1991, **57**, 2171–2178.
20. De Boer, M., Bom, P., Kindt, F., Keurentjes, J. J. B., Sluis, I. V. D., Van Loon, L. C. and Bakker, P. A. H. M., Control of *Fusarium* wilt of radish by combining *Pseudomonas putida* strains that have different disease suppressive mechanisms. *Phytopathology*, 2003, **93**, 626–632.
21. Guetsky, R., Elad, Y. and Dinoor, A., Combining biocontrol agents to reduce the variability of biological control. *Phytopathology*, 2001, **91**, 621–627.
22. Duffy, B. K. and Weller, D. M., Use of *Gaeumannomyces graminis* var. *graminis* alone and in combination with fluorescent *Pseudomonas* spp. to suppress take-all of wheat. *Plant Dis.*, 1995, **79**, 907–911.

ACKNOWLEDGEMENTS. Necessary facilities provided by the Dean, College of Forestry and Hill Agriculture, are acknowledged. The work was carried out utilizing grants received under the Uttaranchal Diversified Agriculture Support Project.

Received 19 March 2005; revised accepted 27 August 2005

Histochemical and biochemical studies of parasite–host interaction of *Cassytha filiformis* Linn. and *Zizyphus jujuba* Lamk.

M. N. Abubacker^{1,*}, M. Prince¹ and Y. Hariharan²

¹PG and Research Department of Botany, National College, Tiruchirapalli 620 001, India

²Department of Biotechnology, Mahendra Arts and Science College, Kalipatti 637 501, India

Parasite–host interaction of *Cassytha filiformis* and *Zizyphus jujuba* through histochemical studies revealed the presence of specialized glandular cells facilitating adhesion of parasite to the host and further specialization to obtain nutrients from phloem tissue of the host. Histochemical studies indicated the presence of high acid phosphatase activity of the parasite, which revealed the digestion of macromolecules/energy transfer and intercellular transport of parasite. Partial photosynthesis activity of parasite was noticed by accepting hydrogen released from photolysis of water through Hill reaction.

Keywords: *Cassytha filiformis*, haustoria, Hill reaction, parasite–host interaction, *Zizyphus jujuba*.

PARASITES are unusual plants, well adapted to their mode of life. More than 2500 species of higher plants are known to live parasitically on other plants. These parasitic plants produce flowers and seeds and belong to several widely separated botanical families¹. They vary greatly in their dependence on their host plants. For example, *Viscum* (mistletoes) have chlorophyll but no roots and therefore depend on their hosts only for water and minerals. *Cuscuta* (dodders) and *Cassytha* (amarbeli) have little chlorophyll and no true roots. Hence they depend on their hosts for water, food and minerals. The most common and serious parasites belong to the following botanical families and genera.

Cuscutaceae: *Cuscuta* (dodders); Scrophulariaceae: *Striga* (witch weed); Orobanchaceae: *Orobanche* (broom rapes); Cassythaceae: *Cassytha* (amarbeli); Loranaceae: *Elytranthe*, *Korthalsella* and *Loranthus*, Viscaceae: *Arceuthobium* (dwarf mistletoes); *Phoradendron* (American true mistletoes) and *Viscum* (European true mistletoes).

Cassytha filiformis Linn. Cassythaceae (Figure 1a) is a twining parasitic, perennial angiosperm which adheres to the host by suckers (haustoria). The plant is characterized by leaves reduced to minute scales, small flower, hermaphrodite, sessile, spicate, perianth tube short and globose, stamens six, ovary globose and fruit a drupe enclosed in the enlarged inflated perianth tube. During winter *Cassytha* seeds grow along with the seeds of the host plant (*Zizyphus*) in infested fields. During the growing season, the

*For correspondence. (e-mail: abubackermn@yahoo.com)

seeds germinate and produce a slender yellowish shoot, but no roots. When in contact with a susceptible host, the shoot encircles the host plant, sends haustoria into it and begins to climb up the plant. The haustoria penetrate the stem or leaf and reach into the vascular tissues, from which they absorb food and water. Soon after contact with the host is established, the base of the *Cassytha* parasite shrivels and dries up, so that the parasite loses all connection with the ground and becomes completely dependent on the host for food and water. The parasite continues to grow and expand while the growth of host plants is suppressed, which may even die. In the meantime, the parasite plant

has developed flowers and produced seeds¹. Many host-parasite interaction studies of *Cuscuta*, *Viscum*, *Orobancha* and *Striga* have been cited in the literature¹⁻¹⁶. However, for *Cassytha* only little information is available. Hence histochemical and biochemical studies of parasite-host interaction of *C. filiformis* and *Zizyphus jujuba* were undertaken.

C. filiformis Linn. Cassythaceae (voucher specimen no. 3976)¹⁷, the parasitic plant with the host *Z. jujuba* Lamk. Rhamnaceae (voucher specimen no. 2557)¹⁷ were collected from Bharathidasan University, Tiruchirapalli campus and used for histochemical and biochemical studies. Microtome




Figure 1. Parasite-host interaction of *Cassytha filiformis* Linn. and *Zizyphus jujuba* Lamk. **a**, *C. filiformis* parasite (P), 20x; **b**, *Z. jujuba* host (H); **c**, Stem anatomy of *C. filiformis*, 200x; **d**, *C. filiformis* (P) form a collar around the host (H), 100x; **e**, Parasite penetrates the phloem (Ph) of host, 100x; **f**, Phosphatase and peroxidase activity along haustoria (Ha), 100x; **g**, Digestion of host cells, 400x.

Table 1. Chlorophyll estimation in *Zizyphus jujuba* (host)

<i>Zizyphus jujuba</i> (host)	(mg/g fresh weight)		
	Chlorophyll <i>a</i>	Chlorophyll <i>b</i>	Total Chlorophyll
Host without parasite interaction	8.92	12.03	20.95
Host with parasite interaction	7.41	9.28	16.69
Chlorophyll reduction (%)	16.93	22.99	20.33

sections of the parasite and host with parasite (suckers) haustoria complex were taken according to the standard procedure¹⁸.

For histochemical studies, sections of host-parasite complex were studied in phloroglucinol-HCl¹⁹ and toluidine blue 'O'²⁰ and photographed.

For histoenzymological studies, sections of the host-parasite complex were studied in acid phosphatase – metal salt method²¹, alkaline phosphatase – tetrazolium method²², succinate dehydrogenase reaction method²³ and peroxide reaction method²⁴.

For biochemical studies, quantitative estimation of chlorophyll *a*, *b* and total chlorophyll²⁵ and photosynthetic activity by Hill reaction studies²⁶ were made.

Stem anatomy of *C. filiformis* has shown a central cylinder of collateral vascular bundles (Figure 1 *b*). Inner cortex is wide and represented by enlarged palisade-like chlorenchymatous cells.

Phloroglucinol-HCl and toluidine blue 'O' histochemical studies revealed the presence of specialized glandular cells, facilitating adhesion of parasite to the host. The glandular *C. filiformis* cells continue to divide and spread like a collar around the host plant (Figure 1 *c*); they then penetrate the phloem (Figure 1 *d*). The results are comparable with earlier reports^{4,27}. Dorr and Kollmann⁴ reported the presence of similar cells in hemp parasitized by *Orobancha ramosa*. Lee and Lee²⁷ reported the same in *Cuscuta australis*.

Haustoria of the parasite showed elongated sieve elements. Penetration of *Z. jujuba* (host) tissue by *C. filiformis* (parasite) is not accompanied by crushing of host cells. Histochemical studies revealed high acid phosphatase and peroxidase activities along haustoria of parasite (Figure 1 *e*). Cortical or vascular tissues of the host indicate digestion of macromolecules (Figure 1 *f*) or energy transfer and intercellular transport for the parasite as indicated by studies^{5,9,16} in different species of *Cuscuta*.

Biochemical studies revealed reduction in chlorophyll *a*, *b* and total chlorophyll pigments of the host due to host-parasite interactions (Table 1), although the parasite (*Cassytha*) is itself partly autotrophic. Estimation of Hill reaction in parasite ensures partial photosynthetic activity by acceptance of hydrogen released from phytolysis of water, as reported by Dinelli *et al.*¹³ in the case of *Cuscuta campestris*. This was demonstrated by the initial reading of blue colour (1.301) changing to pale blue colour

(1.097) when DCPIP (Dichloro Phenol Indo Phenol) market dye for oxidized chlorophyll extract indicating photosynthesis.

The present work indicates that the parasite does not attack the tissues of the host beyond those that are necessary to establish a dynamic relationship with the host vascular tissue. Apparently, *C. filiformis* acquires the needed metabolites from the host vascular system rather than from a combination of host vascular translocates and nutrients gained by the destruction of host tissue.

1. Dorr, I., The haustorium of *Cuscuta* – new structural results. In *Parasitic Higher Plants* (eds Webber, H. C. and Forstreuter, W.), Proceedings of the 4th International Symposium, Philip University, Marburg, Germany, 1987, pp. 163–170.
2. Abu-Shakra, S., Miah, A. A. and Saghir, A. R., Germination of seed of branched broom rape (*Orobancha ramosa* L.). *Hortic. Res.*, 1970, **10**, 119–124.
3. Hawksworth, F. G. and Wiends, D., Biology and taxonomy of the dwarf mistletoes. *Annu. Rev. Phytopathol.*, 1970, **8**, 187–208.
4. Dorr, I. and Kollmann, R., Strukturelle Grundlagen des Parasitismus bei *Orobancha*. I. Wachstum der Haustorialzellen im Wirtsgewebe. *Protoplasma*, 1974, **80**, 245–259.
5. Fer, A., Photosynthesis and respiration in *Cuscuta lupuliformis*, Krock, seedlings during their preparasitic stage. *Physiol. Veg.*, 1976, **14**, 357–365.
6. Pennypacker, B. W., Nelson, P. E. and Wilhelm, S., Anatomic changes resulting from the parasitism of tomato by *Orobancha ramosa*. *Phytopathology*, 1979, **69**, 741–748.
7. Musselmann, L. J., The biology of *Striga*, *Orobancha* and other root parasitic weeds. *Annu. Rev. Phytopathol.*, 1980, **18**, 463–489.
8. Eplee, R. E., *Striga's* status as a plant parasite in the United States. *Plant Dis.*, 1981, **65**, 951–954.
9. Nagar, N. and Sanwal, G. C., Biochemical aspects of parasitism in *Cuscuta reflexa*: Inhibition of cell wall degrading enzymes of *Cuscuta* by non-susceptible plants. In Proceedings of 3rd International Symposium on Parasitic Weeds, Aleppo, CIMMYT, 1984.
10. Ihl, B. and Jacob, F., Functioning of *Cuscuta* haustoria by benzylaminopurine. *Haustorium*, 1990, **24**, 4.
11. Shaw, C. G. and Hennon, P. E., Spread, intensification and upward advance of dwarf mistletoe in thinned, young stands of Western hemlock in Southeast Alaska. *Plant Dis.*, 1991, **75**, 363–367.
12. Thoday, M. G., On the histological relations between *Cuscuta* and its host. *Ann. Bot.*, 1991, **25**, 655–682.
13. Dinelli, G., Bonetti, A. and Tibiletti, E., Photosynthetic and accessory pigments in *Cuscuta campestris* Yuncker and some host species. *Weed Res.*, 1993, **33**, 253–260.
14. Gomez, J. M., Importance of direct and indirect effects in the interaction between a parasitic angiosperm (*Cuscuta epithymum*) and its host plant (*Hormathophylla spinosa*). *Oikos*, 1994, **71**, 97–106.
15. Jeschke, W. D., Baumel, P., Rath, N., Czygan, F. C. and Proksch, P., I. Modelling the flow and partitioning of carbon and nitrogen

- in the holoparasite *Cuscuta reflexa* Roxb. and its host *Lupinus albus* L. II. Flows between host and parasite and within the parasitized host. *J. Exp. Bot.*, 1994, **45**, 801–812.
16. Quick, W. P., Localization of photosynthetic metabolism in the parasitic angiosperm *Cuscuta reflexa*. *Planta*, 1998, **205**, 506–513.
 17. Matthew, K. M., *The Flora of Tamil Nadu Carnatic*, The Rapinat Herbarium, St. Joseph's College, Tiruchirappalli, 1983.
 18. Johansen, D. A., *Plant Microtechnique*, McGraw Hill, New York, 1940.
 19. Gahan, P. B., *Plant Histochemistry and Cytochemistry: An Introduction*, Academic Press, Florida, 1984.
 20. O'Brien, T. P., Feder, N. and McCully, M. E., Polychromatic staining of plant cell walls by toluidine blue O. *Protoplasma*, 1964, **59**, 367–373.
 21. Gomeri, G., An improved histochemical technique for acid phosphatase. *Stain Technol.*, 1950, **25**, 81–85.
 22. Lojda, Z., Gossran, R. and Schiebeler, T. H., *Enzyme Histochemistry*, Springer Verlag, Heidelberg, 1979.
 23. Malik, C. P. and Singh, M. B., *Plant Enzymology and Histochemistry*, Kalyani Publishers, New Delhi, 1980.
 24. Dejong, D. W., Jansen, E. F. and Olson, A. C., Oxidoreductive and hydrolytic enzyme patterns in plant suspension culture. *Exp. Cell. Res.*, 1967, **47**, 139.
 25. Arnon, Copper enzymes in isolated chloroplasts polyphenol oxidase in *Beta vulgaris*. *Plant Physiol.*, 1949, **24**, 1–15.
 26. Hill, R., Oxygen produced by isolated chloroplasts. *Proc. R. Soc. London, Ser. B*, 1939, **127**, 192–210.
 27. Lee, K. B. and Lee, C. D., The structure and development of the haustorium in *Cuscuta australis*. *Can. J. Bot.*, 1989, **67**, 2975–2982.

ACKNOWLEDGEMENTS. We thank DST-FIST, New Delhi for providing infrastructure facilities for the Department of Botany, National College, Tiruchirappalli. We also thank Mr K. Santhanam, Patron Secretary, Mr K. Raghunathan, Secretary, Dr M. Nagarajan, Principal and Dr V. T. Sridharan, Head, Department of Botany, National College for encouragement.

Received 4 February 2005; revised accepted 15 September 2005

Long-term lowland rice and arable cropping effects on carbon and nitrogen status of some semi-arid tropical soils

K. L. Sahrawat^{1,*}, T. Bhattacharyya²,
S. P. Wani¹, P. Chandran², S. K. Ray², D. K. Pal²
and K. V. Padmaja¹

¹International Crops Research Institute for the Semi-Arid Tropics,
Patancheru 502 324, India

²National Bureau of Soil Survey and Land-use Planning,
Nagpur 440 010, India

Samples of surface (0–30 cm) soils were collected from eight sites in the semi-arid tropical regions of India to evaluate and compare the long-term effects of lowland

rice or paddy and non-rice or arable systems on soil organic C (SOC), soil inorganic C (SIC) and total N status. The results showed that soil samples from sites under lowland rice double cropping system had greater organic C and total N content than those from soils under rice in rotation with upland crop or under other arable systems. The SOC:N ratio was wider in soil samples from sites under lowland rice compared to those under other arable systems, which had lower C:N ratios. Samples from soils under lowland rice system tended to have a narrower SIC:N ratio than those under arable systems, indicating a better pedo-environment under paddy rice. Our results support earlier findings that sites under continuous wetland rice cropping accumulate organic matter and contain higher soil organic matter compared to the sites under other arable systems.

Keywords: Arable systems, lowland rice, natural vegetation, nitrogen, pedo-environment.

MAINTENANCE of fertility in soils of the tropics and semi-arid tropics (SAT), a prerequisite for sustainable increase in agricultural productivity, is a major challenge to farmers and researchers alike. Soils, especially those in the SAT are low in organic matter and nutrient reserves. High temperature in the tropics, which results in rapid decomposition and loss of organic matter, is the primary cause of the low organic matter status of soils. Moreover, under dryland agriculture, application of nutrients and organic matter through external inputs is generally low due to socio-economic conditions of the farmers and other factors. Under low or no external input management practices, agricultural productivity depends on the inherent fertility of soils, which is generally low¹.

Jenny and Raychaudhari² studied the effect of climate and cultivation on the nitrogen (N) and organic matter reserves in Indian soils. The approach used was based on collection of soil samples across the country from cultivated fields and forested soils in relation to a climatic grid in which mean annual temperature and mean annual precipitation appeared as independent variables. Based on the analysis of 500 soil samples for organic carbon (C) and total N across India, these authors showed that the climatic effects on soil organic matter and N status are pronounced. Soil N and C increased with increasing mean annual precipitation and decreased with increasing mean annual temperature. Soils in the drier regions had low reserves of organic matter and N compared to those in the humid and sub-humid zones of the country².

Velayutham *et al.*³ and Bhattacharyya *et al.*⁴ carried out studies on carbon stocks in Indian soils. Analysis of thousands of soil samples in the course of these studies helped prioritize research on C sequestration potential in soils of the SAT regions in India. Recently, the concept of quasi-equilibrium values of soil organic C relative to organic C status of forest soils as reference or control has been found

*For correspondence. (e-mail: k.sahrawat@cgiar.org)