

27. Abdul-Baki, A. A. and Anderson, J. D., Viability and leaching of sugars for germinating barley. *Crop Sci.*, 1970, **10**, 31–34.
28. Roberts, E. H. and Ellis, R. H., Water and seed survival. *Ann. Bot.*, 1989, **63**, 39–52.

ACKNOWLEDGEMENTS. D.V. is grateful to CSIR, New Delhi for awarding Senior Research Fellowship during the period of this work. We thank Dr D. K. Joshi, Nuclear Research Laboratory, IARI, New Delhi, for technical help during NMR analysis. The study was supported by the National Agricultural Technology Project (ICAR) under the Team of Excellence in Seed Technology.

Received 16 April 2008; revised accepted 19 February 2009

Species and site effects on leaf traits of woody vegetation in a dry tropical environment

S. K. Pandey, Hema Singh* and J. S. Singh

Department of Botany, Banaras Hindu University, Varanasi 221 005, India

Selected leaf traits (leaf area, leaf weight, specific leaf area and chlorophyll content) from eight woody species at four sites in the dry tropical Vindhyan forest were investigated in order to assess their variability across species and site conditions. The morphological traits such as leaf area and leaf weight were more variable than biochemical traits such as chlorophyll content. There was significant effect of species and site, and the site × species interaction was also significant for all traits. These traits were correlated except for specific leaf area, which was independent of leaf area. A combination of traits could discriminate between the sites. Between-site variability in leaf traits was smaller (1.3–1.5-fold) than between-species variability (1.7–12.5 fold). The larger inter-species variability reflects marked genotypic variability in the leaf traits, while the smaller between-site variability reflects phenotypic plasticity leading to adaptation to site conditions.

Keywords: Dry tropical forest, inter-species variability, leaf traits, woody species.

LEAF traits are often cited as the principal traits to relate plant resource use, biomass and ecosystem functioning^{1–4}. In addition, these traits are easy to quantify and convenient to compare among a large number of plant species. Leaf traits may be divided into two groups: functional traits and structural traits. Functional traits reflect the index of plant growth and metabolism. On the other hand, structural traits are indexes of biological characteristics of different plant species, and reflect the adaptation strategies

of plants to the environment. These traits, among others, include leaf area, leaf dry weight, specific leaf area (SLA) and chlorophyll content. SLA is the ratio of leaf area to leaf dry weight and, being strongly correlated with relative growth rate, maximum rate of photosynthesis³ and competitive ability, is often considered a key trait linked to plant functioning^{5–8}. It has been argued that SLA can provide important clues regarding future changes in community composition due to global change, if dryness is going to be altered in much of the tropics. The high-SLA leaves are productive^{6,9} but are necessarily also short-lived and vulnerable to herbivory^{10,11}. On the other hand, low-SLA leaves perform better in resource-poor environments⁷.

Leaf area plays an important role in light interception, water and nutrient use, growth and yield potential^{12–14}. Leaf size and SLA decline along gradients of decreasing moisture and/or nutrient availability^{15–20}. Lower SLA, due to thicker and/or denser leaves contributes to long leaf survival, nutrient retention, and protection from desiccation²¹, whereas small leaf size reduces boundary layer resistance, and helps maintain favourable leaf temperatures and higher photosynthetic water-use efficiency under the combination of high solar radiation and low water availability^{22,23}.

Chlorophyll is the most important pigment for photosynthesis^{24–26}. Chlorophyll concentration in leaves and canopies can be an indicator of photosynthetic capacity, developmental stage, plant productivity, environmental stress and nutrient management^{27–30}. Measurement of leaf chlorophyll content is also an indirect approach to estimate soil nitrogen^{31–33}.

Since the leaf traits are considered important for understanding vegetation response to a broad range of environmental factors, we examined four leaf traits, viz. leaf area, leaf weight, SLA and chlorophyll content in eight woody species on four sites of a dry tropical forest in the Vindhyan highland. We addressed the questions: (i) How much do the woody species occurring in a dry tropical environment differ in these leaf traits and how much are these leaf traits affected by site conditions? (ii) Can a combination of these leaf traits across species discriminate between the sites?

The study was conducted on four sites, viz. Ranitali, Neruiadamar, Bokrakhari and Hathinala of the Vindhyan dry tropical region (21°29′–25°11′N lat.; 78°15′–84°15′E long.), Sonbhadra District, Uttar Pradesh in 2008. The elevation above the mean sea level ranges between 313 and 483 m. The area experiences a tropical monsoon climate. The sites are located between two meteorological stations, Obra and Renukut. Ranitali site is nearest to Obra and Hathinala site is nearest to Renukoot. Mean annual rainfall is 926 mm at Obra and 1146 mm at Renukoot³⁴. The soils are residual ultisols, sandy loam in texture, reddish to dark grey in colour and are extremely poor in nutrients³⁵. Among the four sites, the mean rainy

*For correspondence. (e-mail: hema.bhu@gmail.com)

RESEARCH COMMUNICATIONS

season gravimetric soil water content was highest at Hathinala site and lowest at Ranitali site (Table 1).

Shorea robusta, *Buchanania lanzan*, *Diospyros melanoxylon*, *Lagerstroemia parviflora*, *Lannea coromandelica*, *Terminalia tomentosa*, *Holarrhena antidysenterica* and *Lantana camara* were selected for the study as these species occurred on all the four sites. Five fully grown leaves from each of the five individual plants were sampled ($n = 800$) for each species in September 2008. The leaves were collected from the unshaded mid-canopy region. After measuring the leaf area by a portable leaf area meter (SYSTRONICS, Leaf Area Meter-211), the leaves were oven-dried at 70°C for 72 h to obtain the dry mass with a precision electronic balance, to the nearest 0.001 g. SLA was calculated as the ratio of leaf area to leaf dry weight. For determination of chlorophyll content, five leaf samples were collected at the same time from the same five individuals and locations as above. The leaf samples were homogenized in 80% acetone, and absorbance at 663 and 645 nm was measured. The chlorophyll concentration was then calculated following Arnon³⁶. Data were subjected to multivariate ANOVA. Tukey's test was used to differentiate between means. Two-tailed Pearson correlation coefficients among leaf traits were calculated. All statistical analyses were done using SPSS (ver. 10) package. Species and sites were ordinated by correspondence analysis (CA) using the combination of leaf traits through Biodiversity-Pro (ver. 2).

All leaf traits examined had wide variability across species and sites (Table 2). The coefficient of variation was maximum for leaf weight (86%) followed by leaf area (82%), SLA (57%) and total chlorophyll content (46%). Thus total chlorophyll concentration had the least variation. The leaf traits across species and sites were significantly correlated with each other (Table 3), except for SLA which was independent of leaf area, although leaf weight which accounted for 75.3% variability in leaf area (Figure 1), showed a significant negative relationship with SLA. The positive relationship between leaf area and chlorophyll concentration is particularly meaningful for the seasonally dry tropics, as the canopies have to perform photosynthetically at their best during the short growth period. Many leaf traits have been found to be correlated in other studies also^{7,37}. At the species level, comparative studies of woody plants from a range of

environments, however, suggest that leaf size and SLA may not be closely linked^{20,38}. The lack of correlation between leaf area and SLA as observed in our study suggests that these two traits could be associated with different ecophysiological strategies or environmental factors, as argued by Ackerly *et al.*³⁹. SLA, for example, could be correlated with sclerophylly or thickness of cuticle, while leaf area could be related to boundary-layer resistance and photosynthetic water-use efficiency³⁹.

ANOVA indicated that all the four leaf traits had a significant species effect (Table 4). The mean leaf area varied from 26.39 cm² in *H. antidysenterica*, which had the minimum total chlorophyll concentration (0.72 mg/g), to 143.05 cm² in *T. tomentosa* (Table 5). On the other hand, mean leaf weight was minimum for *L. camara* (0.11 g/leaf), which had the greatest SLA (269.35 cm²/g), and maximum for *T. tomentosa* (1.38 g/leaf). Minimum SLA was exhibited by *D. melanoxylon* (88.15 cm²/g), which had the maximum total chlorophyll concentration (1.26 mg/g). Thus there was a marked interspecific variation in leaf traits: leaf area 5.4-fold, leaf weight 12.5-fold, SLA three-fold and chlorophyll content 1.7-fold. Other studies also indicate that the morphological traits such as leaf area and leaf mass are more variable than biochemical traits such as chlorophyll content⁴⁰. However, Tukey's test indicated that several species overlapped in individual leaf traits, except for *T. tomentosa* which was significantly different from all other species in terms of leaf area and leaf weight, *L. camara* in terms of leaf weight and SLA, *L. coromandelica* in terms of SLA and *D. melanoxylon* in total chlorophyll concentration (Table 5).

ANOVA also indicated a significant site effect on all four traits (Table 4). When the data on leaf traits were averaged for each site across the eight species, the overall

Table 1. Mean gravimetric soil water, total soil nitrogen and total soil carbon for the different sites, as recorded for the rainy season (R.K. Chaturvedi, unpublished)

Site	Gravimetric soil water (%)	Total soil nitrogen (%)	Total soil carbon (%)
Ranitali	16.10 ± 1.21	0.11 ± 0.01	1.52 ± 0.08
Neruiadamar	17.70 ± 1.09	0.11 ± 0.01	1.53 ± 0.08
Bokarakhari	18.10 ± 1.14	0.07 ± 0.01	1.20 ± 0.1
Hathinala	19.80 ± 1.11	0.09 ± 0.01	1.39 ± 0.18




Figure 1. Relationship between leaf area and leaf weight across the eight species at the four sites. The values were square-root transformed for the analysis. The regression line represents the equation $y = 1.574 + 8.4x$ ($r^2 = 0.753$, $P < 0.0001$, $n = 800$), where y is square-root leaf area and x is square-root leaf weight.

Table 2. Minimum, maximum, mean and standard deviation values for leaf traits across species and sites ($n = 800$)

Descriptive statistics	Minimum	Maximum	Mean	Standard deviation
Leaf area (cm ²)	4.90	314.30	70.26	57.81
Leaf weight (g)	0.03	3.21	0.65	0.56
Specific leaf area (cm ² /g)	9.74	546.81	133.34	76.10
Chlorophyll (mg/g)	0.21	2.54	0.91	0.42

Table 3. Pearson's correlation coefficients between leaf traits. The values for leaf traits were square-root transformed ($n = 800$)

	Leaf area	Leaf weight	Specific leaf area
Leaf weight (cm ²)	0.868**		
Specific leaf area (cm ² /g)	-0.015 ^{NS}	-0.449**	
Chlorophyll content (mg/g)	0.154**	0.149**	-0.081*

* $P < 0.05$; ** $P < 0.01$; NS, Not significant.




Figure 2. Plots of leaf traits against sites; the sites are scaled according to gravimetric soil water content: a, Leaf area; b, Leaf weight; c, Specific leaf area; d, Chlorophyll concentration.

variation was reduced (Table 6). Mean leaf area ranged from 57.21 cm² for the Bokarakhari site to 86.73 cm² for

Table 4. Summary of ANOVA on leaf traits

Main effect	Traits	df	F
Site	LA	3	29.47
	LW	3	13.74
	SLA	3	46.30
	CHL	3	27.85
Species	LA	7	202.67
	LW	7	206.93
	SLA	7	310.29
	CHL	7	50.90
Site × species	LA	21	12.02
	LW	21	7.02
	SLA	21	41.09
	CHL	21	30.14
Error	Each trait	768	

All *F* values were significant at *P* < 0.001. LA, leaf area; LW, Leaf weight; SLA, Specific leaf area; CHL, Chlorophyll content.

the Hathinala site (1.5-fold), mean leaf weight from 0.57 g/leaf at Neruiadamar to 0.77 g/leaf at Hathinala (1.3-fold), mean SLA from 123.10 cm²/g on Bokarakhari to 155.15 cm²/g at Hathinala (1.3-fold), and total chlorophyll concentration from 0.81 mg/g at Neruiadamar to 1.04 mg/g at Hathinala site (1.3-fold). The Hathinala site had the highest mean leaf trait values and was significantly different from all other sites. The Neruiadamar site was also significantly different from all other sites in terms of SLA and the Ranitalli site in terms of chlorophyll concentration (Table 6).

Interaction between species and site yielded significant differences in all the leaf traits (Table 4). The effect of site × species interaction was not consistent across leaf traits and majority of the species did not show marked dynamics with respect to sites (Figure 2). On the wettest site (Hathinala), *S. robusta* and *D. melanoxylon* showed clear peaks in mean leaf area, *T. tomentosa* and *D. melanoxylon* in leaf weight; *L. camara*, *L. coromandelica* and *S. robusta* in SLA, and *B. lanzan* and *D. melanoxylon* in chlorophyll concentration. On the other hand, *H. antidysenterica* showed peaks in leaf area and SLA on the driest site (Ranitalli).

CA on the basis of the set of leaf traits indicated three broad groups of species (Figure 3 a). The groups generally corresponded to leaf size (large leaf size, ≥ 107 cm²: *T. tomentosa*, *S. robusta* and *B. lanza*; smaller leaf size, 26.37–61.48 cm²: *H. antidysenterica*, *L. parviflora*, *D. melanoxylon* and *L. coromandelica*), except for *L. camara* having small leaf size (29.77 cm²) but high SLA (269.35 cm²/g), which stood alone. It is important to note that *L. camara* is a non-native invasive species in the dry tropical forests of India and is spreading fast in open-canopy areas. Global-scale comparisons also show that exotic invasive species generally have significantly higher values of SLA compared to native species⁴¹. Leaves that have high SLA can produce large or more assimilatory surfaces for a given amount of carbon fixed⁴². Also, SLA is positively related to plant growth and efficiency because it relates to leaf thickness and longevity⁴³. Thus exotic species like *Lantana* achieve a greater C gain for a given investment in leaves, resulting in an even greater return on investment and hence faster growth.

A similar analysis on site based-leaf traits indicated that a combination of leaf traits can discriminate between the sites. The sites remained scattered and distinct in the ordination space (Figure 3 b). Between-site variability in leaf traits was smaller (1.3–1.5-fold) than between-species variability (1.7–12.5-fold). The larger inter-species variability reflects marked genotypic variability in the leaf traits, while the smaller between-site variability reflects phenotypic plasticity leading to adaptation to site conditions. For example, smallest cross-species SLA on the driest Ranitalli site may reflect water stress, which is known to reduce SLA⁴⁴. The relationships among the leaf traits could contribute to the relative consistency of com-




Figure 3. a, Correspondence analysis (CA) ordination of species on the basis of leaf traits: *Bla*, *Buchanania lanzan*; *Dme*, *Diospyros melanoxylon*; *Han*, *Holarrhena antidysenterica*; *Lca*, *Lantana camara*; *Lco*, *Lannea coromandelica*; *Lpa*, *Lagerstroemia parviflora*; *Sro*, *Shorea robusta* and *Tto*, *Terminalia tomentosa*. b, CA ordination of four sites on the basis of leaf traits: *Bok*, Bokarakhari; *Hat*, Hathinala; *Ner*, Neruiadamar; *Ran*, Ranitalli.

Table 5. Mean leaf area, leaf weight, specific leaf area and chlorophyll content of eight species across four sites

Species	Leaf area (cm ²)	Leaf weight (g)	Specific leaf area (cm ² /g)	Chlorophyll (mg/g)
<i>Shorea robusta</i>	118.33 ^a	1.09 ^a	108.86 ^a	1.07 ^a
<i>Buchanania lanzan</i>	107.39 ^a	1.031 ^a	108.27 ^a	1.04 ^a
<i>Diospyros melanoxylon</i>	61.48 ^b	0.70 ^b	88.15 ^b	1.26 ^b
<i>Lagerstroemia parviflora</i>	26.51 ^c	0.30 ^c	89.67 ^b	0.82 ^{dc}
<i>Lannea coromandelica</i>	49.13 ^b	0.31 ^c	168.92 ^c	0.90 ^d
<i>Terminalia tomentosa</i>	143.05 ^d	1.38 ^d	118.42 ^a	0.73 ^c
<i>Holarrhena antidysenterica</i>	26.39 ^c	0.26 ^c	114.90 ^a	0.72 ^c
<i>Lantana camara</i>	29.77 ^c	0.11 ^e	269.35 ^d	0.74 ^c

Values suffixed with the same letter in a column were not significantly different from each other.

Table 6. Mean leaf area, leaf weight, specific leaf area and chlorophyll content of plants on a site across species

Site	Leaf area (cm ²)	Leaf weight (g)	Specific leaf area (cm ² /g)	Chlorophyll (mg/g)
Ranitalli	71.67 ^a	0.66 ^a	118.48 ^a	0.93 ^a
Neruiadamar	65.41 ^{ba}	0.56 ^b	136.53 ^b	0.81 ^b
Bokarakhari	57.21 ^b	0.60 ^{ba}	123.10 ^a	0.84 ^b
Hathinala	86.73 ^c	0.76 ^c	155.15 ^c	1.04 ^c

Values suffixed with the same letter in a column were not significantly different from each other.

bined traits at the site level. For example, as SLA increases, leaf weight and chlorophyll concentration decrease. Thus there is a trade-off between the leaf traits.

In conclusion, we suggest that a combination of cross-species leaf traits indeed reflects site conditions in the dry tropical environment. The SLA and chlorophyll concentration are more stable traits than leaf area or leaf weight, because of their lower inter-species and inter-site variability.

- Westoby, M., A leaf height seed (LHS) plant ecology strategy scheme. *Plant Soil*, 1998, **199**, 213–227.
- Vendramini, F., Diaz, S., Gurrich, D. E., Wilson, P. J., Thompson, K. and Hodgson, J. G., Leaf traits as indicators of resource-use strategy in floras with succulent species. *New Phytol.*, 2002, **154**, 147–157.
- Weihner, E., The combined effects of scale and productivity on species richness. *J. Ecol.*, 1999, **87**, 1005–1011.
- Craine, J. M., Froehle, J., Tilman, D. G., Wedin, D. A. and Chapin, F. S., The relationships among root and leaf traits of 76 grassland species and relative abundance along fertility and disturbance gradients. *Oikos*, 2001, **93**, 274–285.
- Reich, P. B., Walters, M. B. and Ellsworth, D. S., Leaf life-span in relation to leaf, plant, and stand characteristics among diverse ecosystems. *Ecol. Monogr.*, 1992, **62**, 365–392.
- Van der Werf, A., Geerts, R. H. E. M., Jacobs, F. H. H., Korevaar, H., Oomes, M. J. M. and De Visser, W., The importance of relative growth rate and associated traits for competition between species during vegetation succession. In *Inherent Variation in Plant Growth, Physiological Mechanisms and Ecological Consequences* (eds Lambers, H., Poorter, H. and Van Vuuren, M. M. I.), Backhuys, Leiden, The Netherlands, 1998, pp. 489–502.
- Wilson, P. J., Thompson, K. and Hodgson, J. G., Specific leaf area and leaf dry matter content as alternative predictors of plant strategies. *New Phytol.*, 1999, **143**, 155–162.
- Garnier, E. *et al.*, Consistency of species ranking based on functional leaf traits. *New Phytol.*, 2001, **152**, 69–83.
- Poorter, H. and Van der Werf, A., Is inherent variation in RGR determined by LAR at low irradiance and by NAR at high irradiance? A review of herbaceous species. In *Inherent Variation in Plant Growth, Physiological Mechanisms and Ecological Consequences* (eds Lambers, H., Poorter, H. and Van Vuuren, M. M. I.), Backhuys, Leiden, The Netherlands, 1998, pp. 309–336.
- Coley, P. D., Bryant, J. P. and Chapin III, F. S., Resource availability and plant anti-herbivore defence. *Science*, 1985, **230**, 895–885.
- Grime, J. P., Cornelissen, J. H. C., Thompson, K. and Hodgson, J. G., Evidence of a causal connection between anti-herbivore defence and the decomposition rate of leaves. *Oikos*, 1996, **77**, 489–494.
- Aase, J. K., Relationship between leaf area and dry matter in winter wheat. *Agron. J.*, 1978, **70**, 563–565.
- Smart, R. E., Principles of grape vine canopy microclimate manipulation with implications for yield and quality: a review. *Am. J. Enol. Vitic.*, 1985, **36**, 230–239.
- Williams, L. E., Growth of ‘Thompson Seedless’ grape vines. I., Leaf area development and dry weight distribution. *J. Am. Hort. Sci.*, 1987, **112**, 325–330.
- Hamann, O., On climatic conditions, vegetation types and leaf size in the Galapagos Islands. *Biotropica*, 1979, **11**, 101–122.
- Dolph, G. and Dilcher, D., Variation in leaf size with respect to climate in the tropics of the western hemisphere. *Bull. Torrey Bot. Club*, 1980, **107**, 154–162.
- Givnish, T. J., Leaf and canopy adaptation in tropical forest. In *Physiological Ecology of Plants of the Wet Tropics* (eds Medina, E., Mooney, H. A. and Vazquez-Yanes, C.), Junk, The Hague, 1984, pp. 51–84.
- Skarpe, C., Plant functional types and climate in southern African Savanna. *J. Veg. Sci.*, 1996, **7**, 397–404.
- Cunningham, S. A., Summerhayes, B. A. and Westoby, M., Evolutionary divergences in leaf structure and chemistry, comparing rainfall and soil nutrient gradients. *Ecol. Monogr.*, 1999, **69**, 569–588.

20. Fonseca, C. R., Overton, J. M., Collins, B. and Westoby, M., Shifts in trait-combinations along rainfall and phosphorus gradients. *J. Ecol.*, 2000, **88**, 964–977.
21. Mooney, H. A. and Dunn, E. L., Convergent evolution of Mediterranean-climate evergreen sclerophyllous shrub. *Evolution*, 1970, **24**, 292–303.
22. Parkhurst, D. and Loucks, O., Optimal leaf size in relation to environment. *J. Ecol.*, 1972, **60**, 505–537.
23. Givinish, J. J. and Vermeiji, G. J., Sizes and shapes of liane leaves. *Am. Nat.*, 1976, **110**, 743–778.
24. Evans, J. R., Photosynthesis and nitrogen relationships in leaves of C₃ plants. *Oecologia*, 1989, **78**, 9–19.
25. Yoder, B. J. and Pettigrew-Crosby, R. E., Predicting nitrogen and chlorophyll content and concentrations from reflectance spectra (400–2500 nm) at leaf and canopy scales. *Remote Sensing Environ.*, 1995, **53**, 199–211.
26. Niinemets, U. and Tenhunen, J. D., A model separating leaf structural and physiological effects on carbon gain along light gradients for the shade-tolerant species *Acer saccharum*. *Plant Cell Environ.*, 1997, **20**, 845–866.
27. Whittaker, R. H. and Marks, P. L., Methods of assessing terrestrial productivity. In *Primary Productivity of the Biosphere* (eds Lieth, H. and Whittaker, R. H.), Springer-Verlag, New York, 1975, pp. 55–118.
28. Darks, S. M., Evans, E. H. and Whittaker, P. A., *Photosynthetic systems. Structure Function and Assembly*, Wiley, New York, 1983.
29. Ahmad, I. S., Reich, J. F., Noguchi, N. and Hansen, A. C., Nitrogen sensing for precision agriculture using chlorophyll maps. ASAE/CSAE-SCGR Annual International Meeting, Toronto, Canada, 18–21 July 1999.
30. Zarco-Tejada, P. J., Miller, J. R., Morales, A., Berjon, A. and Aguera, J., Hyperspectral indices and model simulation for chlorophyll estimation in open-canopy tree crops. *Remote Sensing Environ.*, 2004, **90**, 463–476.
31. Peterson, T. A., Blackmer, T. M., Francis, D. D. and Schepers, J. S., Using a chlorophyll meter to improve N management. *Nebguide G93-1171A. Cooperative Extension Services*, University of Nebraska, Lincoln, 1993.
32. Wood, C. W., Reeves, D. W. and Himelrick, D. G., Relationships between chlorophyll meter readings and leaf chlorophyll concentration, N status, and crop yield: a review. *Proc. Agron. Soc., NZ*, 1993, **23**, 1–9.
33. Blackmer, T. M., Schepers, J. S. and Varvel, G. E., Light reflectance compared with other nitrogen stress measurements in corn leaves. *Agron. J.*, 1994, **86**, 934–938.
34. Pandey, C. B. and Singh, J. S., Rainfall and grazing effects on net primary productivity in a tropical savanna, India. *Ecology*, 1992, **73**, 2007–2021.
35. Singh, J. S., Raghubanshi, A. S., Singh, R. S. and Srivastava, S. C., Microbial biomass acts as a source of plant nutrient in dry tropical forest and savanna. *Nature*, 1989, **338**, 499–500.
36. Aron, D. I., Copper enzymes in isolated chloroplasts. (I) Polyphenoloxidase in *B. vulgaris*. *Plant Physiol.*, 1949, **24**, 1–15.
37. Roche, P., Diaz-Burlinson, N. and Gachet, S., Congruency analysis of species ranking based on leaf traits: which traits are more reliable? *Plant Ecol.*, 2004, **174**, 37–48.
38. Ackerly, D. D. and Reich, P. B., Convergence and correlations among leaf size and function in seed plants: a comparative test using independent contrast. *Am. J. Bot.*, 1999, **86**, 1272–1281.
39. Ackerly, D. D., Knight, C. A., Weiss, S. B., Barton, K. and Starmer, K. P., Leaf size, specific leaf area and microhabitat distribution of chaparral woody plants: contrasting patterns in species level and community level analyses. *Oecologia*, 2002, **130**, 449–457.
40. Vincent, G., Leaf photosynthetic capacity and nitrogen content adjustment to canopy openness in tropical forest tree seedlings. *J. Trop. Ecol.*, 2001, **17**, 495–509.
41. Leishman, M. R., Haslehurst, T. Ares, A. and Baruch, Z., Leaf trait relationships of native and invasive plants: community- and global-scale comparisons. *New Phytol.*, 2007, **176**, 635–643.
42. Chiariello, N. R., Mooney, H. A. and Williams, K., Growth, carbon allocation and cost of plant tissues. In *Plant Physiological Ecology* (eds Percy, R. W. *et al.*), Chapman and Hall, London, 1991, pp. 327–366.
43. Baruch, Z. and Goldstein, G., Leaf construction cost, nutrient concentration and net CO₂ assimilation of native and invasive species in Hawaii. *Oecologia*, 1999, **121**, 183–192.
44. Khurana, E. and Singh, J. S., Germination and seedling growth of five tree species from tropical dryforest in relation to water stress: impact of seed size. *J. Trop. Ecol.*, 2004, **20**, 385–396.

ACKNOWLEDGEMENTS. We thank the Ministry of Environment and Forests, Government of India for providing financial assistance. J.S.S. is supported under NASI Senior Scientist Scheme.

Received 5 January 2009; accepted 19 March 2009

Dimethyl sulphoxide inhibits *in vitro* synthesis of juvenile hormone I and II and stimulates juvenile hormone III by corpora allata of insects[†]

B. G. Unni^{1,*}, Ken E. Peck², M. M. Pytel², K. H. Dahm², G. Bhaskaran², H. Ranjit Singh¹, Y. Kakoty¹, B. Devi¹ and S. B. Wann¹

¹Biotechnology Division, North East Institute of Science and Technology (formerly Regional Research Laboratory), Jorhat 785 006, India

²Department of Biology, Texas A&M University, College Station, Texas 71843, USA

Dimethyl sulphoxide (DMSO) specifically inhibits the rates of biosynthesis of juvenile hormone I (JH I) and juvenile hormone II (JH II) in corpora allata (CA) of 0 day last instar larvae of *Manduca sexta* maintained *in vitro*. Addition of mevalonate (Mev), propionate and homomevalonate (HMeV) to the medium increased the rate of biosynthesis of both JH II and JH III in normal CA. Even with added Mev, DMSO drastically reduced the rate of synthesis of both the higher homologues, but enhanced the rate of JH III synthesis even more with DMSO alone. The mated adult female CA which secretes JH II and JH III, and almost no JH I, also showed similar pattern of response to DMSO in larval CA. Allatotropin-stimulated JH II synthesis was also blocked by DMSO. DMSO inhibited JH II synthesis in *Galleria mellonella*, but the rate of JH III synthesis showed no increase. Likewise, in the CA of

[†]This research communication is dedicated to professors Govindan Bhaskaran and Karl H. Dahm, Department of Biology, Texas A&M University, USA, for guidance towards the field of endocrinology and juvenile hormone research.

*For correspondence. (e-mail: bgunni@yahoo.com)