

19. FAO, Crop evapotranspiration: guidelines for computing crop requirements. FAO Irrigation and Drainage Paper 56, 1998, pp. 57–139.
20. Rao, B. K. and Rajput, T. B. S., Effective rainfall and its significance in crop water requirements in canal command areas. *J. Water Manage.*, 2005, **13**, 36–42.
21. Hajilal, M. S., Rao, N. H. and Sarma, P. B. S., Real time operation of reservoir based irrigation systems. Central Board of Irrigation and Power, New Delhi, Publication no. 256, 1997, p. 370.

ACKNOWLEDGEMENTS. The first author acknowledges the financial assistance from IARI, New Delhi, in the form of Senior Research Fellowship towards his doctoral programme. We thank the officials of Nagarjuna Sager Project, Khammam Division, Andhra Pradesh, for providing necessary data for this study.

Received 5 August 2008; revised accepted 22 May 2009

Physico-chemical, biochemical and microbial characteristics of soils of mangroves of the Andamans: a post-tsunami analysis

S. Ghoshal Chaudhuri^{1,*}, R. Dinesh²,
T. E. Sheeja², R. Raja¹, V. Jeykumar¹ and
R. C. Srivastava¹

¹Division of Natural Resources Management,
Central Agricultural Research Institute (ICAR),
Port Blair, PB #181, Andamans 744 101, India

²Indian Institute of Spices Research (ICAR),
Marikunnu PO, Calicut 673 012, India

The mangroves of the Bay Islands accounting for 18% (383 km²) of the total Indian mangroves were adversely affected by the December 2004 tsunami. Changes in topography, salinity and massive water inflow have led to extensive soil degradation and associated nutrient losses in these mangroves. The major aim of the study was to determine whether the December 2004 tsunami had any effects on soil physico-chemical (pH, electrical conductivity of saturation extract (EC), clay, cation exchange capacity (CEC), organic carbon (OC), total N (TN), Bray phosphorus (P), exchangeable cations (Ca, Mg, K and Na)) and biochemical/microbial parameters (Microbial biomass-C (C_{MIC}), -N (N_{MIC}), N-flush, basal respiration and hydrolytic enzyme activities). The post-tsunami soil samples (disturbed sites) were characterized by higher levels of EC, Na and Mg, while the pre-tsunami soils samples (undisturbed sites) had higher levels of OC, P, K and CEC. The study also revealed marked reductions in microbial biomass and activity in the disturbed sites. C_{MIC}, N_{MIC}, N-flush, basal respiration, and activities of hydrolytic enzymes like BAA-protease,

casein-protease, phosphomonoesterase, β -glucosidase, arylsulphatase, invertase, carboxy methyl cellulase and dehydrogenase were considerably lower in the disturbed sites. Higher levels of metabolic quotient (qCO₂) in the disturbed soils indicated comparatively more stressed soil microbial community with reduced substrate utilization efficiency. Apparently, microbial activity was limited by the supply of biologically available substrates like OC in the disturbed sites. Contrarily, the more direct supply of nutrients from decomposing plant litter and the indirect supply of nutrients from the mineralization of organic matter led to significantly higher microbial activity in the undisturbed sites.

Keywords: Biochemical properties, mangrove forests, microbial biomass carbon, soil enzymes, soil microbial activity.

MANGROVES cover an area of around 15 mha (or 1,50,000 km²) worldwide, with close to 40% of this area found in the countries affected by the December 2004 tsunami. Presently, less than 50% of the area remains, and of this over 50% is degraded due mainly to anthropogenic factors like conversion to fish ponds, agricultural land, etc.¹. Of the country's total area under mangrove vegetation, 70% is recorded on the east coast and 12% on the west coast. The Bay Islands (Andaman and Nicobar) account for 18% of the country's total mangrove area^{2,3}. The insular mangroves exist in the Bay Islands on many tidal estuaries, small rivers, neritic islets and lagoons, accounting for 18% (383 km²) of the total Indian mangroves. As would be expected, the mangroves along the Andamans coast were adversely affected by the 2004 tsunami.

However, the extent of damage is still not clear and it may take sometime before the final impacts are known, since the deposit of silt may clog the pores of the aerial roots of mangroves, and thus suffocate them. Changes in topography, soil salinity and freshwater inflow from upstream may also adversely affect the mangroves and other coastal forests in the longer term (<http://www.fao.org/newsroom/en/news/2005/89119/index.html> accessed on 01/02/2008). One of the major consequences of the tsunami is the extensive soil degradation and associated nutrient losses.

In order to minimize soil degradation and to adopt management techniques that contribute to the maintenance or recovery of soil fertility, the soil quality should be ascertained in order to understand the limits that can be set to its use and treatment. Of all the parameters that determine soil quality, biological and biochemical variables are the most sensitive to changes occurring in a soil^{4,5} and provide rapid and accurate information on changes in soil quality due to their sensitivity to environment stress⁶, role in degradation⁷ and strong influence on microbially mediated processes like nutrient cycling,

*For correspondence. (e-mail: sgc2004@rediffmail.com)

nutrient capacity and aggregate stability⁸. Sufficient data on these parameters will, therefore, be immensely useful for detecting possible degradation and estimating the degree of recovery of degraded mangrove soils during rehabilitation.

This study reports the results of post-tsunami analysis on soil biochemical/microbial properties of mangrove forests of the Bay Islands. The aim was to determine the effect of the tsunami on soil physico-chemical and biochemical/microbial properties of the mangrove forests in the region. For the study, soil cores (0–30 cm) were taken from ten randomly selected spots from within the intertidal zone of ten mangrove forest sites in March 2006. All core samples were collected at low tide. The tsunami hit (disturbed) mangrove sites selected for the study are Manjeery, Burmanalha, Kodyaghat, Sippighat, Wandoor, Shole Bay, Kalataung, Yerata, Barataung and Potataung. The soils were then sieved (<2 mm), analysed for moisture content and stored at 4°C for not more than one week before analyses. Sub-samples for the determination of organic C and total N were sieved to pass a 0.5 mm mesh. Electrical conductivity (EC) was measured in saturated extracts using a conductivity meter. A combined glass-calomel electrode was used to determine the pH of aqueous suspensions (1 : 2.5 solid/liquid ratio).

Before chemical analysis, samples were washed with 60% (v/v) aqueous ethanol to remove salts. Total N and organic C were determined using standard methods⁹. Clay content was determined by the pipette method¹⁰ and cation exchange capacity (CEC) by the method of Gillman¹¹. Exchangeable cations (Ca, Mg, K and Na) were extracted in ammonium acetate¹², followed by determination of Ca, Mg and K by atomic absorption and Na by flame photometry. Bray P was measured using the dilute-acid fluoride extractant¹³ and soil Al and Fe using the spectrophotometric methods described by Barnhisel and Bertsch¹⁴ and Olson and Ellis¹⁵ respectively.

Microbial biomass C (C_{MIC}) and N (N_{MIC}) were determined using the chloroform fumigation technique¹⁶, utili-

zing a factor of 0.45 (refs 17 and 18) and 0.54 (refs 19 and 20) respectively. Nitrogen flush was calculated from the difference between ninhydrin-reactive N in fumigated and non-fumigated samples²¹. Basal respiration (CO_2 evolution) was determined by adjusting the moisture content to 55% of water-holding capacity and pre-incubating the soil samples, as described by Salamanca *et al.*²². Dehydrogenase was assayed using 2,3-5 triphenyl tetrazolium chloride as the substrate²³, urease using urea as the substrate²⁴, BAA-protease and casein-protease using α -benzoyl-*N*-argininamide (BAA) and casein respectively, as substrates²⁵, acid phosphomonoesterase using *p*-nitrophenol phosphate as the substrate²⁶, β -glucosidase using *p*-nitrophenyl- β -D-glucopyranoside as the substrate²⁷, phosphodiesterase using *bis-p*-nitrophenyl phosphate as the substrate²⁸, arylsulphatase using *p*-nitrophenyl sulphate as the substrate²⁹, invertase using saccharose as the substrate³⁰, and carboxy methyl cellulase using carboxymethyl cellulose as the substrate³⁰.

Maxima, minima, means and standard deviations (SD) of the soil variables are expressed on an oven-dry soil basis (105°C). The results were then compared with our earlier unpublished data on soils collected from the same mangrove sites in March 2004. The relationships between the different parameters were determined by the Pearson's correlation test and the interdependence of the soils variables was determined by principal component analysis (PCA).

For convenience, in the subsequent sections the tsunami-hit mangrove forest sites will hitherto be referred to as disturbed sites. Soil pH was acidic and exhibited little variation among the sites (Table 1). Electrical conductivity (EC of the saturation extract) was considerably higher at the disturbed sites (33.8 ± 1.5 dS m^{-1}), indicating accumulation of salts. In contrast, Al_2O_3 and Fe_2O_3 showed very little variation compared to CEC, which declined marginally from 236 ± 31 to 225 ± 23 mmol kg^{-1} due to disturbance. On the contrary, the clay content was lower by 35%, Bray P by 31%, and organic C by 25% at

Table 1. Relevant properties of soils at the undisturbed and tsunami-hit mangrove sites

	Undisturbed sites			Tsunami-hit sites		
	Minimum	Maximum	Mean \pm SD	Minimum	Maximum	Mean \pm SD
pH	4.1	6.7	5.3 \pm 1.00	3.6	6.2	4.8 \pm 0.95
EC (dS m^{-1})	20.3	25.6	23.2 \pm 1.70	31.3	36.2	33.8 \pm 1.5
Total N (mg g^{-1})	2.01	2.81	2.34 \pm 0.24	2.0	2.85	2.36 \pm 0.31
Organic C (mg g^{-1})	14.1	24.6	19.1 \pm 3.4	9.9	22.4	14.3 \pm 4.3
CEC (mmol kg^{-1})	202	286	236 \pm 31	189	275	225 \pm 23
Bray P (mg 100 g^{-1})	19.2	37.4	25.6 \pm 6.1	14.7	20.9	17.7 \pm 2.3
Exchangeable K (Cmol _c kg^{-1})	1.12	1.35	1.25 \pm 0.08	0.81	1.06	0.90 \pm 0.07
Exchangeable Ca (Cmol _c kg^{-1})	6.7	7.9	7.0 \pm 0.6	6.1	7.4	6.7 \pm 0.4
Exchangeable Mg (Cmol _c kg^{-1})	3.2	3.8	3.4 \pm 0.2	4.1	4.9	4.5 \pm 0.2
Exchangeable Na (Cmol _c kg^{-1})	8.2	12.7	10.9 \pm 1.5	10.3	15.8	13.8 \pm 1.7
Al_2O_3 (%)	0.31	1.55	1.12 \pm 0.36	0.33	1.45	1.14 \pm 0.24
Fe_2O_3 (%)	0.21	1.63	1.14 \pm 0.29	0.15	1.34	1.10 \pm 0.14
Clay (%)	19	27	23 \pm 3	11	19	15 \pm 3

RESEARCH COMMUNICATIONS

Table 2. Values of soil microbial and biochemical parameters at the undisturbed and tsunami-hit mangrove sites

	Undisturbed sites			Tsunami-hit sites		
	Minimum	Maximum	Mean \pm SD	Minimum	Maximum	Mean \pm SD
Microbial biomass C ($\mu\text{g g}^{-1}$)	370	504	426 \pm 50	242	352	286 \pm 36
Microbial biomass N ($\mu\text{g g}^{-1}$)	33.2	48.8	39.6 \pm 4.6	23.0	30.0	27.3 \pm 2.6
N flush ($\mu\text{g g}^{-1}$)	7.6	14.9	11.9 \pm 2.8	5.6	12.8	9.8 \pm 1.6
Basal respiration ($\mu\text{g CO}_2\text{-C g}^{-1}$ per day)	11.6	16.2	13.2 \pm 2.3	8.9	13.5	10.9 \pm 1.7
qCO ₂ (mg CO ₂ -C g ⁻¹ biomass C per day)	23.9	47.6	31.0 \pm 5.2	23.4	44.6	38.0 \pm 3.2
Dehydrogenase (nmol TPF g ⁻¹ h ⁻¹)	253	513	412 \pm 86	121	278	212 \pm 42
Catalase (mmol H ₂ O ₂ consumed g ⁻¹ h ⁻¹)	1.76	4.51	3.53 \pm 0.87	0.81	1.46	1.21 \pm 0.19
Phosphomonoesterase ($\mu\text{mol p-nitrophenol g}^{-1}$ h ⁻¹)	11.5	20.4	15.48 \pm 2.90	5.2	9.4	7.34 \pm 1.43
Phosphodiesterase ($\mu\text{mol p-nitrophenol g}^{-1}$ h ⁻¹)	3.5	6.2	5.03 \pm 1.02	3.1	5.7	4.54 \pm 0.95
Arylsulphatase ($\mu\text{mol p-nitrophenol g}^{-1}$ h ⁻¹)	0.51	1.00	0.73 \pm 0.18	0.46	1.23	0.68 \pm 0.11
Urease ($\mu\text{mol NH}_3\text{-N g}^{-1}$ h ⁻¹)	6.0	17.0	12.8 \pm 3.0	5.8	15.7	11.7 \pm 3.1
BAA-protease ($\mu\text{mol NH}_3\text{-Ng}^{-1}$ h ⁻¹)	6.33	8.74	7.39 \pm 0.90	4.66	8.18	6.35 \pm 1.09
Casein-protease ($\mu\text{mol tyrosine g}^{-1}$ h ⁻¹)	2.3	4.6	3.6 \pm 0.4	1.2	2.5	2.0 \pm 0.4
β -Glucosidase ($\mu\text{mol p-nitrophenol g}^{-1}$ h ⁻¹)	3.4	6.7	5.5 \pm 1.1	2.1	5.2	3.7 \pm 0.9
CM cellulase ($\mu\text{mol glucose g}^{-1}$ h ⁻¹)	0.61	0.94	0.78 \pm 0.11	0.25	0.76	0.43 \pm 0.10
Invertase ($\mu\text{mol glucose g}^{-1}$ h ⁻¹)	7.8	15.7	12.4 \pm 2.7	5.4	13.7	9.8 \pm 2.1

the disturbed sites. Gomez and Fortes³¹ also observed greater organic matter levels in the surface soils of undisturbed mangrove forests of Pagbilao, Quezon. Lower clay and organic C levels at the disturbed sites indicated washing away of finer soil particles, organic fractions and litter by the flood. On the contrary, greater levels of organic C and nutrients like P and K at the undisturbed sites are due in large part to nutrient regeneration from fallen leaves, twigs, buds, flowers, decaying roots, etc³². Among the exchangeable cations, Ca did not show much variation among the sites, while K was greater at the undisturbed sites. On the contrary, Mg and Na levels were markedly higher at the disturbed sites. This finding is consistent with the possibility that overflowings during tsunami were relatively high in Na and Mg ions, which then accumulated near or on the soil surface.

Among the biochemical parameters, C_{MIC} level at the undisturbed sites was on an average 426 \pm 50, which decreased to 286 \pm 36 $\mu\text{g g}^{-1}$, indicating a 33% decline (Table 2), possibly due to decreased microbial activity in response to lower levels of organic C and nutrients in the disturbed sites. Earlier studies on total litter productivity at various mangrove sites of the Andamans indicate considerably lower annual litter fall at the disturbed mangrove sites (472 g dry wt m⁻²) compared to the undisturbed sites (830 g dry wt m⁻²)³³. Since microbial biomass in forest soils is largely governed by litter fall³⁴, greater microbial biomass at the undisturbed sites reflects greater accumulation of plant residues and organic C, which are substrates for soil microbes³⁵. This is reflected in the existence of a strong correlation between microbial biomass and organic C ($r = 0.76$ at $P < 0.05$; $n = 200$). This is expected since both are part of the organic matter pool³⁶ and microbial biomass is influenced more quickly by organic inputs than by changes in organic matter³⁷.

Similar to C_{MIC}, the average levels of N_{MIC} and N flush at the disturbed sites (27.3 \pm 2.6 and 9.8 \pm 1.6 $\mu\text{g g}^{-1}$ respectively) were lower by 31 and 18% compared to the undisturbed sites (39.6 \pm 4.6 and 11.9 \pm 2.8 $\mu\text{g g}^{-1}$ respectively; Table 2). Since N flush is a measure of the microbial biomass, a small biomass would have a low N flush³⁸. Therefore, greater N flush values at the undisturbed sites indicate greater microbial activity and microbial biomass as reflected by the high correlation between C_{MIC} and N flush ($r = 0.78$; $P < 0.01$). Lower levels of C_{MIC} and microbial activity led to lower basal respiration rates in the disturbed soils. Average CO₂ evolution rates declined by 17% (10.9 \pm 1.7 $\mu\text{g CO}_2\text{-C g}^{-1}$ per day) in the disturbed sites compared to the undisturbed sites (13.2 \pm 2.3 $\mu\text{g CO}_2\text{-C g}^{-1}$ per day). On the contrary, metabolic quotient (qCO₂) was higher by 23% in the disturbed sites (38.0 \pm 3.2) compared to the undisturbed sites (31.0 \pm 5.2). Higher qCO₂ in the disturbed soils indicated microbial community with less catabolic activity³⁹ and reduced efficiency of substrate utilization or high energy requirements and comparatively more stressed microbial community which was less metabolically efficient⁴⁰.

The results pertaining to soil enzyme activities were also identical to those of microbial biomass. The activities of these enzymes were reduced to varying degrees at the disturbed sites (Table 2). Accumulation/stabilization of organic matter⁴¹ coupled with greater microbial activity⁴² might explain the greater levels of enzyme activities at the undisturbed sites. On the contrary, loss in nutrients, organic matter and dissolved organic substrates due to forced water flow would have reduced microbial activity and therefore decreased enzyme activity in the disturbed sites. Numerous studies have shown that soil microbial community is influenced by organic C, which

is the principal factor influencing growth and activity of microbial community⁴³. Also, in soils under fluvial systems, the sediments are transported in suspension and in bed-load together with bacteria adhering to the transported particles⁴⁴ leading to reduced microbial load and enzyme activity. Besides, reduced microbial/enzyme activity at the disturbed sites would also be due to excess salt accumulation (as evidenced by markedly higher EC levels) and osmotic desiccation leading to microbial cell lysis, and a salting-out effect modifying the ionic conformation of the active site of the enzyme protein⁴⁵.

In order to clarify the structure of interdependence of various biochemical parameters and soil variables, a joint principal components analysis was done (Table 3). The first factor was loaded by C_{MIC} , N_{MIC} , basal respiration, N flush, dehydrogenase, catalase, β -glucosidase, invertase, casein-protease, OC, Bray P and K, indicating the size and activity of the microbial community and the logical dependence of soil microbes on nutrient content. The high loadings of invertase and casein-protease also suggest that the activities of both these enzymes depend on the microbial community than on their extracellular accumulation in soil. Phosphomonoesterase CM-cellulose, organic C and total N were combined as the second factor characterizing accumulation of hydrolytic enzymes and

organic matter in mangrove soils. The third factor was defined by qCO_2 reflecting specific metabolic activity of soil microbes. The high loadings of urease and BAA-protease in the third factor confirmed the independence of the degradation of low molecular weight nitrogen compounds in these soils.

Overall, it is apparent that microbial biomass and activity were markedly higher at the undisturbed sites possibly due to greater levels of labile soil organic matter⁴⁵. Apparently destruction of mangrove vegetation by the tsunami led to decreased litter additions to the soil in the disturbed sites. Consequently, clay content, nutrient (Bray P, exchangeable K) and organic C levels in the soils decreased at the disturbed sites. Besides, the levels of soil C_{MIC} , N_{MIC} , N flush, CO_2 evolution rates and enzyme activities were also reduced by varying degrees. Decreased litter addition/accumulation coupled with loss in organic matter and therefore, microbial activity seems to be the likely reason for lower levels of microbial biomass and enzyme activities in the sites ravaged by the tsunami. The study reveals a clear cause-and-effect mechanism between tsunami and soil quality deterioration at these mangrove sites. This would mean a decrease in microbial-mediated processes like nutrient cycling, nutrient capacity and aggregate stability in the disturbed soils. While the results are important in detecting the extent of soil degradation and could be used for post-tsunami reconstruction of the mangrove ecosystem in the Bay islands, such degradation could be transient in regions where mangrove forest restoration is considered imperative. However, in regions where little effort is made for post-tsunami reconstruction the consequences could be serious, leading to long-term damage to the mangrove ecosystem.

Table 3. Oblique solution primary pattern matrix of the principal components analysis (orthotran/varimax transformation; $n = 200$) for the different physico-chemical and biochemical and microbial properties of the mangrove soils

	Factor 1	Factor 2	Factor 3
C_{MIC}	0.72	n.s.	n.s.
N_{MIC}	0.74	n.s.	n.s.
Basal respiration	0.64	n.s.	n.s.
N flush	0.69	n.s.	n.s.
qCO_2	n.s.	n.s.	0.74
Dehydrogenase	0.75	n.s.	n.s.
Catalase	0.66	n.s.	n.s.
Phosphomonoesterase	n.s.	0.71	n.s.
Phosphodiesterase	n.s.	n.s.	n.s.
Arylsulphatase	n.s.	n.s.	n.s.
Urease	n.s.	n.s.	0.64
BAA-protease	n.s.	n.s.	0.68
β -Glucosidase	0.65	n.s.	n.s.
CM cellulase	n.s.	0.71	n.s.
Invertase	0.66	n.s.	n.s.
Casein -protease	0.62	n.s.	n.s.
PH	n.s.	n.s.	n.s.
CEC	n.s.	n.s.	n.s.
Clay	n.s.	n.s.	n.s.
Organic C	0.64	0.55	n.s.
Total N	n.s.	0.70	n.s.
Bray P	0.64	n.s.	n.s.
Exchangeable K	0.75	n.s.	n.s.
Al_2O_3	n.s.	n.s.	n.s.
Fe_2O_3	n.s.	n.s.	n.s.
Explained variance (%)	41.9	24.2	18.3

n.s., Not significant.

1. AAAS, Proceedings on Human Population and Water: Fisheries, and Coastal Areas: Science and Policy Issues Symposium, American Association for the Advancement of Science, Washington DC, 1995.
2. Krishnamurthy, K., Choudhury, A. and Untawale, A. G., Status report: mangroves in India, Ministry of Environment and Forests, Govt of India, New Delhi, 1987.
3. Kathiresan, K., Quo vadis, mangroves? *Seshaiyana*, 1995, **3**, 25–28.
4. Dinesh, R., Suryanarayana, M. A., Ghoshal Chaudhuri, S. and Sheeja, T. E., Influence of leguminous cover crops on the general and specific biochemical characteristics in soils of a plantation. *Soil Till. Res.*, 2004, **77**, 69–77.
5. Bhattacharyya, P., Tripathy, S., Chakrabarti, K., Chakraborty, A. and Banik, P., Fractionation and bioavailability of metals and their impacts on microbial properties in sewage irrigated soil. *Chemosphere*, 2008, **72**, 543–550.
6. Vanhala, P. and Ahtiainen, J. H., Soil respiration, ATP content and photobacterium toxicity test as indicators of metal pollution in soil. *Environ. Toxicol. Water Qual.*, 1994, **9**, 115–121.
7. Yakovchenko, V. I., Sikora, L. J. and Kauffman, D. D., A biologically based indicator of soil quality. *Biol. Fertil. Soils*, 1996, **21**, 245–251.
8. Doran, J. W. and Parkin, T. B., Defining and assessing soil quality. In *Defining Soil Quality for a Sustainable Environment* (eds

- Doran, J. W. *et al.*), SSSA Publication No 35, Madison, WI, 1994, pp. 3–21.
9. Allen, S. E., *Chemical Analysis of Ecological Materials*, Blackwell, Oxford, UK, 1989.
 10. Gee, G. W. and Bauder, J. W., Particle-size analysis. In *Methods of Soil Analysis, Part 1, Physical and Mineralogical Methods* (ed. Klute, A.), Agronomy Monograph, ASA, SSSA, Madison, WI, 1986, 2nd edn, vol. 9, pp. 383–411.
 11. Gillman, G. P., A proposed method for the measurement of exchange properties of highly weathered soils. *Aust. J. Soil Res.*, 1979, **17**, 129–139.
 12. Thomas, G. W., Exchangeable cations. In *Methods of Soil Analysis, Part 2, Chemical and Microbiological Properties* (eds Page, A. L., Miller, R. H. and Keeney, D. R.), ASA, SSSA, Madison, WI, 1982, pp. 159–165.
 13. Olsen, S. R. and Sommers, L. E., Phosphorus. In *Methods of Soil Analysis, Part 2, Chemical and Microbiological Properties* (eds Page, A. L., Miller, R. H. and Keeney, D. R.), ASA, SSSA, Madison, WI, 1982, pp. 403–430.
 14. Barnhisel and Bertsch, P. M., Aluminium. In *Methods of Soil Analysis, Part 2, Chemical and Microbiological Properties* (eds Page, A. L., Miller, R. H. and Keeney, D. R.), ASA, SSSA, Madison, WI, 1982, pp. 275–300.
 15. Olson, R. V. and Ellis Jr, R., Iron. In *Methods of Soil Analysis, Part 2, Chemical and Microbiological Properties* (eds Page, A. L., Miller, R. H. and Keeney, D. R.), ASA, SSSA, Madison, WI, 1982, pp. 301–312.
 16. Vance, E. D., Brookes, P. C. and Jenkinson, D. S., An extraction method for measuring soil microbial biomass C. *Soil Biol. Biochem.*, 1987, **19**, 703–707.
 17. Wu, J., Joergensen, R. G., Pommerening, B., Chausod, R. and Brookes, P. C., Measurement of soil microbial biomass C by fumigation-extraction: an automated procedure. *Soil Biol. Biochem.*, 1990, **22**, 1167–1169.
 18. Joergensen, R. G., The fumigation-extraction method to estimate soil microbial biomass: extraction with 0.01 M CaCl₂. *Agric. Res.*, 1995, **48**, 319–324.
 19. Brookes, P. C., Landman, A., Pruden, G. and Jenkinson, D. S., Chloroform fumigation and the release of soil nitrogen; a rapid direct extraction method for measuring microbial biomass nitrogen in soil. *Soil Biol. Biochem.*, 1985, **17**, 837–842.
 20. Joergensen, R. G. and Mueller, T., The fumigation-extraction method to estimate soil microbial biomass; calibration of the k_{EN} value. *Soil Biol. Biochem.*, 1996, **28**, 33–37.
 21. Joergensen, R. G. and Brookes, P. C., Ninhydrin-reactive measurements of microbial biomass in 0.5 M K₂SO₄ soil extracts. *Soil Biol. Biochem.*, 1990, **22**, 1023–1027.
 22. Salamanca, E. F., Raubuch, M. and Joergensen, R. G., Relationships between soil microbial indices in secondary tropical forest soils. *Appl. Soil Ecol.*, 2002, **21**, 211–219.
 23. Casida Jr, L. E., Klein, D. A. and Santoro, R., Soil dehydrogenase activity. *Soil Sci.*, 1964, **98**, 371–378.
 24. Kandeler, E. and Gerber, H., Short-term assay of soil urease activity using colorimetric determination of ammonium. *Biol. Fertil. Soils*, 1988, **6**, 68–72.
 25. Gil-Sotres, F., Trasar-Cepeda, M. C., Ciardi, C., Ceccanti, B. and Leirós, M. C., Biochemical characterization of biological activity in very young mine soils. *Biol. Fertil. Soils*, 1992, **13**, 25–30.
 26. Tabatabai, M. A. and Bremner, J. M., Use of *p*-nitrophenyl phosphate for assay of soil phosphatase activity. *Soil Biol. Biochem.*, 1969, **1**, 301–307.
 27. Eivazi, F. and Tabatabai, M. A., Glucosidases and galactosidases in soils. *Soil Biol. Biochem.*, 1988, **20**, 601–606.
 28. Browman, M. G. and Tabatabai, M. A., Phosphodiesterase activity of soils. *Soil Sci. Soc. Am. J.*, 1978, **42**, 284–290.
 29. Tabatabai, M. A. and Bremner, J. M., Arylsulfatase activity of soils. *Soil Sci. Soc. Am. Proc.*, 1970, **34**, 225–229.
 30. Schinner, F. and von Mersi, W., Xylanase, CM-cellulase and invertase activity in soil: an improved method. *Soil Biol. Biochem.*, 1990, **22**, 511–515.
 31. Gomez, P. and Fortes, M. D., Transport of organic matter in the mangroves of Pagbilao, Quezon. Report on microbial aspects of nutrient cycling in mangrove environments, UNDP/UNESCO (RAS/79/002), Manila, Philippines, p. 24.
 32. Reyes, R. C. and Fortes, M. D., Litter production and leaf litter decomposition rates of mangrove trees in Pagbilao, Quezon. In Report on microbial aspects of nutrient cycling in mangrove environments, UNDP/UNESCO (RAS/79/002), Manila, Philippines.
 33. Dagar, J. C. and Sharma, A. K., Multiple viviparity in mangroves. *J. Andaman Sci. Assoc.*, 1989, **5**, 72–73.
 34. Trofymow, J. A., Detrital carbon fluxes and microbial activity in successional Douglas-fir forests. In Proceedings of a Workshop on Structure, Process and Diversity in Successional Forests of Coastal British Columbia (eds Trofymow, J. A. and MacKinnon, A.), Victoria, British Columbia, 1998, vol. 72, pp. 51–53.
 35. Kandeler, E., Stemmer, M. and Klimanek, E. M., Response of soil microbial biomass, urease and xylanase within particle size fractions to long-term soil management. *Soil Biol. Biochem.*, 1999, **31**, 261–273.
 36. Wick, B., Kühne, R. F., Vielhauer, K. and Vlek, P. L. G., Temporal variability of selected soil microbiological and biochemical indicators under different soil quality conditions in south-western Nigeria. *Biol. Fertil. Soils*, 2002, **35**, 155–167.
 37. Ocio, J. A. and Brookes, P. C., An evaluation of methods for measuring the soil microbial biomass in soils following recent additions of wheat straw and the characterization of the biomass that develops. *Soil Biol. Biochem.*, 1990, **22**, 685–694.
 38. Bolton Jr, H., Elliott, L. F., Papendick, R. I. and Bezdicek, D. F., Soil microbial and selected soil enzyme activities: effect of fertilization and cropping practices. *Soil Biol. Biochem.*, 1985, **17**, 297–302.
 39. Ghollara, M. and Raiesi, F., The adverse effects of soil salinization on the growth of *Trifolium alexandrinum* L. and associated microbial and biochemical properties in a soil from Iran. *Soil Biol. Biochem.*, 2007, **39**, 1699–1702.
 40. Yuan, B.-C., Li, Z.-Z., Liu, H., Gao, M. and Zhang, Y.-Y., Microbial biomass and activity in salt affected soils under arid conditions. *Appl. Soil Ecol.*, 2007, **35**, 319–328.
 41. Stemmer, M., Gerzabek, M. H. and Kandeler, E., Invertase and xylanase activity of bulk soil and particle-size fractions during maize-straw decomposition. *Soil Biol. Biochem.*, 1999, **31**, 9–18.
 42. Parham, J. A., Deng, S. P., Raun, W. R. and Johnson, G. V., Long-term cattle manure application in soil. I. Effect on soil phosphorus levels, microbial biomass C, and dehydrogenase and phosphatase activities. *Biol. Fertil. Soils*, 2002, **35**, 328–337.
 43. D'Ascoli, R. *et al.*, Impact of river overflowing on trace element contamination of volcanic soils in south Italy: Part II. Soil biological and biochemical properties in relation to trace element speciation. *Environ. Pollut.*, 2006, **144**, 317–326.
 44. Fischer, H., Sukhodolov, A., Wilczek, S. and Engelhardt, C., Effects of flow dynamics and sediment movement on microbial activity in a lowland river. *River Res. Appl.*, 2003, **19**, 473–482.
 45. Steenwerth, K. L., Jackson, L. E., Calderón, F. J., Stromberg, M. R. and Scow, K. M., Soil microbial community composition and land use history in cultivated and grassland ecosystems of coastal California. *Soil Biol. Biochem.*, 2002, **34**, 1599–1611.
 46. Dinesh, R., Ramanathan, G. and Harjit Singh, Influence of chloride and sulphate ions on soil enzymes. *J. Agron. Crop Sci.*, 1995, **175**, 129–133.

Received 27 March 2008; revised accepted 4 May 2009