

Increased xylanase activity in *Aspergillus niger* through mutation
Anjali Sharma*, A. K. Jaitly and Pankaj Rai*****

Author: Anjali Sharma, Research Scholar, Invertis University, Bareilly
*sharmanjali3012@gmail.com,

Corresponding authors:

1. Dr. A.K.Jaitly, Professor in Department of Plant Science, MJPRU, Bareilly
**jaitly1958@gmail.com,
2. Dr. Pankaj Rai, Assistant Professor in Department of Microbiology, Invertis University, Bareilly ***pankajraibhu@gmail.com

Increased xylanase activity in *Aspergillus niger* through mutation
Anjali Sharma*, A. K. Jaitly and Pankaj Rai*****

Microbiology Lab, Invertis University, Bareilly

Email; *sharmanjali3012@gmail.com, **jaitly1958@gmail.com,
***pankajraibhu@gmail.com

Abstract: *Aspergillus niger* is used for xylanase production on agricultural waste as substrate under broth culture. Rice straw and sugarcane baggase have been found the most potential substrate for xylanase production. Two different mutagens were used: UV radiation for different time duration and 5-Bromouracil of different concentrations. Mutants so formed were selected on the basis of morphological and colony characteristics. Selected mutants were checked for their stability, requirement of amino acid and xylanase activity. Tested mutants showed 4 fold increases in xylanase activity from wild type

Keywords: Agricultural waste; Rice straw, Sugarcane baggase, Mutation, Xylanase.

Introduction: Microbial enzymes are one of the fastest growing fields in biotechnology as these enzymes have several industrial applications [1]. Most common enzymes include cellulase, mannanase, arabinase, xylanase, etc. Hemi-cellulose is the second most abundant renewable biomass in nature after cellulose. Xylan is the major hemi-cellulose component and accounts for 20-35% of plant cell wall dry weight [2]. Xylan provides potential raw materials for obtaining fermentable sugars which can be converted into several valuable products. Xylanases are hydrolytic enzymes which cleave the β -1, 4 backbone of the hemi-cellulose. Xylanases and associated debranching enzymes, produced by a large variety of microorganisms including bacteria, actinomycetes, yeast and fungi [3].

Xylan consist of beta-1,4- linked D -xylopyranose as a backbone and short chains of o-acetyl, α -L-arabinofuranosyl, and α -D-glucuronyl residues. Enzymes such as endo-1,4-beta-xylanase, beta-D-xylosidase, beta-D-xylosidase, alpha-L-arabinofuranosidases, alpha-glucuronidases and acetyl esterases acts synergistically to hydrolyze xylan (Poutanen et.al, 1987). Pre-treatment with enzymes (mainly cellulase-free xylanase) has been reported as an environmentally benign and less expensive [4] method that aids in the spread of hardwoods and softwoods. [5].

Enzymatic conversion of hemi-cellulose produces high value products like furfural, xylitol, bio fuels and in artificial low calorie sweeteners [6]. Breakdown of hemicelluloses in wheat flour making the dough softer and easier to knead [7]. Micro-organisms such as bacteria and fungi producing xylanases are more favoured as they are easily available and structurally stable as well as can also be easily genetically manipulated. It has been reported that fungi are more efficient in producing xylanase in comparison to bacteria, however some species of *Bacillus* has been reported to secrete important extracellular enzymes and protein in the medium.

Mutation plays an important role in improving fungal strains for specific work. Mutation may take place spontaneously or may be induced. Spontaneous mutation is slow and their rate depends on the growth conditions of organism and the mutation frequency is low (8). Induced mutation could be carried either by physical or chemical methods. Physical mutation could be carried by UV radiation, gamma radiation or X-rays (9) while chemical mutation by using EtBr (Ethidium bromide), 5-Bromouracil, EMS (Ethyl Methyl Sulphonate) etc. Xylanase activity of *Aspergillus niger* has been increased up to 118% by mutation. UV radiation has been reported as very effective in industries and do not require any equipment. As a result, the current study aims to increase xylanase activity in *Aspergillus niger* by mutation in order to investigate its industrial potential (10). As it has been reported that xylanase is produced by different *Aspergillus* species such as *Aspergillus nidulans*, *A. niger* and *A. oryzae*. Different strains of *Aspergillus niger* have been reported for the production of Xylanase through mutagenesis. Mutagenesis involves two different methods which includes physical as well chemical; in physical method fungus is treated with UV radiations for different time interval and in chemical method fungus is grown on the medium containing chemical mutagen such as 5-Bromouracil , Ethylmethyl Sulphonate etc. In the commercial development of microbial fermentation processes, industrial strain enhancement is critical. According to Rowlands (1984) (11), mutagenic methods can be tweaked in terms of mutagens and dose. Effects can

be taken into account, and mutagenesis can be intensified or guided to produce the highest frequency of suitable mutant types among the isolates to be tested. Most of the research on increasing xylanase activity by mutation and protoplast fusion have been conducted on mesophilic and thermophilic strains acquired from composting materials, agricultural wastes and natural environments.

Materials and methods:

Organism: *Aspergillus niger*, isolated in the Microbiology laboratory of Plant Science department has been used for xylanase production.

Culture medium used: Minimal medium was used with the following composition - MgSO₄.4H₂O (0.25gm); K₂HPO₄ (0.50gm); Starch (7.5gm); Agar (10.250gm) in 500ml distilled water. Complete medium was also used in this study, the composition was same as minimal medium excepting it also includes all amino acids.

Chemical and physical mutagenesis: 5-Bromouracil, a base analogue and UV radiation have been used as mutagen. Different concentrations of 5- Bromouracil i.e., 200µg, 400µg, 600µg and 800µg were used to treat the freshly grown mycelium/ spores of *Aspergillus niger*. 5 ml of spore suspension was treated by UV radiation for different time intervals from 30-180 min with a gap of 30 min each. For mutant selection, 1 ml of spore suspension was plated with both complete and minimum medium after treatment. Mutants differ from wild type in their morphology, colony pattern, diameter, colour, size of somatic and reproductive structures were selected and tested for requirement of specific amino acids for the growth and xylanase activity. Stability of the mutants was checked by successive culturing.

Enzyme preparation and xylanase Activity: 25 ml YpSs broth medium supplemented with 1 gm of rice straw powder was used for the production of xylanase. Above medium was inoculated with the test fungus and incubated at 30°C and pH 6.0 for 5 days at constant temperature incubators. Triplicates were taken for each strain of *Aspergillus niger* wild type and mutants. After incubation culture was centrifuged at 5000rpm for 5 minutes and supernatant was used as enzyme to evaluate xylanase activity by DNS method (12) (13). The enzyme activity was evaluated using following formula:

Enzyme activity = OD X 1/ enzyme volume X 1/ substrate volume X 1/ Incubation time X Retention coefficient

Substrate: 1% xylan in acetate phosphate buffer with pH 6.0 was used as a substrate.

Standard graph: Standard curve was prepared using stock solution of 1 mg/ml of Xylose. One ml of different concentrations from 0µg-200µg of Xylose prepared from stock solution in different test tubes and boiled after adding 3ml of DNS reagent. OD was taken at 540 nm against spectro-zero after cooling it.

Spectro-zero: 0.9 ml of substrate and 0.1 ml of buffer was incubated at 50 °C for 30 min in a test tube. 3 ml of DNS reagent was added and boiled for 5 min and was taken to set zero at 540nm.

Reaction: 0.2 ml enzyme and 1.8 ml substrates was taken in a test tube and kept at 50° C for 30 min in test tube. After reaction 1 ml of the reaction mixture was taken in a test tube, 3 ml of DNS reagent was added and boiled for 5 min. After, cooling OD was taken at 540 nm against spectro-zero.

Results: Mutants formed by physical and chemical mutation were selected and named according to the source of mutagens. Mutants differed significantly from wild type in terms of morphology, colony layout, diameter, and colour (Table 1).

Table 1: Morphological difference *Aspergillus niger* wild and mutants

S.no	Isolates	Characteristics
1.	Wild	Dark black, dense colony , outer colony black
2.	Mutants AnB-1	Light black, sparse colony, outer whitish black
3.	AnB-2	Light black, sparse colony, outer whitish black
4.	AnB-3	Light black, sparse colony, outer whitish black
5.	AnU-1	Black , scattered colony, outer creamish white colony
6.	AnU-2	Black , scattered colony, outer creamish white colony

Mutants obtained differ in the requirement of amino acid (Table 2) and has been characterized as auxotrophic mutants, which will be used for further improvement in xylanase activity through protoplast fusion.

Table 2: Growth of mutants on minimal medium in presence/absence of different amino acids

S.no	Amino acid	Wild	AnB-1	AnB-2	AnB-3	AnU-1	AnU-2
1.	Cysteine	+	+	+	+	+	+
2.	Arginine	+	+	-	-	-	+
3.	Lysine	+	-	+	+	+	-
4.	Valine	+	-	+	+	+	+
5.	Glycine	+	+	-	-	-	-
6.	Proline	+	+	-	+	+	+
7.	Serine	+	+	+	-	+	+
8.	Histidine	+	-	+	+	-	+
9.	Methionine	+	+	-	+	-	-

10.	Tryptophan	+	+	+	+	+	+
-----	------------	---	---	---	---	---	---

(+) Growth with presence of amino acid; (-) No growth with absence of amino acid

Xylanase activity of the wild type and mutants has been undertaken and results have been presented in Table 3. All the tested mutants showed an increase in xylanase activity over wild type; however mutant AnB-3 showed the highest activity of 1289.18 IU/ml followed by AnU-1 (1182.79 IU/ml) on day 5, showed an increase of 86.43% and 75.05% over wild type.

Table 3: Xylanase activity of *Aspergillus niger* wild and mutant strain

S.no	Isolates	Day	Source	Enzyme activity (IU/ml)	% Increase
1.	Wild	5	-	691.48±0.76	-
2.	AnB-1	5	Chemical	1163.04±0.52	68.19%
3.	AnB-2	5	Chemical	1091.87±0.67	57.88%
4.	AnB-3	5	Chemical	1289.18±0.88	86.43%
5.	AnU-1	5	Physical	1182.79±0.31	71.05%
6.	AnU-2	5	Physical	998.37±0.84	44.38%

Discussion: Mutants of *Aspergillus niger* differ in their culture characteristics and as well as in requirement of amino-acid, they were regarded as auxotrophic mutants. Many workers have also produced auxotrophic mutants in same procedure. (14)(15)(16). Mutants selected showed an increase in xylanase activity (Table 3). Amongst the all mutants, mutant AnB-3 and AnU-1 exhibited 86.43% and 71.05% increase in enzyme activity. It may be due to improved change in gene sequence which resulted in the formation of more active protein other workers also found similar results. However, more work is needed to look in the industrial applications of this xylanase. Further work to improve xylanase activity by fusing these auxotrophic strains using protoplast fusion is under process (17) (18)

Conclusion: : *Aspergillus niger* wild and mutants formed during present study have been proved to be more effective in utilization of rice straw waste, thus shall be helpful in minimizing pollution caused by burning of rice straw. Mutants formed very more effective in utilization of rice straw than the wild *Aspergillus niger*. The aim of this study was to use agricultural waste for the production

Acknowledgement: The authors are grateful to the *Council of Science and Technology, U.P.*, for providing financial assistance for above work.




Fig.1 (a) *Aspergillus niger* wild, (b) *Aspergillus niger* after treatment with physical mutant i.e UV radiation and (c) *Aspergillus niger* after treatment with chemical mutant i.e, 5-Bromouracil.

References:

1. Pathak, S.S., Sandhu, S.S. and Rajak, R.C. Mutation studies on fungal Glucoamylase: A Review- *IJPBS* 2005, 5, 297-308.
2. Polizeli, M.L.T.M et.al., Xylanases from fungi: properties and industrial applications – *Applied Microbiology and Biotechnology* 2005, 67, 577-591.
3. Solimon, H.M, Sherief A.D.A and Tanash.A.B.EL - Production of Xylanase by *Aspergillus niger* and *Trichoderma viride* using some Agricultural residues- *International Journal of Agricultural Research* 2012, 7(1), 46-57.

4. Bajpai, P Application of enzymes in the pulp and paper industry. *Biotechnol Prog* 1999, 15, 147–157.
5. Angayarkanni, J., Palaniswamy, M., Pradeep, B.V., Swaminathan, K. Biochemical substitution of fungal xylanases for prebleaching of hardwood kraft pulp. *Afr J Biotechnol* 2006, 5, 9291–9296.
6. Polizeli, M.L.T.M., Rizzatti, A.C.S., Terenzi, H.F., Jorge, J.A. and Amorim, D.S. Xylanases from fungi: properties and industrial applications. *Appl Microbiol Biotechnol*, 2005, 67, 577–591.
7. Betini, J.H.A., Michelin, M., Peixoto-Nogueira, S.C., Jorge, J.A., Terenzi, H.F., and Polizeli, M.L.T.M.. Xylanases from *Aspergillus niger*, *Aspergillus niveus* and *Aspergillus ochraceus* produced under solid-state fermentation and their application in cellulose pulp bleaching; *Bioprocess and Biosystems Engineering*, 2009, 32, 819–824.
8. Waites M.J., Morgan N.L., John S.R. and Higton G. Industrial Microbiology- An Introduction. *Wiley Blackwell Publishing*, 2002, 1,
9. Agrawal R., Deepika N.U. and Joseph R. Strain improvement of *Aspergillus sp.* and *Penicillium sp.* by induced mutation for biotransformation of alpha penine to verbinol. *Biotech Bioeng*, 1999, 63, 249-252.
10. Desai, D.I. and Iyer, B.D. Utilization of Corn Cob Waste for Cellulase-Free Xylanase Production by *Aspergillus niger* DX-23: Medium Optimization and Strain Improvement. *Waste and Biomass Valorization* ,2017, 8, 103–113.
11. Rowlands, R.T. Industrial strain improvement: Mutagenesis and random screening procedure. *Enzy. Microb. Tech.* 1984, 6, 3-10.
12. Miller. Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar. *American Chemical Society Anal. Chem.* 1959, 31, 3, 426–428.
13. Khan A , Tremblay D, Le Duy (1986). Assay of xylanase and xylosidase activities in bacterial and fungal cultures, *Enzyme Microb. Technol*, 8, 373-377.
14. Romanowska, I., Polak, J. and Bielecki S. Isolation and properties of *Aspergillus niger* IBT-90 xylanase for bakery. *Appl Microbiol Biotechnol* , 2006, 69, 665-671.
15. Kango, N., Agrawal, S.C. and Jain. P.C. Xylanase Production by thermophilic fungi from Soil and Decomposing organic matter. *Microbiology and Biotechnology for Sustainable Development* , 2004, B-13, 293-299
16. Radha, S., Himakiran Babu, R., Sridevi, A., Prasad, N.B.L. and Narasimha, G. Development of mutant fungal strains of *Aspergillus niger* for enhanced production of acid protease in submerged and solid state fermentation. *European Journal of Experimental Biology*, 2012, 2(5) , 1517-1528.
17. Kirimura, K., Yaguchi, T. and Usami, S. Intraspecific Protoplast Fusion of Citric acid producing strains of *Aspergillus niger*. *J. Ferment. Technol.* 1986, 6, 473-479.
18. Elbouami, F., Isolation and regeneration of protoplasts from mycelium of *Fusarium solani*. *African Crop Science Journal*, 2001, 9(2), 351-358.