

Spatio-temporal analysis of drought and aridity in Gomti basin

R Sarma¹ and D K Singh²

¹Senior Research Fellow, Centre for Environment Science and Climate Resilient Agriculture, Indian Agricultural Research Institute, New Delhi- 110012 (email: rikisarmah@gmail.com)

²Principal Scientist, Water Technology Centre, Indian Agricultural Research Institute, New Delhi-110012.

Running title: Drought and aridity in Gomti basin

ABSTRACT

This study analyses the drought events for the Gomti basin in Uttar Pradesh, India using the Standardized Precipitation Index (SPI) at the spatial and temporal scales. Daily precipitation data for 14 districts in the Gomti basin for 41 years (1971-2011) was used to calculate the SPI for 1-month, 4-month, 6-month and 12-month time scales. Results for the 6-month (June-November) and 12-month time scales were similar and drought years were observed in 1972, 1979, 1987, 1993, 1994, 2002 and 2010. The 4-month SPI was analyzed for the main monsoon months in the Gomti basin, i.e. from June to September. Results showed that significant drought occurred during the monsoon months of 1979, 1987, 1993, 2002, 2009 and 2010. For drought analysis at the spatial scale, the Kriging interpolation method available in ArcMap was used. The 12-month SPI showed that the frequency of severe and extreme drought were more in the upper regions of the basin during 1971-2000 while drought frequency was more in the central and lower regions of the basin during 2001-2011. Further, the de Martonne aridity index was calculated for the period 1971-2007 and its correlation with the 1-month SPI for the period 1971-2007 was evaluated.

Key words: Spatial and temporal drought analysis, Standardized Precipitation Index, Kriging interpolation, de Martonne aridity index, Gomti basin.

Agriculture is one of the main sources of livelihood in Uttar Pradesh, contributing a large share to the state's output¹. Farming operations depend heavily on rainfall and when precipitation amounts are inadequate, irrigation plays a crucial role in determining crop yields. The Gomti river basin that lies in Uttar Pradesh has a predominantly agrarian economy with 80% of the crops being grown under full or supplementary irrigation². Drought and aridity are very important parameters that assist in estimating the irrigation demand.

Drought is a natural hazard that has significant negative impacts on the environment, agriculture and socio-economic conditions of the people. Drought is the manifestation of long

spell of deficit in precipitation. Depending on the approach used to measure drought, it can be classified as – meteorological drought, hydrological drought, agricultural drought and socio-economic drought³. The India Meteorological Department declares a drought year for the country based on the deficit of rainfall. When the rainfall deficiency is more than 10% in 20 to 40% of the country, then IMD terms that year as All India Drought Year (now termed “Deficient Year”). When the spatial coverage of drought is more than 40% then IMD declares All India Severe Drought Year (now termed “Large Deficient Year”)^{4,5}. A number of drought indices have been developed and used by scientists around the world⁶. Of these, the Standardized Precipitation Index (SPI) is the most commonly used to quantify precipitation deficit. McKee *et al.*⁷ have developed the SPI for the classification of drought in a region. This classification has been accepted internationally to define meteorological drought. The World Meteorological Organization uses the same criteria to define drought. The SPI can be determined using rainfall data alone at variable timescales (1-month, 3-month, 6-month, 9-month, 12-month and 24-month).

The Standardized Precipitation Index makes drought assessment easier and has a wide range of meteorological, hydrological and agricultural applications⁸. Mishra and Desai⁹ studied the frequency of drought occurrence in the Kansabati river basin using the SPI and reported severe drought during the 1980s. Frequency of droughts for the period of 1951-2007 in the Hunan Province in China was studied by Du *et al.*¹⁰ using the SPI at various time scales. They reported that during that period, more dry years were experienced in the upper reaches of the major rivers in the Hunan Province than the lower and middle reaches. Santos *et al.*¹¹ used the SPI at 1, 3, 6 and 12 month time scales to characterize drought events in mainland Portugal from September 1910 to October 2004. Pai *et al.*¹² compared two drought indices – the Per cent of Normal Precipitation (PNP) and SPI for 458 districts in India covered by southwest monsoon. Their results showed that the SPI is a better drought index than PNP.

Ntale and Gan¹³ compared the SPI with the Palmer Drought Severity Index and the Bhalme-Mooley Index for East Africa and concluded that the SPI is more suitable for analyzing East African droughts since it is more easily adapted to the local climate and only requires precipitation data. Potop *et al.*¹⁴ used the SPI along with the percentage of long term precipitations and the Ped Drought Index for the period from 1961-2007 for the Czech Republic to evaluate drought episodes and their potential impacts on winter and spring cereal crops.

Aridity has different definitions across different disciplines such as climatology, meteorology, hydrology and agriculture^{15,16}. Penck¹⁷ first defined an arid region as a place where annual evaporation exceeds precipitation. Beran and Rodier¹⁸ defined aridity of a given region as the deficiency of available water in an ecosystem. Among the numerous aridity indices developed for the determination of irrigation demands¹⁹, the de Martonne aridity index is one of the most commonly used since it only requires mean monthly temperature and mean monthly rainfall data for a period. Zhang *et al.*²⁰ used the SPI and the de Martonne aridity index for the Pearl River basin for the period from 1960-2005 to classify anomalously wet and dry conditions. Shahid²¹ studied the spatio-temporal variation in aridity and dry period in terms of irrigation demand for the 1959-2009 time period for Bangladesh using the de Martonne aridity index. His study revealed that dryness in Bangladesh has decreased in the last fifty years and irrigation was required for more than six months in western Bangladesh. Livada and Assimakopoulos²² compared the SPI and de Martonne aridity index for 51 years for Greece and found a satisfactory correlation between the two indices. SPI and the de Martonne aridity index were also used to observe variability in drought and aridity in the Barlad catchment in Romania and its potential impacts on hydrology²³. De Martonne's aridity index was used along with five other indices by Lungu *et*

*al.*²⁴ for Dobrudja territory to demonstrate that increasing aridity in the region is a real hazard.

From the above, it is clear that analysis of spatial and temporal trends of drought and aridity are important to understand the climatic variability and irrigation requirements in a region. In this paper the Standardized Precipitation Index and the de Martonne aridity index were determined for Gomti basin to describe the spatial and temporal variability in drought severity and aridity and irrigation requirements respectively. Additionally, a correlation between the two indices was evaluated.

Data and methods

The Gomti river basin is part of the Indo-Gangetic plains of India (Figure 1). The Gomti basin comprises of 31240 km² area in the state of Uttar Pradesh in India²⁵. The climate of the Gomti basin ranges from semi-arid to sub-humid tropical. The average annual rainfall in the basin varies from 850 to 1100 mm. Approximately 75% of the total rainfall occurs in the monsoon season (middle of June to middle of October)²⁶. Districts in the Gomti basin are Bara Banki, Faizabad, Hardoi, Jaunpur, Kheri, Lucknow, Pilibhit, Pratapgarh, Rae Bareli, Shahjahanpur, Sitapur, Sultanpur, Unnao and Varanasi.

Data used

Daily meteorological data consisting of minimum and maximum temperature for the period of 1971-2007 and daily rainfall data for the period of 1971-2011 for each district of Gomti basin available with NICRA (National Innovations on Climate Resilient Agriculture, <http://www.nicra-icar.in/nicrarevised/>) web portal were used. Monthly and annual means were then computed from the daily data for each district of the Gomti basin.

Methods of analysis The Standardized precipitation index was used to assess the drought frequency at different time scales. The de Martonne aridity index was used to assess the

episodes of dryness and irrigation demand. Descriptions of these are presented in the following paragraphs.

Standardized Precipitation Index (SPI)

Precipitation deficit affects soil moisture, surface and subsurface flow, reservoir storage and groundwater recharge. Long term deficit in precipitation may result in drying of streams, reservoirs, aquifers and severe soil moisture deficit. Quantification of precipitation deficit at different time scales is required to study its influence on soil moisture, surface and subsurface flow, reservoir storage and groundwater recharge. The Standardized Precipitation Index was developed by McKee *et al.*⁷ to quantify the precipitation deficits at different time scales. According to McKee *et al.*⁷ drought occurs when the value of SPI is continuously negative and reaches -1.0 or less. They defined the criteria for drought intensity based on the SPI (Table 1). Drought ends when SPI becomes positive. SPI is computed from Equation 1 and 2.

$$SPI = S \frac{t - (c_2 t + c_1) + c_0}{((d_3 t + d_2) + d_1) t + 1}, \quad t = \sqrt{\ln \frac{1}{H(x)^2}} \quad \dots\dots\dots (1)$$

$$G(x) = \frac{1}{\beta^\gamma \Gamma(\gamma)} \int_0^x x^{\gamma-1} e^{-x/\beta} dx, \quad x > 0, \Gamma(\gamma) = \int_0^\infty x^{\gamma-1} e^{-x} dx \quad \dots\dots\dots (2)$$

where, x: rainfall values, β, γ : scale parameter and shape parameter of the Γ function, S: positive and negative coefficients, c_0 : 2.515517, c_1 : 0.802853, c_2 : 0.010328, d_1 : 1.432788, d_2 : 0.189269, d_3 : 0.001308, $G(x)$: probability of precipitation distribution. $S = 1$ when $G(x) > 0.5, H(x) = 1 - G(x)$ and $S = -1$ when $G(x) \leq 0.5, H(x) = G(x)$.

Aridity index of E. de Martonne

Martonne E. de²⁷ introduced an aridity index based on temperature and precipitation only. The monthly values of the index are calculated as:

$$I_i = \frac{12P_i}{T_i + 10} \quad \dots\dots\dots (3)$$

where, P_i : monthly precipitation amount in mm, T_i : monthly mean air temperature in °C. Since irrigation requirement depends both on temperature and rainfall, this index is used to

identify the months when irrigation is necessary. This index may also be used to identify dry episodes in a region. According to the index, if $I_i < 30$, dry conditions may be observed and irrigation may be necessary. In general, irrigation becomes necessary when $I_i < 20$.

Results and discussion

Standardized Precipitation Index (SPI)

The 12-month and the 6-month SPI (June-November) were calculated for each of the 14 districts of the Gomti basin. A dry year occurs if $SPI \leq -1.00$ (Table 1). During the period 1971-2011, at the basin level, drought years were observed for 7 years based on both the 12-month and 6-month SPI, i.e. in 1972, 1979, 1987, 1993, 1994, 2002 and 2010 (Figure 2 (a) & (b)). The drought conditions were mapped using the Kriging interpolation technique in ArcMap 10.0. At the spatial scale, based on the 12-month SPI, the frequency of severe and extreme drought were observed more for the upper districts of the basin from 1971 to 2000. During the early 2000s, the central region experienced more severe and extreme drought while the lower districts of the basin experienced more severe and extreme drought in 2010. Figure 3 demonstrates this drought frequency for (a) upper region of the basin (districts Pilibhit, Shahjahanpur, Kheri, Sitapur, Hardoi), (b) central region (Bara Banki, Lucknow, Unnao, Faizabad, Pratapgarh, Sultanpur, Rae Bareli) and (c) lower region (Jaunpur, Varanasi).

The 4-month SPI for the monsoon months in Gomti basin (June-September) was calculated and drought frequency was observed more during the 2000s. Significant drought years were observed in 1979, 1987, 1993, 2002, 2009 and 2010. Kriging interpolation for these years is presented in Figure 4.

De Martonne aridity index (I_i)

The aridity index developed by de Martonne identifies dry period in terms of irrigation demand, i.e. the months when irrigation is necessary. For the Gomti basin, irrigation was

necessary for more than 8 months ($I_i < 20$) for all districts of the basin for the period 1971-2007. During this period, it was observed that for all the districts, the value of the index has decreased on a monthly basis. This decrease still kept the index above 20 for some months but lowering of its value implies that rainfall in some districts of the basin is decreasing over the last 37 years. At the spatial scale, aridity index was lower for the districts in the lower region of the basin while the upper and central regions had higher values.

The comparison of the 1-month SPI and the de Martonne aridity index I ($I = I_i/12$) showed that both indices were positively correlated (correlation coefficient ranging from 0.30 to 0.38 for all 14 districts). Although the correlation is linear, the indices are closer together for lower values and more scattered as the values increase (Figure 5).

Discussion

Analysis of the 6-month SPI (June-November) and the 12-month SPI for Gomti basin yielded similar results. This indicates that the drought period in any particular year in the basin is primarily governed by the rainfall pattern from June to November. Analysis of rainfall trends using the Mann-Kendall test by Abeyasingha *et al.*²⁸ showed that 1987 and 2002 experienced negative rainfall anomaly/ drought which is also reported in the present study.

June-September are the main monsoon months in the basin and analysis of drought during these months is important from an agricultural perspective. Jha *et al.*²⁹ reported an increasing trend in dryness during the summer monsoon season in their study for the period 1951-2006 for the Gangetic plain. In the present study, based on the 4-month SPI, more drought years were observed during the 2000s although severity of the drought was more in 1979, 1987 and 1993.

The de Martonne index for the basin showed that the lower region of the basin experienced more dry episodes and aridity for the period 1971-2007. This result is consistent with the

findings of Abeysingha *et al.*²⁸ which stated that rainfall has decreasing trend in the downstream areas of Jaunpur and Varanasi.

Finally, correlation between the 1-month SPI and the de Martonne aridity index (I) was found to be positive. This correlation is stronger for lower values of SPI and I indicating that during a drought condition, irrigation demand naturally increases. For higher values of SPI, i.e. during wet conditions, irrigation demand automatically decreases. Since the de Martonne aridity index is dependent on both temperature and rainfall, the scattered higher values of I may be attributed to increasing trend of mean temperature but both increasing and decreasing trend of rainfall for different districts in the basin^{28,30}.

The Gomti basin experienced drought in 1972, 1979, 1987, 1993, 1994, 2002 and 2010. Of these, only 1972, 1979, 1987 and 2002 were declared as major drought years in the country^{31,32}. This suggests that drought at the regional scale and at the country level may not always occur in the same year. At the regional scale, there is poor spatial and temporal distribution of rainfall. There may also be local climate variations which may affect the rainfall distribution. These could be the possible reasons for drought occurrence in the basin during 1993, 1994 and 2010, even when drought was not observed in the entire country. Since changes in temperature and precipitation are not globally uniform and considerable spatial and temporal variations in climate parameters exist at the regional level^{33,34}, it is important to analyse historical climate data at the regional level also. Such studies at regional scales are very helpful in defining historical drought years for smaller regions and for assessment of future irrigation requirements.

Conclusions

The spatial and temporal variability of drought and aridity was studied for the Gomti basin in Uttar Pradesh. Significant drought years were observed in 1972, 1979, 1987, 1993, 1994, 2002 and 2010. The Kriging interpolation method showed that during 1971-2000, the upper

regions of the basin experienced more severe and extreme drought events. The central and lower regions experienced more drought during 2001-2011. Based on the de Martonne aridity index, it was found that irrigation in the basin was necessary for more than 8 months in a year. During the period 1971-2007, aridity was more for the lower regions of the basin. The 1-month SPI and the de Martonne aridity index showed a positive correlation. Understanding drought intensity and severity plays a vital role in agriculture, particularly in estimating irrigation requirements. It is suggested that such studies may be undertaken for other basins in India.

References

1. Roy, R. and Ahmad, H., State Agricultural Profile of Uttar Pradesh (2011-12). Agro-Economic Research Centre, University of Allahabad, 2013.
2. Abeysingha, N. S., Singh, M., Sehgal, V. K., Khanna, M., Pathak, H., Jayakody, P. and Srinivasan, R., Assessment of water yield and evapotranspiration over 1985 to 2010 in the Gomti River basin in India using the SWAT Model. *Curr. Sci.*, 2015, **108**(12), 2202-2212.
3. Wilhite, D. A. and Glantz, M. H., Understanding the Drought Phenomenon: The Role of Definitions. *Water Int.*, 1985, **10**(3), 111-120.
4. India Meteorological Department, Forecaster's Guide. India Meteorological Department, Pune, 2008.
5. Koshy, J. and Vasudeva, V., No more droughts in India, says IMD. *The Hindu*, January 12, 2016. Retrieved from <https://www.thehindu.com/>. Accessed on 30th October 2018.
6. World Meteorological Organization, Standardized Precipitation Index User Guide (M. Svoboda, M. Hayes and D. Wood). WMO-No. 1090, Geneva, 2012.
7. McKee, T. B., Doesken, N. J. and Kliest, J., The relationship of drought frequency and duration to time scales, in Proceedings of the 8th Conference on Applied Climatology, 17-22 January, Anaheim, CA. American Meteorological Society: Boston, MA, USA, 1993, pp. 179-184.
8. Hayes, M., Wilhite, D. A., Svoboda, M. and Vanyarkho, O., Monitoring the 1996 drought using the Standardized Precipitation Index. *B. Am. Meteorol. Soc.*, 1999, **80**, 429-438.
9. Mishra, A. K. and Desai, V. R., Spatial and temporal drought analysis in the Kansabati river basin, India. *Int. J. River Basin Management*, 2005, **3**(1), 31-41.
10. Du, J., Fang, J., Xu, W. and Shi, P., Analysis of dry/wet conditions using the standardized precipitation index and its potential usefulness for drought/flood monitoring in Hunan Province, China. *Stoch. Env. Res. Risk A.*, 2013, **27**(2), 377-387.
11. Santos, J. P., Portela, M. M. and Pulido-Calvo, I., Regional frequency analysis of droughts in Portugal. *Water Resour. Manage.*, 2011, **25**, 3537-3558.
12. Pai, D. S., Sridhar, L., Guhathakurta, P. and Hatwar, H. R., District-wide drought climatology of the southwest monsoon season over India based on standardized precipitation index (SPI). *Nat. Hazards*, 2011, **59**(3), 1797-1813.
13. Ntale, H. K. and Gan, T. Y., Drought indices and their application to East Africa. *Int. J. Climatol.*, 2003, **23**, 1335-1357.

14. Potop, V., Turkott, L., Koznarova, V. and Mozny, M., Drought episodes in the Czech Republic and their potential effects in agriculture. *Theor. Appl. Climatol.*, 2010, **99**, 373-388.
15. Quan, C., Han, S., Utescher, T., Zhang, C. and Liu, Y., Validation of temperature-precipitation based aridity index: Paleoclimatic implications. *Palaeogeogr., Palaeoclimatol. Palaeoecol.*, 2013, **386**, 86-95.
16. Maliva, R. and Missimer, T., Arid Lands Water Evaluation and Management. *Environmental Science and Engineering*. Springer, 2012, pp. 21-39, DOI: 10.1007/978-3-642-29104-3_2.
17. Penck, A. Versuch einer Klimaklassifikation auf physiogeographischer Grundlage. *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, 1910, **12**, 236-246.
18. Beran, M. A. and Rodier, J. A., Hydrological aspects of drought, studies and reports in hydrology. UNESCO-WMO, 39. Presses Universitaires de France, Paris, 1985.
19. World Meteorological Organization, Drought and agriculture. WMO Note 138 Publ WMO-392 Geneva- Switzerland, 127, 1975.
20. Zhang, Q., Xu, C. and Zhang, Z., Observed changes of drought/wetness episodes in the Pearl River basin, China, using the standardized precipitation index and aridity index. *Theor. Appl. Climatol.*, 2009, **98**, 89-99.
21. Shahid, S., Spatio-temporal variation of aridity and dry period in term of irrigation demand in Bangladesh. *Am. Eurasian J. Agric. Environ. Sci.*, 2010, **7**(4), 386-396.
22. Livada, I. and Assimakopoulos, V. D., Spatial and temporal analysis of drought in Greece using the Standardized Precipitation Index (SPI). *Theor. Appl. Climatol.*, 2007, **89**, 143-153.
23. Borcan, M., Cheval, S. and Chendes, V., Observed variability of drought and aridity and its impact on the hydrological regime in the Barlad catchment (Romania). *Geophys. Res. Abstr.*, 2015, **17**.
24. Lungu, M., Panaitescu, L. and Nita, S., Aridity, climatic risk phenomenon in Dobrudja. *Present Environment and Sustainable Development*, 2011, **5**(1), 179-189.
25. Mali, S., Crop water footprint assessment for sustainable agriculture in selected sub-basins of River Ganga, PhD thesis, Post-Graduate School, Indian Agricultural Research Institute, New Delhi, India, 2015.
26. Rai, R. K., Upadhyay, A., Sarkar, S., Upadhyay, A. M. and Singh, V. P., GIUH based transfer function for Gomti River basin of India. *J. Spat. Hydrol.*, 2009, **9**, 24-50.
27. de Martonne E., Une nouvelle fonction climatologique: l'indice d' aridité. *La Météorologie*, 1926, **2**, 449-458.
28. Abeysingha, N. S., Singh, M., Sehgal, V. K., Khanna, M. and Pathak, H., Analysis of Rainfall and Temperature Trends in Gomti river basin. *J. Agric. Phys.*, 2014, **14**(1), 56-66.
29. Jha, S., Sehgal, V. K., Raghava, R. and Sinha, M., Trend of standardized precipitation index during Indian summer monsoon season in agroclimatic zones of India. *Earth Syst. Dynam. Discuss.*, 2013, **4**, 429-449.
30. Hingane, L. S., Rupa, Kumar K., and Murty, V. R., Long-term trends of surface air temperature in India. *Int. J. Climatol.*, 1985, **5**, 521-528.
31. Indian Institute of Tropical Meteorology, Pune. Monsoon On-Line. <https://www.tropmet.res.in/~kolli/MOL/Monsoon/Historical/air.html>. Accessed on 30th October, 2018.
32. Jamwal, N., India's criteria for classifying drought-hit regions are causing many cases to go unreported. January 27, 2018. <https://scroll.in/article/866262/indias-criteria-for-classifying-drought-hit-regions-are-causing-many-cases-to-go-unreported>. Accessed on 30th October, 2018.

33. Yue, S. and Hashino, M., Long term trends of annual and monthly precipitation in Japan. *J. Am. Water Resour. As.*, 2003, **39**(3), 587-596.
34. Sarma, R. and Singh, D. K., Homogeneity analysis of climatic parameters for the Gomti river basin. *Indian J. Soil Conserv.*, 2018, **46**(1), 25-33.

List of tables and figures

Table 1. Dryness/wetness categories based on SPI values^{6,7}

Figure 1. Location of the Gomti river basin in India²⁵

Figure 2. Drought years observed based on (a) 12-month SPI (b) 6-month SPI

Figure 3. Drought frequencies for (a) upper region (b) central region and (c) lower region of Gomti basin

Figure 4. June-September 4-month SPI for Gomti basin

Figure 5. Scatter plots and correlation lines between de Martonne aridity index and SPI for all districts of Gomti basin

Table 1. Dryness/wetness categories based on SPI values^{6,7}

SPI	Category
≥ 2.0	Extremely wet
1.50 to 1.99	Severely wet
1.00 to 1.49	Moderately wet
0.99 to -0.99	Near normal
-1.00 to -1.49	Moderately dry
-1.50 to -1.99	Severely dry
≤ -2.0	Extremely dry




Figure 1. Location of the Gomti river basin in India²⁵







Figure 2. Drought years observed based on (a) 12-month SPI (b) 6-month SPI






(a)



(b)



(c)

Figure 3. Drought frequencies for (a) upper region, (b) central region and (c) lower region of Gomti basin















Figure 4. June-September 4-month SPI for Gomti basin








Figure 5. Scatter plots and correlation lines between de Martonne aridity index and SPI for all districts of Gomti basin